

Ambys Lab

海人 Umihito アクアリウム

SciMoun


The Axolotl News Letters

～Axolotl Research Reports and Updates～

Vol 8.0

Feature 1: Cognition and Behavior in Axolotls
— Presented at an International Conference!

Feature 2: Secrets of Longevity Learned from the Olm!



2026/4/19 初版

Presented by
HAL lab_Axolotl (since2020～)

Commercial use of all images and text without permission is prohibited. This work is licensed under a Creative Commons Attribution–NonCommercial–NoDerivatives (CC BY-NC-ND) license.

Edited by:
Sugiyama
Axolotl Haruka

動物意識の大阪国際会議 (Animal Consciousness Conference in Osaka) **No.7**

How Does the Axolotl Feel?

アホロートルはどのように感じるのか？

○**Sugiyama Axolotl Haruka**[1], Yoshichika Yokoe[1,2,3,4], F.N. Santako[1], Ryuki Tatsuoka[4], Axolotl Taro[3]

([1] HAL lab (Axolotl Ecology Research Unit), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology, [5] Ambys Lab, [6] Hal lab farm)

E-mail adress sugiyama.haruka.axoltl@gmail.com **Twitter URL** https://twitter.com/lab_new2


CC BY NC ND

1). Question: Can the Amphibians (*Urodeles, including Ambystoma mexicanum*) "Perceive / Recognize"?

Various environmental stimuli provided by the owner humans.




2). Axolotl perception system-1



3). Axolotl perception system-2

B: culture density & Enemy-response system, Growth control (Sugiyama, 2023)




High density with another species (*Hynobius retardatus*)

Low density culture


High density culture

4). Axolotl perception system-3

C: Photo-response system, circadian clock (Sugiyama, 2022-2025)




5). Axolotl recognition (learning) system



6). Summary; "How Does the Axolotl Feel?"

Axolotls' "Feel"

- **Perception (mechanical response)**
  - ✓ Photo-stimulation response (circadian-clock)
    → **eye (central nervous system)**
  - ✓ Salt-stimulation in the water
    → **olfactory system / lateral line (electro-sensory receptor)**
- **Recognition (learning)**
  - ✓ trial-and-error learning
    → **olfaction, lateral line, eye → memory-integration to establish a territory**



CC BY NC ND

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

【総説】魚類・両生類における認知と行動 (Part.1)


Review: Cognition and Behavior in Fish and Amphibians

著: 杉山 遥 [1]; 横江 誼衡 [1, 2, 3, 4]; F.N. 三太子 [1]

[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー

Authors: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]; F.N. Santako[1]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology



※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

導入

動物の認知能力と行動様式は、外界からの感覚入力を神経系で処理し、適応的な行動へとつなげるプロセスに基づいている。視覚・聴覚・嗅覚はその中心的役割を担い、生態環境や生活史に応じて進化的に多様化してきた。こうした感覚と認知の進化は、動物行動学・神経生物学・比較心理学の分野において長年の関心を集めてきた。哺乳類においては、大脳皮質の高度な発達により、色覚や音声認識、嗅覚による社会的認知が精緻に統合され、学習・記憶と結びついた複雑な行動が可能となっている。魚類は、網膜と視蓋を中心とした視覚処理や、側線系と内耳による水中音認識を進化させ、嗅覚とあわせて産卵回帰や餌探索といった生態行動を制御する。一方、両生類は、脊椎動物の進化的中間段階に位置し、多様な感覚様式と認知行動を示す点で特に注目される。

有尾類（アホロートル、タイガーサラマンダー、サンショウウオ類など）は夜行性への適応を反映して視覚よりも嗅覚に依存する傾向が強く、個体認知や繁殖行動における化学的シグナルが重要である。無尾類（アフリカツメガエル、ウシガエル、ヒキガエルなど）は鳴き声を介した社会的コミュニケーションを発達させ、聴覚機能が個体間相互作用の基盤を形成している。さらにオオサンショウウオのように、匂いと水流感覚を統合して縄張り認知や配偶行動を行う種も存在し、感覚特性と行動様式の多様性が際立つ。

加えて、扁形動物や環形動物といった原始的な無脊椎動物にも、光受容や機械感覚を通じた行動応答が見られることは、複雑な脳構造を持たない動物においても、外界刺激に基づく認知と行動が進化的に保存されていることを示している。

本総説では、哺乳類から魚類・両生類、さらに無脊椎動物に至るまでの比較を通じて、認知と行動を支える感覚系と脳領域の進化的特性を概観する。特に両生類は、視覚・嗅覚を介した個体認知と、聴覚に基づく社会的認知を兼ね備える点で、脊椎動物における認知進化を理解するための重要な比較モデルとなる。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Introduction

Cognitive abilities and behavioral patterns in animals are grounded in the processing of sensory inputs by the nervous system and their translation into adaptive responses. Vision, audition, and olfaction play central roles in this process and have diversified through evolution according to ecological demands and life histories. The evolution of sensory and cognitive systems has therefore been a central theme in ethology, neurobiology, and comparative psychology.

In mammals, the advanced development of the cerebral cortex enables sophisticated integration of color vision, auditory recognition, and olfactory cues in social contexts, supporting complex behaviors linked to learning and memory. Fish, by contrast, rely heavily on visual processing through the retina and optic tectum, as well as mechanosensory and auditory inputs from the lateral line system and inner ear. These sensory modalities, combined with olfaction, regulate key ecological behaviors such as spawning migrations and foraging.

Amphibians, positioned as an intermediate group in vertebrate evolution, are of particular interest due to their diverse sensory capacities and cognitive behaviors.

Among caudates (e.g., axolotl, tiger salamander, and various newts), adaptation to nocturnality has favored reliance on olfactory cues, which play crucial roles in individual recognition and reproductive behavior. In anurans (e.g., African clawed frog, bullfrog, toads), acoustic communication is central to social interactions, with auditory function forming the basis of mate choice and territorial behaviors. The Japanese giant salamander, for example, integrates olfactory and mechanosensory cues from water flow to regulate territoriality and mating. These examples highlight the diversity of sensory strategies and their behavioral correlates in amphibians.

Furthermore, in more primitive invertebrates such as flatworms and annelids, light perception and mechanosensation contribute to basic behavioral responses, indicating that cognition and behavior based on environmental stimuli are evolutionarily conserved even in organisms lacking complex brains.

This review provides a comparative overview of sensory systems and neural regions underlying cognition and behavior across mammals, fish, amphibians, and invertebrates. Amphibians, in particular, stand out as key comparative models, as they integrate olfactory and auditory recognition with visual processing, offering critical insights into the evolutionary trajectory of vertebrate cognition.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

動物の認知と行動：視覚・色覚と脳領域を中心にした比較

---

1. 哺乳類における認知と行動

哺乳類は高度に発達した**大脳新皮質**を有し、複雑な感覚処理や行動調節を可能にしている。

- **視覚経路と脳領域**
  網膜で受容された光刺激は外側膝状体（LGN）を経て一次視覚野（V1）に投射される。V1 には**方位選択性カラムや両眼視差カラム**が存在し、形状や奥行きを解析する。情報は V2、V4、MT へと分岐し、**V4 は色覚処理、MT は運動視**に特化している。さらに腹側経路（「何」経路）は側頭葉に至り物体認識を担い、背側経路（「どこ」経路）は頭頂葉に至り空間認知や運動制御を担う。
- **色覚の進化**
  霊長類は三色型色覚（L/M/S 錐体）を進化させ、果実や社会的シグナル（顔色など）の識別を可能にした。一方、齧歯類などは夜行性適応のため杆体依存が強く、色覚能力は低い。
- **高次認知**
  前頭前野は視覚情報と記憶・情動を統合し、**意思決定、社会的行動、感情調節**を担う。

---

2. 魚類における認知と行動

魚類は水中環境に特化した視覚システムを進化させた。

- **視覚処理**
  網膜からの入力は**視蓋（optic tectum）へ投射され、これは初期脊椎動物において大脳皮質に相当する主要視覚中枢である。多くの魚類は四色型～紫外感受性**を備え、複雑な水中環境に適応している。
- **行動機能**
  視蓋は捕食・回避・縄張り行動を統合する。シクリッド類では**色覚が配偶者選択や種識別**に不可欠であり、サケ科では光・嗅覚・地磁気などを統合して回遊を行う。

---

3. 両生類における認知と行動

両生類は水生から陸生への移行過程に位置し、独自の視覚行動システムを持つ。

- **視覚神経回路**
  網膜入力は主に視蓋に投射され、特に運動検出に優れる。カエルでは小さな動く獲物に反応する「**虫検出ニューロン**」が存在する。
- **色覚と行動**
  紫外光を含む複数の錐体型を持ち、摂食や求愛に利用される。アフリカツメガエル (*Xenopus laevis*) は UV 視覚を餌探索に用いることが知られている。
- **発生段階の変化**
  変態期に網膜回路や視蓋ネットワークが再編成され、幼生期（水中捕食）から成体期（陸上捕食・繁殖）へと適応する。

---

4. 扁形動物（Platyhelminthes）における認知と行動

扁形動物は単純ながら再生能力や初歩的な学習能力を示す。

- **視覚**
  眼点（ocelli）は光強度と方向を感知し、脳神経節に信号を送る。色識別はほぼできないが、負の走光性を示す。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **認知能力**
  プラナリアは**条件学習（光や電気刺激と餌の連合）**を示し、神経可塑性を持つ。再生後も記憶が残存する例が報告されている。

---

5. 環形動物（Annelida）における認知と行動

環形動物は分節化した神経系をもち、単純ながら高度な反射行動を実現する。

- **感覚器**
  皮膚光受容器により昼夜リズムを感知する。多毛類では眼点が発達し、光・色識別も可能。
- **神経回路**
  各体節に神経節があり反射制御を行う。前部脳は方向運動や群集行動・交尾行動を統合する。

比較的まとめ

- **哺乳類**： 新皮質を基盤に高度な認知を発揮
- **魚類・両生類**： 視蓋を中心に感覚と行動が直結
- **扁形動物・環形動物**： 単純回路ながら光受容・学習・社会行動を実現

進化の過程で、単純な光応答から**統合的な知覚・認知行動**へと発展してきたことがわかる。

参考文献

1. Kandel ER, Schwartz JH, Jessell TM. 『神経科学の原理』 McGraw-Hill, 2013.
2. Hubel DH, Wiesel TN. 猫の視覚野における受容野と機能構造. *J Physiol*. 1962.
3. Land MF, Nilsson D-E. 『Animal Eyes』 Oxford Univ. Press, 2012.
4. Roth G, Wullimann MF. 『Brain Evolution and Cognition』 Wiley, 2001.
5. Ingle DJ. カエルの二重視覚システム. *Science*. 1973.
6. Fritsches KA, Marshall NJ, Warrant EJ. サンゴ礁魚類における視覚と行動. In: Sale PF (ed.), *Ecology of Fishes on Coral Reefs*. Academic Press, 2015.
7. Inoue D, Yamashita M, Yoshizawa M. 両生類の視覚システムとその進化. *Zool Sci*. 2020.
8. Baguñà J, Saló E. 『Platyhelminthes: Development, Biology, and Evolution』 CRC Press, 2010.
9. Bullock TH, Horridge GA. 『無脊椎動物の神経系の構造と機能』 Freeman, 1965.
10. Arendt D. 神経系集中化の進化. *Philos Trans R Soc Lond B Biol Sci*. 2008.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Cognition and Behavior in Animals: A Comparative Perspective on Visual and Neural Mechanisms

---

1. Cognition and Behavior in Mammals

Mammals possess a highly developed **neocortex**, which supports complex sensory processing and behavioral regulation.

- **Visual pathways and brain regions**
  Retinal input is transmitted via the lateral geniculate nucleus (LGN) to the primary visual cortex (V1). Within V1, **orientation-selective and ocular dominance columns** extract basic features such as orientation and binocular disparity. Signals diverge into higher areas: V4 for **color perception** and MT for **motion vision**. The ventral (“what”) stream projects to the temporal lobe for object recognition, while the dorsal (“where/how”) stream projects to the parietal lobe for visuospatial and motor integration.
- **Evolution of color vision**
  Primates developed trichromatic vision (L/M/S cones), enabling fruit detection and social signal recognition (e.g., facial coloration). By contrast, nocturnal rodents are predominantly rod-dependent and exhibit limited color vision.
- **Higher cognition**
  The prefrontal cortex integrates visual input with memory and emotion to regulate **decision-making, social interactions, and emotional control**.

---

2. Cognition and Behavior in Fish

Fish have evolved sensory systems specialized for aquatic environments.

- **Visual processing**
  Retinal signals are primarily relayed to the **optic tectum**, which serves as the major visual center in teleosts, analogous in part to cortical processing in mammals. Many species exhibit **tetrachromatic or ultraviolet sensitivity**, facilitating adaptation to complex underwater light conditions.
- **Behavioral functions**
  The tectum integrates visual input for prey capture, predator avoidance, and territorial defense. In cichlids, **color vision is critical for mate choice and species recognition**, while salmonids integrate visual, olfactory, and geomagnetic cues for migration.

---

3. Cognition and Behavior in Amphibians

Amphibians represent a transitional group from aquatic to terrestrial life, with specialized visual mechanisms.

- **Visual neural circuits**
  Retinal projections primarily target the optic tectum, which is highly specialized for motion detection. Frogs possess “**bug-detector neurons**” that respond selectively to small moving prey.
- **Color vision and behavior**
  Multiple cone types, including UV-sensitive cones, enable prey capture and mate recognition. *Xenopus laevis* uses UV vision to optimize foraging.
- **Developmental transitions**
  During metamorphosis, retinal circuits and tectal networks are reorganized to shift from larval (aquatic feeding) to adult (terrestrial hunting and reproductive) behavior.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. Cognition and Behavior in Platyhelminthes (Flatworms)

Flatworms exhibit simple but informative sensory systems with regenerative capacity.

- **Vision**
  Ocelli detect light intensity and direction and project to the cerebral ganglia. They lack true color vision but exhibit negative phototaxis.
- **Cognitive abilities**
  Planarians demonstrate **associative learning**, forming conditioned responses to light or electric stimuli paired with food. Remarkably, memory traces may persist after regeneration, suggesting distributed neural plasticity.

---

5. Cognition and Behavior in Annelids

Annelids exhibit segmented nervous systems capable of reflex control and simple cognition.

- **Sensory systems**
  Cutaneous photoreceptors mediate circadian and diel rhythms. Polychaetes possess ocelli that enable detection of both light and basic spectral differences.
- **Neural circuits**
  Segmental ganglia regulate reflexive locomotion, while the anterior brain coordinates **directional movement, swarming, and mating behavior**.

Comparative Summary

- **Mammals:** Neocortex-based higher cognition and sensory integration.
- **Fish & Amphibians:** Optic tectum-centered processing linking visual cues directly to survival behaviors.
- **Flatworms & Annelids:** Simple neural architectures still allow **light perception, associative learning, and coordinated social/sexual behaviors**.

This comparative framework illustrates an evolutionary trajectory from basic photoreception to the emergence of **integrated perception, cognition, and behavior** across the animal kingdom.

References

1. Kandel ER, Schwartz JH, Jessell TM. *Principles of Neural Science*. McGraw-Hill, 2013.
2. Hubel DH, Wiesel TN. Receptive fields and functional architecture in the cat's visual cortex. *J Physiol*. 1962.
3. Land MF, Nilsson D-E. *Animal Eyes*. Oxford University Press, 2012.
4. Roth G, Wullimann MF. *Brain Evolution and Cognition*. Wiley, 2001.
5. Ingle DJ. Two visual systems in the frog. *Science*. 1973.
6. Fritsches KA, Marshall NJ, Warrant EJ. Vision and behavior of reef fishes. In: Sale PF (ed.), *Ecology of Fishes on Coral Reefs*. Academic Press, 2015.
7. Inoue D, Yamashita M, Yoshizawa M. Visual system of amphibians and its evolution. *Zool Sci*. 2020.
8. Baguñà J, Saló E. *Platyhelminthes: Development, Biology, and Evolution*. CRC Press, 2010.
9. Bullock TH, Horridge GA. *Structure and Function in the Nervous Systems of Invertebrates*. Freeman, 1965.
10. Arendt D. The evolution of nervous system centralization. *Philos Trans R Soc Lond B Biol Sci*. 2008.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

有尾類（イモリ・サンショウウオ）の認知と行動

1) 視覚・色覚（行動との相関）

基本生理

- 網膜：桿体優勢（薄暗所で高感度）＋複数種類の錐体。水生・昼行性の種ほど錐体比率が高め。
- 視覚中枢：主に**中脳視蓋（optic tectum）**で統合（哺乳類の皮質的視覚処理に相当する役割）。
- 空間分解能は高くないが、**小型・移動物体の検出**に非常に優れる（獲物追尾に直結）。

色覚

- S（短波長）～L（長波長）域の複数錐体をもち、**種・生活様式により UV～青～緑～赤の感受域が変動**。
- 完全夜行性・地中性では機能的色覚は弱くなりがち。逆に**水生・昼行性のイモリ（例：アカハライモリ等）**では、明所下での色識別が餌選好・配偶行動に寄与。
- **警告色・腹面の強色彩**（例：アカハラ）を示す種は、捕食者に対する**威嚇・忌避（腹見せ姿勢）**に色を用いる。一方、**同種間の微妙な体色差**が繁殖期のコンディション合図になりうる。

行動対応

- **採餌**：視覚誘発の捕食（「小さく・速く・断続的に動く」対象へ強反応）。背景コントラストや短波長光は餌検出を助けることがある。
- **配偶**：多くは化学・触覚主導だが、**求愛姿勢・婚姻色**など視覚的シグナルが補助的に働く種あり。
- **捕食回避**：素早い運動検出→逃避反応。腹面の赤色提示（ポーズ）で抑止。
- **ナビゲーション**：ランドマーク利用や明暗勾配を用いた環境定位が可能。

---

2) 聴覚・振動感知（行動との相関）

聴覚の特徴

- **外耳・鼓膜を欠く**ため、空気音の聴取は弱いが、**内耳（columella）と肩帯に連結する“オペルクラ系（operculum”）により低周波振動・地面伝播振動**に感受性。
- 水生期・水生種では**側線系**が発達し、水流・振動を検知して**捕食・回避**に利用。

行動対応

- **捕食者検知**：足場や水底を伝わる微小振動で捕食者接近を早期察知。
- **採餌**：水流の乱れから生体の動きを検出（側線）。
- **コミュニケーション**：有声化は稀だが、**姿勢・接触・化学信号**と併用して低頻度振動（踏み鳴らし様）を使う例も。

---

3) 化学感覚・認識能力（行動の中核）

嗅覚・ヤコブソン器（鋤鼻器）

- 有尾類の**社会認知は化学感覚が中核**。水中では溶存化学物質、陸上では揮発性化合物を利用。
- **個体識別・性成熟・繁殖状態・縄張り**・種識別・**捕食者の匂い学習**など、ほぼ全域をカバー。

行動対応

- **配偶行動**：雄のフェロモン（頸部や精神腺由来）→雌の受容性上昇、**テールストラドリング・spermatophore 受け渡し**の成功率を上げる。
- **縄張り・競争**：痕跡化学シグナルで相手の**性別・サイズ・優劣**を推定、闘争・回避を調節。
- **捕食回避学習**：天敵の匂いに対する**条件づけ**が成立し、長期記憶を形成。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4) 学習・記憶・空間認知

- **連合学習（古典・道具立て）**：光・振動・匂いと餌／回避刺激の結びつけが可能。
- **迷路学習・ランドマーク利用**：T 迷路・十字迷路での到達学習、**ランドマーク再定位・経路再計画**を示す。
- **可逆学習（リバーサル）**：規則変更への適応（以前の刺激価の反転）も学習可能。
- **時間・場所の手がかり**：摂餌の時刻予測や、繁殖場への**帰巣（homing）**が報告される種あり。
- **磁気受容の示唆**：一部のイモリで、**地磁気方位への選好**が示される報告（定位の補助）。

---

5) 生活史・種差によるモジュール性

- **幼生 vs 成体**：幼生は側線・水中視覚・化学感覚で捕食・回避／**変態で回路が再編**し、陸上適応（嗅覚・触覚の比重上昇）。
- **水生 vs 陸生**：水生種は側線・視覚（とくに動き検知）が強く、陸生・森林性では**嗅覚・触覚主導**。
- **昼行 vs 夜行**：昼行は錐体比率↑・色覚の行動利用↑、夜行は桿体系主導（明暗・動き中心）。

---

6) 実験での評価指標（実務のヒント）

- **視覚**：小型移動刺激への追尾、色選好試験（明所・暗所、背景コントラスト統制）。
- **振動／側線**：周波数別振動刺激に対する驚愕反応・回避閾値の計測。
- **化学認知**：Y 迷路での匂い選択、フェロモン添加での求愛行動変化。
- **学習・記憶**：T 迷路・リバーサル課題、天敵匂いの条件づけ保持期間。
- **生活史**：変態前後の感覚重み付けの遷移（視覚→嗅覚／触覚）を縦断評価。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Cognition in Newts and Salamanders (Urodeles): Vision, Hearing/Vibration, Recognition, and Behavior

Vision & Color

- **Retina**: rod-dominant (dim-light sensitivity) plus multiple cone classes; cone proportion higher in aquatic/diurnal species.
- **Central processing**: the **optic tectum** is the principal visual hub, specialized for **small, moving prey**.
- **Color**: several cone types (often spanning S to L; UV sensitivity in some taxa). Color use scales with ecology—stronger under photopic, aquatic conditions.
- **Behavioral links**:

  – **Foraging**: robust orienting to small moving items; contrast and short-wavelength cues can facilitate prey detection.

  – **Courtship**: primarily chemical/tactile, but **visual postures and nuptial displays** can modulate receptivity.

  – **Antipredator**: rapid motion-triggered escape; **ventral aposematic display** (e.g., bright bellies) to deter predators.

  – **Navigation**: landmark use and light gradients assist spatial orientation.

Hearing & Vibration

- No external ear/tympanum; airborne hearing is limited. The **columella–operculum system** and body conduction enhance **low-frequency and substrate-borne** sensitivity.
- Aquatic species/larvae use the **lateral line** to detect water movements.
- **Behavioral links**: predator detection via ground/water vibrations; prey localization through hydrodynamic cues; limited vibratory signaling may complement chemical/tactile displays.

Chemosensory Recognition (Core of Social Cognition)

- **Olfactory & vomeronasal systems** dominate **individual/sex/species/state recognition**.
- **Courtship**: male pheromones (e.g., mental glands) elevate female receptivity; guide spermatophore uptake.
- **Territoriality**: chemical marks inform size/sex/dominance, reducing escalated fights.
- **Predator learning**: rapid conditioning to predator odors; long-lasting memory.

Learning, Memory, Spatial Cognition

- **Associative learning** (classical/instrumental) with visual, chemical, or vibratory cues.
- **Maze and landmark learning**, **reversal learning**, and **homing** reported in several species.
- **Magnetoreception** has been suggested in some newts for orientation.

Life-History Modularity

- **Larva → Adult**: reweighting from lateral-line/aquatic vision to olfaction/tactile systems after metamorphosis.
- **Aquatic vs Terrestrial; Diurnal vs Nocturnal**: predictably shifts reliance among color vision, motion detection, chemosensation, and tactile/vibration sensing.

Practical Assays

- Vision: small-object tracking; color-choice under controlled illuminance.
- Vibration: startle/avoidance thresholds across low frequencies; lateral-line assays.
- Chemosensory: Y-maze odor preference; pheromone-induced courtship changes.
- Learning: T-maze, reversal tasks; predator-odor conditioning retention.
- Developmental comparisons pre-/post-metamorphosis.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

有尾類（イモリ・サンショウウオ）の認知能力：視覚・聴覚（振動）・化学感覚と行動

0. 俯瞰：有尾類の共通骨格

- **視覚**：網膜は桿体優勢（薄明・低照度対応）＋複数の錐体。脳の主要視覚中枢は**中脳視蓋（optic tectum）**。
- **色覚**：SWS1（UV～近紫外）・SWS2（青）・RH2（緑）・LWS（黄～赤）オプシンが一般的（発現・実効は種・生活史で変動）。
- **聴覚／振動**：外耳・鼓膜は欠如。**コルメラ（columella）～オペルキュラム（operculum）系**で**低周波・基質振動**を感受。水生期・水生種では**側線系**が水流・振動検出に寄与。
- **化学感覚**：嗅上皮＋**鋤鼻器（vomeronasal organ）**が高度に発達。社会認知・繁殖に中核。
- **代表的行動相関**：小型移動物体への強い視覚誘発捕食／フェロモン誘起求愛／振動による捕食者察知／帰巣・方位選好（ランドマーク・匂い・地磁気の統合）。

---

1. 種別プロファイル（感覚特性 × 行動）

A) アカハライモリ *Cynops pyrrhogaster*（日本：半水生）

- **視覚・色覚**：日中～薄暮活動で錐体の寄与が比較的大。SWS～LWS まで幅広く、**明所での色識別**が餌選好・社会シグナルの補助に。腹面の**赤色（警告色）**は防御ポーズで提示。
- **振動／側線**：水中では側線が有効、底床振動で回避反応。
- **化学認知**：雄が**ソデフリン（sodefrin）**等のペプチド系フェロモンを放出→雌の受容性上昇、テールストラドリング～**精包（spermatophore）受け渡し**を促進。
- **行動**：小型移動刺激への追尾、匂いによる餌・配偶者識別、腹見せ忌避。

B) アカウミイモリ属／ヨーロッパ系イモリ

- **スムーズニュート *Lissotriton vulgaris***（半水生）：
  - 視覚での**求愛ディスプレイ**（尾の波打ち）＋化学信号。明所下で視覚シグナルの効果↑。
  - 水中側線で獲物検出。
- **オオイモリ *Triturus cristatus*（グレートクレステッドニュート）**：
  - 繁殖期にオスの**背びれ・尾の誇示**（視覚）＋接触・化学。
  - 色コントラスト・輪郭がディスプレイの視認性を高める。

C) アメリカイモリ *Notophthalmus viridescens*

- **生活史**：水生幼生 → **陸生エフト（赤色）** → 水生成体。
- **視覚**：水生期は小型移動刺激に強反応。陸生期は明暗・コントラスト依存が増す。
- **化学・認知**：個体識別・捕食者匂い学習が顕著。
- **方位**：**地磁気方位選好**の報告（帰巣・定位の補助）。

D) サンショウウオ属・ファイアサラマンダー等

- **トウホク・クロサンショウウオ（*Hynobius* spp.／森林性）**：
  - 夜行・地上生活で**嗅覚優位**。視覚は動き検知中心。振動で捕食者察知。
- **ファイアサラマンダー *Salamandra salamandra***：
  - **黄黒の警告色**（毒腺）を視覚提示。化学で個体・テリトリー認識。

E) ウーパールーパー（アホロートル）*Ambystoma mexicanum*（終生幼形成熟：水生）

- **視覚**：暗所適応強く色覚は限定的だが、**動き検知・コントラスト**で採餌。
- **側線**：発達。水流・振動で獲物・捕食者認知。
- **化学**：高い。餌・個体識別、ストレス／病的水質の回避。
- **行動**：低照度での探索、振動・匂いで採餌精度↑。

F) プレソドン科（*Plethodon* spp.／北米：完全陸生・肺なし）

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **視覚**：夜行・森林性で桿体系寄り。
- **化学**：雄の**メンタルグランド**（頤部腺）由来フェロモン（PMF/PRF 等）で雌受容性変化。
- **社会行動**：化学痕跡で縄張り・競争回避。嗅・触が中心、視覚は補助。

---

2. 感覚—行動対応の要点（横断的）

感覚系	生理・器官	主な行動	代表例
視覚（色・動き）	桿体＋複数錐体（SWS1/2, RH2, LWS）、視蓋が主中枢	採餌（小物体・動き）、警告色提示、求愛ディスプレイ	*Cynops*（腹赤の提示）、*Triturus*（背びれ誇示）
振動・側線	operculum–columella 系、側線（幼生・水生種）	捕食者検知、獲物定位（流水検出）、驚愕回避	*Ambystoma mexicanum*, *Notophthalmus*
化学（嗅・鋤鼻）	嗅上皮・鋤鼻器、フェロモン腺	配偶者受容、個体・種・状態識別、縄張り	*Cynops*（ソデフリン）、*Plethodon*（メンタル腺）
空間認知	視覚ランドマーク、匂い、（一部）地磁気	帰巣・繁殖池移動、経路再計画、リバーサル学習	*Notophthalmus*, *Ambystoma* spp.

---

3. 実験プロトコル（目安・研究実務に使える条件）

3-1 視覚（色・動き）

- **色選好テスト（Y 迷路）**
  - **照明**：狭帯域 LED（例：UV 365–380 nm／青 440–460／緑 520–540／赤 620–640）。
  - **照度**：明所 150–300 lx、水生種は 50–150 lx から調整。
  - **水質**：導電率・pH・濁度一定（色の見え方を安定化）。
  - **制御**：輝度等価化（photometric matching）、側偏好の交互入替、ランダム化。
  - **指標**：選択率・滞在時間・到達潜時。
- **動き検出／オプトモータ応答**
  - **刺激**：移動グレーティング（空間周波数 0.05–0.3 cyc/deg、速度 10–40 deg/s）。
  - **指標**：追従泳動、頭部サッケード、捕食スナップ頻度。

3-2 振動／側線

- **基質振動**
  - **装置**：加振テーブル（10–200 Hz）。**加速度**をキャリブレーション（例：0.01–0.1 g の段階）。
  - **指標**：驚愕・逃避開始閾値、潜時、移動距離。
- **水流検出（側線）**
  - **刺激**：パルス水流（ピエゾ／ポンプ）、流速 0.5–5 cm/s。
  - **指標**：定位角度、捕食スナップの的中率、探索軌跡。

3-3 化学認知（嗅・鋤鼻）

- **Y 迷路匂い選好**
  - **刺激**：同種雄フェロモン抽出液／餌抽出液／捕食者抽出液。
  - **流量**：各アーム等流（乱流回避）、導入濃度段階化。
  - **指標**：アーム選択率、滞在時間、探索軌跡。
- **求愛誘起**
  - **添加**：雄抽出フェロモン（*Cynops*：ソデフリン系、*Plethodon*：PMF/PRF）。
  - **指標**：雌の追従・テールストラドリング受容、精包受容率。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

3-4 学習・記憶

- **T 迷路／十字迷路**：餌・匂い手掛かりで到達学習、**リバーサル課題**で行動柔軟性評価。
- **捕食者匂い条件づけ**：CS（新規匂い）＋US（捕食者匂い）→回避反応の保持期間。
- **帰巣／方位**：アリーナ内の地磁気回転（ヘルムホルツコイル）で方位選好をテスト（**倫理・遮蔽配慮**必須）。

※上記の数値は「典型的レンジ」の目安です。個体差・種差が大きいため、**予備実験で最適化**してください。

---

4. 実務 Tips（飼育・再現性）

- **環境**：昼夜照明（スペクトル含む）・水質（EC, pH, 溶存有機物）を記録・統制。
- **生活史**：幼生・エフト・成体で感覚重みが変わるため**段階別に測定**。
- **動機づけ**：給餌前後で反応が変化。**標準化した摂食スケジュール**を。
- **福祉**：低ストレス取り扱い・適切な休息・温度域の遵守。
- **統計**：反復測定（個体内要因）＋混合効果モデルで個体差を吸収。

参考文献（抜粋）

- Kikuyama S. et al. (1995) A female-attracting peptide pheromone in the Japanese newt. *Nature*.
- Houck LD., Arnold SJ., Feldhoff RC., et al. (2008–) Courtship pheromones in plethodontid salamanders.
- Phillips JB. (1986–) Magnetic orientation in the eastern red-spotted newt.
- Roth & Wullimann (2001) *Brain Evolution and Cognition*. Wiley.
- Land & Nilsson (2012) *Animal Eyes*. OUP.
- Ingle DJ. (1973) Two visual systems in the frog. *Science*.（両生類視蓋の動き検出）
- Petranka JW. (1998) *Salamanders of the United States and Canada*. Smithsonian.
- Døving KB., Trotier D.（嗅・鋤鼻の書籍・総説）
- Northcutt RG.（側線・脳区画の比較神経解剖）

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Urodeles (Newts and Salamanders): Sensory–Behavior Coupling

Species Profiles

- **Cynops pyrrhogaster (Japanese fire-bellied newt)**
  Visual detection of **small moving prey**; conspicuous **ventral redness** for aposematic display. Strong **pheromonal control** of courtship (e.g., **sodefrin** peptides), enhancing tail-straddling and spermatophore uptake. Lateral line and substrate-borne vibration aid predator detection.
- **Lissotriton vulgaris / Triturus cristatus (European newts)**
  Male **visual displays** (crest, tail-fanning) augment chemical/tactile cues in courtship. Color contrast and outline enhance display detectability under photopic conditions.
- **Notophthalmus viridescens (Eastern newt)**
  Aquatic larva → **terrestrial eft** → aquatic adult. Robust visual prey capture in water; strong **chemical recognition** (conspecifics, predators). Evidence for **magnetoreception** in orientation/homing.
- **Ambystoma mexicanum (axolotl, paedomorphic aquatic)**
  Low-light vision with limited color; **lateral-line–based hydrodynamic sensing** is prominent. Chemosensation supports foraging and stress-avoidance decisions.
- **Plethodontids (e.g., *Plethodon* spp., lungless, terrestrial)**
  Nocturnal, forest-floor specialists with **chemosensory dominance**. Males deliver **mental-gland pheromones** (e.g., PMF/PRF families) to elevate female receptivity. Visual cues are secondary.

Sensory Systems and Behavior

- **Vision/Color**: Rod-dominant retina plus multiple cones (SWS1/2, RH2, LWS). Optic tectum implements motion- and small-object detection. Color use scales with ecology (aquatic/diurnal > terrestrial/nocturnal).
- **Hearing/Vibration**: No tympanum; **columella–operculum apparatus** and body conduction yield low-frequency/substrate sensitivity. **Lateral line** in larvae/aquatic forms for hydrodynamic cues.
- **Chemosensation**: Olfactory + vomeronasal systems underpin social recognition, courtship, territoriality, and predator learning.
- **Spatial Cognition**: Landmark use, odor trails, and in some species **magnetic field** cues support homing.

Practical Assays (recommended ranges)

- **Color choice (Y-maze)**: Narrow-band LEDs (UV 365–380, blue 440–460, green 520–540, red 620–640 nm); 150–300 lx photopic; luminance-matched arms; endpoints = arm choice, dwell time, latency.
- **Optomotor/motion**: Moving gratings 0.05–0.3 cyc/deg at 10–40 deg/s; track pursuit, saccades, snap counts.
- **Substrate vibration**: Shaker plate 10–200 Hz; calibrated acceleration 0.01–0.1 g; endpoints = startle threshold, escape latency/distance.
- **Hydrodynamic (lateral line)**: Pulsed jets 0.5–5 cm/s; endpoints = orientation angle, strike accuracy.
- **Chemosensory**: Y-maze with conspecific pheromones/food/predator extracts; equal flow in arms; endpoints = preference, dwell time, trajectory.
- **Courtship induction**: Add male pheromones (e.g., sodefrin/PMF/PRF) and score female following, tail-straddling, spermatophore uptake.
- **Learning**: T-/plus-mazes for acquisition and **reversal**; predator-odor conditioning with retention tests; magnetic arena (Helmholtz coils) with strict shielding and welfare oversight.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Notes for Reproducibility

- Control **illumination spectrum**, **water quality** (EC, pH, turbidity), and **feeding state**.
- Stratify by **life stage** (larva/eft/adult) and ecology (aquatic vs terrestrial).
- Use mixed-effects models to handle individual variability; ensure adequate replication.

References (selected)

- Kikuyama S. et al. 1995. Female-attracting peptide in *C. pyrrhogaster*. *Nature*.
- Houck LD., Feldhoff RC. Courtship pheromones in plethodontids (multiple works).
- Phillips JB. Magnetic orientation in *N. viridescens* (series).
- Land & Nilsson. *Animal Eyes*. OUP.
- Roth & Wullimann. *Brain Evolution and Cognition*. Wiley.
- Ingle DJ. 1973. Two visual systems in frogs. *Science*.
- Petranka JW. *Salamanders of the United States and Canada*. Smithsonian.
- Døving & Trotier (olfactory/vomeronasal reviews), Northcutt RG (comparative neuroanatomy).

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

有尾類・無尾類の認知能力と行動

1. **メキシコサラマンダー（アホロートル *Ambystoma mexicanum*）**
   - **視覚**：桿体優位、暗所適応型。動き・コントラスト検知が中心。
   - **嗅覚・側線**：餌認知や水質感知に強力。微細な水流や振動を感知。
   - **行動**：夜行性、嗅覚と視覚を組み合わせ餌探索。

2. **タイガーサラマンダー *Ambystoma tigrinum***
   - **視覚**：陸生時は光刺激に敏感。
   - **振動・嗅覚**：基質振動・捕食者匂い学習に反応。
   - **行動**：環境学習による帰巣性。

3. **蝦夷サンショウウオ *Hynobius retardatus***
   - **視覚**：夜行性で動き認識が主。
   - **嗅覚**：繁殖期のフェロモン認知。
   - **行動**：産卵池の選択、捕食者匂い回避学習。

4. **セトウチサンショウウオ *Hynobius setouchi***
   - **視覚**：森林・渓流環境に適応、暗所で動き検知。
   - **嗅覚**：繁殖期にフェロモンで雌雄間認知。
   - **行動**：産卵時に水域環境（匂い・温度）を敏感に選択。

5. **東京サンショウウオ *Hynobius tokyoensis***
   - **視覚**：同属と同じく夜行性視覚。
   - **嗅覚**：水質や同種フェロモンの認識が繁殖行動の鍵。
   - **行動**：都市近郊でも微妙な環境変化を嗅覚で検知。

6. **イベリアトゲイモリ *Pleurodeles waltl***
   - **視覚・嗅覚**：動き検知と化学認知を併用。
   - **行動**：外敵に対し毒腺防御（リブ突出）。

7. **オオサンショウウオ *Andrias japonicus***
   - **視覚**：水中の動きに対する反応は弱め。夜行的。
   - **嗅覚**：極めて発達し、捕食（魚・甲殻類）やナワバリ認知に重要。
   - **行動**：河川の巣穴に定住、嗅覚による環境認知が優位。

8. **アフリカツメガエル *Xenopus laevis***
   - **視覚**：動き認知中心。
   - **聴覚**：水中音・クリック音による繁殖行動。

9. **ネッタイツメガエル *Xenopus tropicalis***
   - **視覚**：浅水環境適応。
   - **行動**：繁殖時のクリック音に反応。

※各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

10. ウシガエル *Lithobates catesbeianus*

- **視覚**：優れた動体視力。獲物（昆虫、小魚）を遠距離から捕捉。
- **聴覚**：発達した鼓膜と共鳴嚢、繁殖期の鳴き声によりオス間競合・雌誘引。
- **行動**：縄張り形成、鳴き声を通じた社会的認知が強調される。

11. ヒキガエル *Bufo japonicus*

- **視覚**：暗所で動きを識別、色覚は限定的。
- **嗅覚**：繁殖期に雄が雌を識別する重要手段。
- **行動**：帰巣性が強く、繁殖池に戻る習性あり。

比較まとめ

- **サンショウウオ類**：嗅覚依存度が高く、繁殖期の環境選択や捕食者回避に関与。
- **カエル類**：聴覚・鳴き声による社会的認知が顕著。
- **大型種（オオサンショウウオ、ウシガエル）**：嗅覚や聴覚が極めて発達し、縄張りや繁殖行動を強く支配。

参考文献

- Wells, K.D. (2007). *The Ecology and Behavior of Amphibians*. Univ. Chicago Press.
- Roth & Wullimann (2001). *Brain Evolution and Cognition*. Wiley.
- Elepfandt A. (1996). Hearing in *Xenopus laevis*. *J Comp Physiol A*.
- Kikuyama S. et al. (1995). Sodefrin pheromone in *Hynobius*. *Nature*.
- Browman H.I., & Hawryshyn C.W. (1994). Visual pigments in amphibians. *J Exp Biol*.
- Okada Y. (1931). Studies on *Andrias japonicus*. Zoological Magazine (Tokyo).

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Cognition and Behavior in Salamanders, Newts, and Frogs

1. Axolotl (*Ambystoma mexicanum*)

- **Vision**: Rod-dominant, adapted to dim light, motion/contrast-based prey recognition.
- **Olfaction & Lateral Line**: Strong for prey and water chemistry detection.
- **Behavior**: Nocturnal, relies on olfactory + visual cues for feeding.

2. Tiger Salamander (*Ambystoma tigrinum*)

- **Vision**: Sensitive to light in terrestrial phase.
- **Chemosensory/Vibration**: Learns predator odors; substrate vibration sensing.
- **Behavior**: Homing ability via olfactory and landmark learning.

3. Hokkaido Salamander (*Hynobius retardatus*)

- **Vision**: Motion-based, nocturnal.
- **Olfaction**: Reproductive pheromones important.
- **Behavior**: Breeding site selection and predator avoidance learning.

4. Setouchi Salamander (*Hynobius setouchi*)

- **Vision**: Suited for dim stream habitats.
- **Olfaction**: Pheromone-based mate recognition.
- **Behavior**: Sensitive site selection during breeding.

5. Tokyo Salamander (*Hynobius tokyoensis*)

- **Vision**: Motion-based recognition.
- **Olfaction**: Detects conspecific pheromones and water chemistry.
- **Behavior**: Adjusts to microenvironmental cues in urban habitats.

6. Iberian Ribbed Newt (*Pleurodeles waltl*)

- **Vision/Chemosensory**: Combines motion and chemical cues.
- **Behavior**: Unique rib-protrusion defense triggered by combined stimuli.

7. Japanese Giant Salamander (*Andrias japonicus*)

- **Vision**: Limited motion sensitivity, nocturnal.
- **Olfaction**: Highly developed; critical for feeding and territory recognition.
- **Behavior**: Cave-dwelling, olfaction-dominant perception.

8. African Clawed Frog (*Xenopus laevis*)

- **Vision**: Motion-based.
- **Hearing**: Underwater sound and mating calls.

9. Tropical Clawed Frog (*Xenopus tropicalis*)

- **Vision**: Shallow water adaptation.
- **Behavior**: Call-response in reproduction.

10. American Bullfrog (*Lithobates catesbeianus*)

- **Vision**: Excellent motion detection; captures prey at distance.
- **Hearing**: Well-developed tympanum; mating calls regulate territory and courtship.
- **Behavior**: Strong territoriality, acoustic communication central.

11. Japanese Toad (*Bufo japonicus*)

- **Vision**: Detects movement in dim light.
- **Olfaction**: Important for male mate recognition.
- **Behavior**: Strong homing instinct to breeding ponds.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Comparative Summary

- **Salamanders/Newts**: Olfactory-driven, essential for reproduction and predator avoidance.
- **Frogs/Toads**: Auditory communication dominates, especially in social and mating contexts.
- **Large species (e.g., *Andrias*, Bullfrog)**: Specialized sensory dominance (olfaction or hearing) linked to territoriality and reproduction.

References

- Wells, K.D. (2007). *The Ecology and Behavior of Amphibians*. Univ. Chicago Press.
- Roth & Wullimann (2001). *Brain Evolution and Cognition*. Wiley.
- Elepfandt A. (1996). Hearing in *Xenopus laevis*. *J Comp Physiol A*.
- Kikuyama S. et al. (1995). Pheromonal communication in *Hynobius*. *Nature*.
- Browman H.I., & Hawryshyn C.W. (1994). Visual pigments in amphibians. *J Exp Biol*.
- Okada Y. (1931). Studies on *Andrias japonicus*. Zoological Magazine (Tokyo).

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

日本語版：有尾類・無尾類の認知能力と行動の比較表

種名	視覚	聴覚	嗅覚	その他感覚	主な行動特性
メキシコサラマンダー (*Ambystoma mexicanum*)	暗所適応、動き検知中心	弱い	強い、餌探索に重要	側線で水流感知	夜行性、嗅覚＋視覚で捕食
タイガーサラマンダー (*Ambystoma tigrinum*)	陸生期に光刺激敏感	弱い	捕食者匂い学習	振動感知	帰巣性、環境学習
蝦夷サンショウウオ（*Hynobius retardatus*)	動き検知、夜行性	弱い	フェロモン認知	—	産卵池選択、回避学習
セトウチサンショウウオ (*Hynobius setouchi*)	渓流暗所適応	弱い	繁殖期フェロモン重要	—	産卵時の環境選択
東京サンショウウオ（*Hynobius tokyoensis*)	夜行性視覚	弱い	同種フェロモン認知	—	都市環境適応、繁殖行動
イベリアトゲイモリ (*Pleurodeles waltl*)	動き検知	弱い	化学認知発達	皮膚毒腺	防御行動（リブ突出）
オオサンショウウオ（*Andrias japonicus*)	動体視力弱め	弱い	非常に発達	水流・匂い統合	巣穴生活、縄張り認知
アフリカツメガエル（*Xenopus laevis*)	動き検知	水中音に敏感	中程度	側線感覚	鳴き声で繁殖行動
ネッタイツメガエル（*X. tropicalis*)	浅水適応	鳴き声応答	中程度	—	繁殖クリック音応答
ウシガエル（*Lithobates catesbeianus*)	優れた動体視力	発達した鼓膜、鳴嚢	中程度	—	強い縄張り性、鳴き声社会性
ヒキガエル（*Bufo japonicus*)	暗所動き検知	鳴き声による繁殖行動	繁殖期に重要	—	強い帰巣性

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl


English Version: Comparative Table of Cognition and Behavior in Amphibians

Species	Vision	Hearing	Olfaction	Other Senses	Key Behavioral Traits
Axolotl (*Ambystoma mexicanum*)	Rod-dominant, motion detection	Weak	Strong, prey detection	Lateral line	Nocturnal, combined olfactory-visual feeding
Tiger Salamander (*A. tigrinum*)	Light-sensitive in terrestrial phase	Weak	Learns predator odors	Vibration sensing	Homing ability, environmental learning
Hokkaido Salamander (*H. retardatus*)	Motion-based, nocturnal	Weak	Pheromone recognition	—	Breeding site choice, avoidance learning
Setouchi Salamander (*H. setouchi*)	Suited for dim streams	Weak	Pheromone recognition	—	Sensitive breeding site selection
Tokyo Salamander (*H. tokyoensis*)	Nocturnal vision	Weak	Conspecific pheromone recognition	—	Urban adaptation, reproductive behavior
Iberian Ribbed Newt (*Pleurodeles waltl*)	Motion detection	Weak	Developed chemical sensing	Skin toxin glands	Rib-protrusion defense
Japanese Giant Salamander (*Andrias japonicus*)	Limited motion sensitivity	Weak	Highly developed	Flow + odor integration	Cave-dwelling, territorial recognition
African Clawed Frog (*Xenopus laevis*)	Motion-based vision	Sensitive to underwater sound	Moderate	Lateral line	Acoustic mating behavior
Tropical Clawed Frog (*X. tropicalis*)	Adapted to shallow waters	Call-response	Moderate	—	Reproductive acoustic responses
American Bullfrog (*Lithobates catesbeianus*)	Excellent motion detection	Advanced tympanum, vocal sac	Moderate	—	Territoriality, social acoustic signals
Japanese Toad (*Bufo japonicus*)	Dim-light motion detection	Acoustic signals for reproduction	Important in breeding	—	Strong homing instinct

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl


ハルラボの知見

アホロートルは、視覚のみならず音や振動といった感覚刺激を通じて周囲を認識し、学習している傾向がある。暗所よりも明所において餌や飼育者、給餌場所をより的確に識別していることから、視力の程度は必ずしも明瞭ではないものの、一定の視覚情報が行動判断の基盤として利用されていると考えられる。このことは、アホロートルが多様な感覚入力を統合して環境適応を行っていることを示唆しており、視覚が限定的であっても行動制御に一定の寄与を果たしていると推察される。



補足図 1: アホロートルの認知・学習能力 1　(day50 → day100 個体)


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。



補足図 2: アホロートルの認知・学習能力 2　(day50 → day100 個体)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

今回の調査結果から、アホロートルは視覚および側線機能を喪失した場合でも、最終的には音刺激を手がかりとして周囲を認識することが明らかとなった。



補足図 3: アホロートルの認知・学習能力 3　(day50 → day100 個体)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl

【総説】 両生類における抗酸菌症と対策


Review: Mycobacteriosis in Amphibians and Its Control

著: 横江 誼衡 [1, 2, 3, 4]; F.N. 三太子 [1]; 杉山 遥 [1]

[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー

Authors: Yoshichika Yokoe[1,2,3,4]; Sugiyama Axolotl Haruka[1]; F.N. Santako[1]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology



※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

導入

アホロートル（Ambystoma mexicanum）が痩せていく原因については、これまで複数の要因が知られている。本稿では過去の報告を整理し直し、栄養失調、悪性腫瘍、抗酸菌症（魚結核）、その他の感染症、ならびに急激な変態促進（甲状腺異常）を主要因としてまとめる。特に重要なのは、十分に餌を摂取していても回復できない場合であり、この背景には抗酸菌症と甲状腺異常が関与している可能性が高い。

抗酸菌症は元気に見える個体でも進行性の削痩を引き起こすことがあり、伝播性があるため隔離対応が必須となる。ただし、発症しない個体は一生発症しない場合もある。また魚類からアホロートルへ伝播するリスクがあるため、活餌の使用時には薬浴トリートメントが必須である。治療に関しては塩浴の効果は乏しく、抗酸菌はグラム染色で分類が困難であり、抗生物質に対する感受性も低い。そのため、現時点ではメチレンブルーやイソジンによる薬浴治療が推奨されており、特にイソジンがより高い有効性を示すと考えられる。ただし、メチレンブルーの直接的な有効性については議論の余地がある。

臨床的には、初期段階で完全に回復できなかった個体はその後の復帰が困難であることが多い。罹患個体は当初は元気で餌も摂取するが、徐々に腸が萎縮し、消化や排便がうまくいかなくなる。結果として栄養失調が進行し、さらに感染により壊死した組織の修復に大量のエネルギーが消費されることで、痩せが持続・進行する。本稿では、これらの知見を踏まえ、特に抗酸菌症についてさらに掘り下げて論じる。

Introduction

The causes of emaciation in axolotls (Ambystoma mexicanum) have been attributed to several factors. In this review, we reorganize previous reports and highlight the major causes, including malnutrition, malignant tumors, mycobacteriosis (fish tuberculosis), other infectious diseases, and accelerated metamorphosis (thyroid dysfunction). Of particular importance are cases in which affected animals fail to recover despite adequate food intake. Such cases are most likely associated with mycobacterial infection or thyroid dysfunction.

Mycobacteriosis can induce progressive wasting even in axolotls that appear otherwise healthy. Because the disease is transmissible, isolation of affected individuals is essential. Nevertheless, some animals may never develop clinical disease throughout their lifetime. Transmission from fish to axolotls has also been suggested, underscoring the importance of prophylactic treatment of live feed by medicated baths. Regarding therapeutic approaches, salt baths are generally ineffective. Mycobacteria are difficult to classify by Gram staining and exhibit poor susceptibility to conventional antibiotics. For this reason, immersion therapy using methylene blue or povidone-iodine (isodine) has been recommended, with povidone-iodine considered more effective. The direct efficacy of methylene blue, however, remains debated.

Clinically, axolotls that fail to recover completely in the early stages rarely regain normal condition. Infected individuals often remain active and continue feeding initially but gradually develop intestinal atrophy, leading to impaired digestion and defecation. As a result, nutritional deficiency worsens, and a significant proportion of energy is diverted to the repair of necrotic tissues damaged by infection, causing persistent and progressive emaciation. This article will focus on a deeper exploration of mycobacteriosis in axolotls in light of these clinical and pathological observations.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

始めに: アホロートル（ウーパールーパー）の 尻尾が反り返る（反り上がる） のは、よく観察される「体調不良のサイン」のひとつです。以下のような原因が考えられます。

主な原因

1. **水質の悪化**
   - アンモニア、亜硝酸、硝酸塩の蓄積や pH の急変が大きなストレスに。
   - 水質が悪いと、まずエラの縮みや血流不良が起こり、尾の反り返りなど異常姿勢が出やすい。
2. **水温の上昇**
   - アホロートルは低水温（16–20℃前後）を好む。
   - 25℃を超える環境では代謝が上がりすぎてストレス状態になり、体のコントロールが乱れることがある。
3. **低酸素状態**
   - 酸素不足やフィルターの不調でストレスが高まり、神経筋に異常な反応が出ることがある。
4. **栄養不足・代謝異常**
   - カルシウム不足、ビタミン欠乏、長期の偏食などが神経・筋肉の異常につながる可能性。
5. **感染症や中毒**
   - **細菌感染や真菌感染、または水槽内の化学物質（塩素、金属、薬剤）によって神経・筋肉系に影響が出る。**
   - 内臓の異常や中毒でも姿勢の異常が見られる。
6. **ストレスや環境要因**
   - 他の個体との同居によるストレス、急な照明、水流が強すぎる環境なども関与する。

まとめ

- **尻尾の反り返りは「異常のサイン」**。
- 特に多いのは **水質悪化や水温上昇**。
- 続く場合は感染症や栄養不良の可能性も考えるべき。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

参考文献・情報源　References

出典	内容	尻尾が反り返る原因と関係する点
Ambystoma Genetic Stock Center (University of Kentucky) — FAQ: *“We house axolotls for research purposes. Several of these have developed scoliosis … most appear in the base of the tail … nutritional, … calcium deficiency”* ambystoma.uky.edu	研究施設での飼育アホロートルでの尾基部の尾の偏向、脊柱変形の記録	栄養不良、特にカルシウムの欠乏が原因の可能性が示唆されている。環境や発育の問題。
Caudata Culture Articles — Why Larvae Die	両生類幼生で“curled tail”（尾が丸まる／くるっと反る尾先など）を死亡直前の一般的な症状として記載 caudata.org	尾の先端や尾部分の湾曲は強いストレスや水質悪化、感染症などと関連。
Amphibian survival, growth and development in response to elevated ammonium (Griffis-Kyle KL et al., 2007) PubMed	両生類幼生をアンモニウム高濃度下で育てた実験。発育遅延、形態異常など多数報告	水中のアンモニア／アンモニウム濃度が高いと形態異常を含む様々なストレス症状が見られる可能性。尾の湾曲なども含む可能性あり。
Direct effect of ammonia on three species of north … (Wiley, 2009) ワイリーオンラインライブラリー	アンモニア曝露が両生類に与える影響：孵化率・発育など	水質のアンモニアが発育異常を引き起こすという証拠。尾の反りも発育異常の一種と見なせる場合がある。
Axolotl Planet: “The #1 Guide to Axolotl Sickness and Health”	ペット飼育者向けガイド。「もし尾が異常に反っているなら、水質とアンモニア・窒素サイクルをすぐチェックせよ」と助言 Axolotl Planet	実務的な飼育経験で、尾の湾曲がストレス・水質問題のサインとして扱われている。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

アホロートル尾部の反り返りと抗酸菌症の関連性

1. **非特異的ストレス症状としての尾の反り返り**
    - 文献や飼育レポートでは、尾の湾曲はしばしば **全身状態の悪化サイン** とされます。
    - 水質悪化やアンモニア、低栄養などが原因となりますが、**感染症による慢性消耗** も同じように尾の異常姿勢を引き起こすことがあります。
2. **抗酸菌症と体形異常**
    - 両生類や魚類での抗酸菌感染（*Mycobacterium marinum*, *M. chelonae* など）は、慢性の経過を取り、**痩せ・沈うつ・浮腫・局所の肉芽腫** を伴います。
    - 長期にわたる慢性炎症は、筋肉や脊椎周辺の組織に影響を与え、尾の動きや形状に異常をきたすことがあります。
3. **水質・免疫・感染の三角関係**
    - アホロートルは免疫抑制に弱く、**水質ストレス → 免疫低下 → 抗酸菌感染** という流れが起こりやすい。
    - 反り返る尾は「水質ストレス」のサインであり、その背景に**抗酸菌症のような慢性疾患**が隠れている可能性もある。

参考文献

- **Whipps, C. M., & Kent, M. L. (2006).** "Mycobacteriosis in fishes: a review." *Veterinary Journal*, 171(3), 294–309.
  → 魚類での抗酸菌症の病態を整理。肉芽腫や体形異常を引き起こすことがある。
- **Green, S. L. (2001).** "Amphibian Medicine and Captive Husbandry." Krieger Publishing.
  → 両生類における抗酸菌症の記載。慢性消耗性疾患であり、体形の異常も起こり得る。
- **Ambystoma Genetic Stock Center FAQ**（University of Kentucky）
  → 尾の変形は栄養・代謝異常に加え、感染も原因のひとつとされる。

まとめ

- 尾の反り返り＝必ずしも抗酸菌症ではないが、**水質・栄養・免疫異常 → 二次的に抗酸菌感染が顕在化**するケースを考えられる。
- 特に慢性経過や痩せ、皮膚病変、結節を伴う場合は、**抗酸菌症を疑う一因** となる。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Introduction: In axolotls (Ambystoma mexicanum), a tail that bends upward (tail curling) is a commonly observed sign of poor health. The following causes are often considered.

Main Causes

1. Poor Water Quality

- Accumulation of ammonia, nitrite, nitrate, or sudden pH shifts causes severe stress.
- When water quality deteriorates, gill shrinkage and impaired blood circulation often occur first, followed by abnormal postures such as tail curling.

2. Elevated Water Temperature

- Axolotls prefer cool water (around 16–20 °C).
- At temperatures above 25 °C, metabolism increases excessively, causing stress and loss of bodily control.

3. Low Oxygen Levels

- Oxygen depletion or filter malfunction increases stress, which may trigger abnormal neuromuscular responses.

4. Nutritional Deficiency / Metabolic Disorder

- Calcium deficiency, vitamin shortages, or long-term imbalanced diets may cause neuromuscular abnormalities.

5. Infections or Toxicity

- Bacterial or fungal infections, or exposure to chemicals in the tank (chlorine, metals, medications), may affect the neuromuscular system.
- Organ dysfunction or poisoning may also manifest as abnormal posture.

6. Stress and Environmental Factors

- Stress from cohabitation with other individuals, sudden changes in lighting, or excessive water flow can also contribute.

Summary

- Tail curling is a **warning sign of abnormality**.
- The most frequent causes are poor water quality and high temperature.
- If the condition persists, infection or nutritional deficiency should be considered.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

References / Sources

Source	Content	Relevance to Tail Curling
Ambystoma Genetic Stock Center (University of Kentucky) — FAQ: “We house axolotls for research purposes. Several of these have developed scoliosis … most appear in the base of the tail … nutritional, … calcium deficiency” (ambystoma.uky.edu)	Records of tail deformities and spinal abnormalities in research axolotls	Suggests nutritional deficiency, especially calcium, as a possible cause; also linked to environment and development
Caudata Culture Articles — Why Larvae Die (caudata.org)	Notes “curled tail” as a general pre-death symptom in amphibian larvae	Associates tail curling with stress, poor water quality, and infections
Griffis-Kyle KL et al., 2007. Amphibian survival, growth and development in response to elevated ammonium. PubMed	Experimental exposure of amphibian larvae to high ammonium; reported developmental delays and deformities	High ammonium levels in water can cause stress symptoms including body deformities and potentially tail curling
Wiley Online Library (2009). Direct effect of ammonia on three species of north …	Effects of ammonia exposure on amphibians: hatching rates, development, deformities	Evidence that ammonia in water induces developmental abnormalities, including tail curling
Axolotl Planet — “The #1 Guide to Axolotl Sickness and Health”	Husbandry guide for pet owners; advises immediate water quality and nitrogen cycle checks if the tail appears abnormally curled	Based on practical husbandry experience, tail curling is treated as a sign of stress or water quality problems

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Relationship Between Axolotl Tail Curling and Mycobacteriosis

1. **Non-specific Stress Symptom**
    - Literature and husbandry reports describe tail curling as a **general indicator of declining health**.
    - While poor water quality, ammonia, or malnutrition are common causes, chronic wasting from infection can also result in abnormal posture.
2. **Mycobacteriosis and Body Deformities**
    - In amphibians and fish, infections by *Mycobacterium marinum*, *M. chelonae*, and related species follow a **chronic course**, with emaciation, lethargy, edema, and localized granulomas.
    - Prolonged inflammation can affect muscles and tissues around the spine, leading to tail movement or shape abnormalities.
3. **Water Quality – Immunity – Infection Triangle**
    - Axolotls are particularly vulnerable to immunosuppression.
    - **Water quality stress → lowered immunity → opportunistic mycobacterial infection** is a plausible sequence.
    - Tail curling may thus signal water quality stress, behind which a chronic condition such as mycobacteriosis could be hidden.

Additional References

- Whipps, C. M., & Kent, M. L. (2006). *Mycobacteriosis in fishes: a review.* **Veterinary Journal, 171(3), 294–309.**
  → Summarizes pathology of mycobacteriosis in fish; granulomas and body deformities can occur.
- Green, S. L. (2001). *Amphibian Medicine and Captive Husbandry.* Krieger Publishing.
  → Includes descriptions of mycobacteriosis in amphibians; noted as a chronic wasting disease that may involve body deformities.
- Ambystoma Genetic Stock Center FAQ (University of Kentucky).
  → Notes tail deformities can be caused by nutritional or metabolic disorders, but also by infections.

Final Note

- Tail curling **does not necessarily indicate mycobacteriosis**, but water quality, nutrition, and immune problems may predispose axolotls to opportunistic mycobacterial infections.
- Particularly in chronic cases with emaciation, skin lesions, or nodules, **mycobacteriosis should be considered as a differential diagnosis**.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

抗酸菌症（Mycobacteriosis）とは

1. 定義と病原体

抗酸菌症（Mycobacteriosis）は、**抗酸菌（Mycobacterium 属）** によって引き起こされる慢性・進行性の感染症の総称です。抗酸菌は細胞壁にミコール酸を含むため、酸性アルコールで脱色されにくい「抗酸性」を示し、**Ziehl–Neelsen 染色**や**蛍光染色（Auramine-O）**で特徴的に観察されます。

代表的な病原種:

- ヒト・哺乳類：*M. tuberculosis complex*（M. tuberculosis, M. bovis など）、*M. leprae*, 非結核性抗酸菌（NTM: M. avium complex, M. kansasii, M. abscessus など）
- 魚類・両生類：*M. marinum*, *M. fortuitum*, *M. chelonae* など

---

2. ヒトにおける抗酸菌症

主な疾患

- **結核（Tuberculosis, TB）**：M. tuberculosis による世界的主要感染症。肺結核を中心に、髄膜炎・骨関節結核・腎結核など全身性疾患を起こす。
- **らい病（Leprosy）**：M. leprae による末梢神経障害・皮膚病変。
- **非結核性抗酸菌症（NTM 感染症）**：M. avium complex (MAC), M. abscessus など。慢性呼吸器感染、リンパ節炎、皮膚感染（“水槽肉芽腫”）を引き起こす。

特徴

- 慢性進行性で治療に時間を要する。
- 多剤耐性菌が問題。
- HIV 感染者や免疫不全者では高リスク。

---

3. 家畜における抗酸菌症

主な病型

- **ウシ結核（Bovine TB, M. bovis）**：ウシを中心に、ブタ・シカ・野生動物へも感染。ヒトへの人獣共通感染症リスクあり。
- **鳥型抗酸菌症（Avian mycobacteriosis, M. avium subsp. avium）**：ニワトリ・七面鳥・インコ類などで慢性消耗性疾患。

特徴

- 慢性消耗、体重減少、結節性病変（リンパ節・肝・脾）。
- 家畜衛生上、検査・淘汰・消毒が義務づけられている。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. 魚類における抗酸菌症

病原体

- 主に *M. marinum*, *M. fortuitum*, *M. chelonae*。

臨床像

- 慢性消耗（痩せる）
- 皮膚潰瘍、鱗の脱落、鰭の壊死
- 内臓結節（肝・脾・腎）

特徴

- 「魚結核（Fish tuberculosis）」とも呼ばれる。
- ヒトにおいては“水槽肉芽腫（Fish tank granuloma）”の原因となり、魚からの人獣共通感染に注意。

---

5. 両生類における抗酸菌症

報告例

- アホロートル（Ambystoma mexicanum）、カエル類などで *M. marinum*, *M. fortuitum* 感染例。
- 痩せ、皮膚潰瘍、眼球突出、内臓結節が見られる。

特徴

- 感染しても長期間無症状の個体が存在。
- 症状進行後は栄養失調・消耗により予後不良。
- 飼育・研究施設での拡散リスクが高く、器具・水槽の消毒が必須。

---

6. 感染経路と病理学的特徴

- **経口感染**：汚染餌・水からの摂取。
- **経皮感染**：外傷や鰓・皮膚からの侵入。
- **病理像**：類上皮細胞や多核巨細胞を伴う肉芽腫形成。中心壊死や線維化も。

---

7. 消毒と管理

- 次亜塩素酸ナトリウム（200–500 ppm, 30 分以上）
- 熱処理（100°C, 5 分以上）
- 乾燥（60°C 以上数時間）
- 紫外線殺菌（補助的手段）

※ 抗酸菌は細胞壁が強固で耐性が高いため、十分な濃度・時間が必要。

まとめ

- 抗酸菌症は **ヒト・家畜・魚類・両生類に共通して発生する人獣共通感染症** であり、慢性消耗性疾患を引き起こす。
- 病原体は種によって異なるが、共通して「痩せる・潰瘍・結節形成」という臨床像が見られる。
- 種同定は困難な場合が多いが、感染が疑われた段階で **消毒・隔離・適切な廃棄** を徹底することが推奨される。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Mycobacteriosis in Humans, Livestock, Fish, and Amphibians

1. Definition and Pathogens

Mycobacteriosis refers to a group of chronic and progressive infections caused by **acid-fast bacteria of the genus Mycobacterium**. These organisms possess a lipid-rich cell wall containing mycolic acids, which confer resistance to acid-alcohol decolorization. Consequently, they are characteristically detected by **Ziehl–Neelsen staining** or **auramine-O fluorescence staining**.

Major pathogens:

- **Humans and mammals**: *Mycobacterium tuberculosis complex* (*M. tuberculosis, M. bovis*), *M. leprae*, and nontuberculous mycobacteria (NTM) such as *M. avium complex, M. kansasii, M. abscessus*.
- **Fish and amphibians**: *M. marinum, M. fortuitum, M. chelonae*.

---

2. Mycobacteriosis in Humans

Major diseases

- **Tuberculosis (TB)**: Caused by *M. tuberculosis*, a major global health threat. Affects mainly the lungs but can involve meninges, bones, kidneys, and other organs.
- **Leprosy (Hansen's disease)**: Caused by *M. leprae*, leading to peripheral neuropathy and skin lesions.
- **Nontuberculous mycobacteriosis (NTM disease)**: Caused by *M. avium complex, M. abscessus*, etc. Manifestations include chronic pulmonary disease, lymphadenitis, and cutaneous infections such as "fish tank granuloma."

Characteristics

- Chronic and progressive course requiring long-term therapy.
- Emergence of multidrug-resistant strains is a major problem.
- High-risk in immunocompromised hosts, including HIV patients.

---

3. Mycobacteriosis in Livestock

Major conditions

- **Bovine tuberculosis (*M. bovis*)**: Affects cattle, pigs, deer, and wildlife. Recognized as a zoonotic infection transmissible to humans.
- **Avian mycobacteriosis (*M. avium* subsp. *avium*)**: Occurs in chickens, turkeys, and parrots, causing chronic wasting disease.

Characteristics

- Progressive emaciation, granulomatous lesions in lymph nodes, liver, and spleen.
- Strict veterinary hygiene measures: testing, culling, and disinfection.

---

4. Mycobacteriosis in Fish

Pathogens

- *M. marinum, M. fortuitum, M. chelonae*.

Clinical signs

- Chronic emaciation ("wasting").
- Skin ulcers, scale loss, fin necrosis.
- Visceral granulomas (liver, spleen, kidney).

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Characteristics

- Often referred to as **“fish tuberculosis”**.
- Zoonotic potential: causes **fish tank granuloma** in humans, typically via skin abrasions.

---

5. Mycobacteriosis in Amphibians

Reported cases

- Axolotls (*Ambystoma mexicanum*) and frogs have been reported to be infected by *M. marinum* and *M. fortuitum*.
- Clinical signs: weight loss, skin ulcers, exophthalmos, visceral granulomas.

Characteristics

- Some individuals remain asymptomatic for long periods.
- Once clinical disease progresses, prognosis is poor due to malnutrition and tissue degeneration.
- High risk of spread in laboratory or breeding facilities, necessitating strict disinfection of water tanks and equipment.

---

6. Transmission Routes and Pathology

- **Oral route**: ingestion of contaminated feed or water.
- **Percutaneous route**: entry through wounds, skin, or gills.
- **Histopathology**: granuloma formation with epithelioid cells, multinucleated giant cells, caseous necrosis, and fibrosis.

---

7. Disinfection and Control

- **Sodium hypochlorite**: 200–500 ppm available chlorine, ≥30 minutes.
- **Boiling water**: 100 °C for ≥5 minutes.
- **Drying**: ≥60 °C for several hours.
- **Ultraviolet irradiation**: for reducing bacterial load in water (not sufficient for surface disinfection).

Note: Mycobacteria are highly resistant due to their lipid-rich cell wall, requiring adequate concentration and contact time for inactivation.

Conclusion

- Mycobacteriosis is a **chronic zoonotic disease** that affects humans, livestock, fish, and amphibians.
- Despite differences in species and causative agents, common clinical features include **emaciation, ulceration, and granuloma formation**.
- Species-level identification of pathogens is often difficult, especially in fish and amphibians, but once mycobacteriosis is suspected, **disinfection, isolation, and proper disposal** should be implemented immediately.
- This is considered an essential **risk management measure** to protect both animal and human health.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

参考文献　References

1. Sasaki M. *Sterilizing activity of povidone-iodine against tubercle bacilli*. Chemotherapy, 1981; 29(8):880–885.
2. Olivier G. *Mycobacteriosis in fish: a review*. Veterinary Bulletin, 1992; 62:1–30.
3. Decostere A, Hermans K, Haesebrouck F. *Mycobacteriosis in fish: a review*. Veterinary Microbiology, 2004; 99(3–4):159–166.
4. Falkinham JO. *Epidemiology of infection by nontuberculous mycobacteria*. Clin Microbiol Rev, 1996; 9(2):177–215.
5. Jacobs JM, Stine CB, Baya AM, Kent ML. *A review of mycobacteriosis in marine fish*. J Fish Dis, 2009; 32(2):119–130.
6. Shioda C, Uchida K, Nakayama H. *Pathological features of olfactory neuroblastoma in an axolotl (Ambystoma mexicanum)*. J Vet Med Sci, 2011; 73(11):1531–1534.
7. Harshbarger JC, Chang SC, DeLanney LE, Rose FL, Green DE. *Cutaneous mastocytomas in neotenic salamanders (Ambystoma mexicanum and A. tigrinum)*. J Cancer Res Clin Oncol, 1999; 125(1):57–61.
8. Decostere A, et al. *Disinfection and control strategies against mycobacteria in aquaculture*. Aquaculture, 2004; 240(1–4):19–30.
9. World Health Organization (WHO). *Global tuberculosis report 2023*. Geneva: WHO.
10. Ministry of Health, Labour and Welfare (Japan). *Guidelines for zoonotic diseases*. 2019.
11. Fisheries Agency of Japan. *Manual for fish disease management*. 2020.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

魚類の抗酸菌症（Mycobacteriosis）

- 起因菌：Mycobacterium marinum, M. fortuitum, M. chelonae など。
- 特徴：慢性的に魚がやせる、皮膚潰瘍、眼の突出、鰭の損耗など。
- ヒトにも感染し（いわゆる「水槽肉芽腫」）、**人獣共通感染症（ズーノーシス）**として扱われています。

厚労省のガイドラインの位置づけ

- 厚生労働省や水産庁の文書では、**魚類抗酸菌症は人獣共通感染症に分類**され、
  水槽・器具・排水の取り扱いには注意が必要とされています。
- 「必ずこの方法で消毒せよ」と一律に義務づけているわけではなく、
  感染源の拡散防止とヒト感染予防の観点から強く推奨されている という位置づけです。
- 実際のガイドラインでは以下が記載されています：
  - 使用した水槽や器具は **次亜塩素酸ナトリウム（漂白剤）や熱湯、乾燥などで消毒**。
  - 飼育水や死魚はそのまま下水に流さず、**消毒処理後に廃棄**。
  - 感染魚の治療は困難なため、**隔離または安楽死処分が推奨**されることも多い。

実務的な意味

- **法的に強制される義務ではない**（例：届出義務疾病ではない）。
- しかし、
  1. 他の魚への水平感染を防ぐ
  2. ヒト（飼育者）の皮膚感染を防ぐ
     この 2 点から、**消毒は実質的に必須に近い対応**と考えられます。

まとめ

- 魚の抗酸菌症は **法的義務ではなく「強く推奨される対応」**。
- 厚労省のガイドラインは「必ず消毒せよ」という命令ではなく、**人獣共通感染症リスクを防ぐための安全対策**。
- 魚や器具をそのまま放置・廃棄すると感染拡散や人へのリスクが残るため、**現実的には消毒を行うべきです**。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

両生類における抗酸菌症とその対策

1. 概要

両生類における抗酸菌症は、**抗酸菌属（Mycobacterium）の酸抗性菌**によって引き起こされる慢性感染症です。原因菌には *M. marinum*, *M. fortuitum*, *M. chelonae* などが含まれます。発生頻度は魚類や哺乳類に比べると少ないものの、**アホロートル（*Ambystoma mexicanum*）**、カエル類（*Rana pipiens*, *Xenopus spp.*）などで報告されています。

2. 臨床症状

- **消耗症候群（やせ）**：初期には食欲があるものの、次第に痩せ細っていく。
- **皮膚病変**：潰瘍、結節、色素消失。
- **眼症状**：重症例では眼球突出（exophthalmos）がみられる。
- **内臓病変**：肝臓、脾臓、腎臓などに肉芽腫性炎症。
- **経過**：慢性的に進行し、進行例では予後不良。無症状の保菌個体も存在。

3. 病態

- **感染経路**：
  - 汚染された生餌や水の摂取
  - 傷口や鰓からの侵入
- **病理学的特徴**：上皮様細胞や多核巨細胞からなる肉芽腫形成、壊死巣。
- **宿主応答**：再生能の高いアホロートルなどでは進行が遅れる場合もあるが、発症を防ぐことは困難。

4. 診断

- **臨床的疑い**：やせ、皮膚潰瘍、眼の異常など。
- **病理組織**：Ziehl–Neelsen 染色や Fite 染色により肉芽腫内の抗酸菌を検出。
- **微生物学的検査**：培養や PCR 同定は可能だが、一般施設では困難。
- **実際**：多くの症例は臨床像と組織所見から「抗酸菌症疑い」と診断される。

5. 対策・管理

- **隔離**：感染・疑い個体はすぐに隔離。
- **消毒**：
  - **次亜塩素酸ナトリウム**：有効塩素 200〜500 ppm、30 分以上浸漬。
  - **熱湯処理**：100℃で 5 分以上（小型器具向き）。
  - **乾燥**：60℃以上で数時間。日光乾燥も有効。
  - **ポビドンヨード (PVP-I, イソジン)**：in vitro では数秒〜数分で抗酸菌を不活化。薬浴応用には安全性確認が必要。
- **淘汰・廃棄**：重症や回復困難な個体は安楽死とし、焼却または消毒後に廃棄。
- **環境管理**：
  - 未処理の生餌使用を避ける（抗酸菌の保菌リスクあり）。
  - 生餌は塩水浴や薬剤処理でトリートメント。
  - 飼育水は廃棄前に消毒。

6. 公衆衛生・人獣共通感染症の懸念

- *M. marinum* などはヒトにも感染し、「水槽肉芽腫（fish tank granuloma）」を引き起こす。
- 特に免疫不全者はリスクが高いため、飼育者は手袋を使用し、器具の定期消毒を徹底する必要がある。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

7. まとめ

両生類の抗酸菌症は重篤で多くの場合致死的であり、動物福祉・研究・保全活動に大きな影響を与える。また人獣共通感染症としてヒトへの感染リスクもある。菌種レベルでの同定は難しいため、**疑いの時点で抗酸菌症として管理する**ことが推奨される。早期発見・迅速な隔離・徹底した消毒が発生拡大を防ぐ鍵となる。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

Mycobacteriosis in Amphibians and Its Control

1. Overview

Mycobacteriosis in amphibians is a chronic infectious disease caused by **acid-fast bacteria of the genus Mycobacterium**, including *M. marinum*, *M. fortuitum*, and *M. chelonae*. Although relatively rare compared to fish and mammals, cases have been reported in **axolotls (*Ambystoma mexicanum*)**, frogs (*Rana pipiens*, *Xenopus spp.*), and other amphibians.

2. Clinical Features

- **Wasting syndrome**: Progressive emaciation despite preserved appetite in the early stages.
- **Cutaneous manifestations**: Ulcerative lesions, nodules, or loss of skin pigmentation.
- **Ocular signs**: Exophthalmos (eye protrusion) reported in severe infections.
- **Visceral lesions**: Granulomatous inflammation in liver, spleen, and kidneys.
- **Course of disease**: Chronic, often with poor prognosis once advanced. Some animals remain asymptomatic carriers.

3. Pathogenesis

- **Transmission routes**:
  - Ingestion of contaminated live feed or water.
  - Entry through skin wounds or gills.
- **Histopathology**: Granuloma formation with epithelioid cells, multinucleated giant cells, and necrotic tissue.
- **Host response**: In amphibians with high regenerative capacity (e.g., axolotls), the immune response may delay but not prevent disease progression.

4. Diagnostic Approach

- **Clinical suspicion**: Wasting, ulcers, ocular changes.
- **Histopathology**: Acid-fast bacilli detected in granulomatous lesions by Ziehl–Neelsen or Fite staining.
- **Microbiology**: Culture and PCR-based identification are possible but rarely successful in standard laboratory settings.
- **Practical diagnosis**: Most cases are labeled as “suspected mycobacteriosis” based on symptoms and histology.

5. Control and Management

- **Isolation**: Infected or suspected individuals should be immediately separated to prevent spread.
- **Disinfection**:
  - **Sodium hypochlorite**: 200–500 ppm, ≥30 minutes immersion of equipment.
  - **Boiling water**: 100 °C for ≥5 minutes for small instruments.
  - **Drying**: ≥60 °C for several hours; effective in reducing bacterial survival.
  - **Povidone-iodine (PVP-I)**: Reported to inactivate *Mycobacterium* spp. within seconds to minutes in vitro; potential use in water baths requires safety validation for amphibians.
- **Culling and disposal**: Severely affected or moribund individuals should be euthanized humanely, and carcasses disposed of via incineration or chemical disinfection.
- **Environmental management**:
  - Avoid use of untreated live feed (possible carriers).
  - Treat live feed by saltwater baths or chemical rinses.
  - Replace or disinfect contaminated tank water before disposal.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

6. Public Health and Zoonotic Concern

- Mycobacteria such as *M. marinum* are zoonotic and can cause **“fish tank granuloma”** in humans, typically via hand wounds.
- Amphibian handlers, especially those with immunosuppression, should wear gloves and disinfect equipment regularly.

7. Conclusion

Mycobacteriosis in amphibians is a serious, often fatal disease with implications for animal health, conservation programs, and zoonotic safety. Because species-level identification of causative mycobacteria is difficult, **any suspected case should be managed as mycobacteriosis** with strict hygiene, disinfection, and isolation measures. Preventive practices, early detection, and rapid intervention are critical to mitigating outbreaks in amphibian colonies.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

抗酸菌に対するメチレンブルーの効果

1. **酸性菌（acid-fast bacteria, Mycobacterium spp.）の染色法におけるメチレンブルーの役割**
    - Ziehl–Neelsen 染色や Kinyoun 染色などでは主染色 (primary stain) に **カーボルフクシン**、脱色後の対染色 (counterstain) として **メチレンブルー (methylene blue)** が使われます。脱色工程後、抗酸菌 (acid-fast bacteria) は赤 (carbol-fuchsin) を保持するため、背景または非抗酸菌がメチレンブルーに染まることで対比がつきます。 Leica Biosystems+2Microbiology Info.com+2
    - このことから、「メチレンブルーは抗酸菌を染める／主染色するための染料ではない」という点がわかります。すなわち、抗酸菌はメチレンブルー単独では赤く染まらず、対染色で背景色を提供するだけの用途。つまり、「メチレンブルー単独では抗酸菌の染色に十分ではない」という意味で「効かない」という表現が部分的に当てはまります。 NCBI+2Microbiology Info.com+2
2. **メチレンブルーを光増感剤（photosensitizer）として使う実験**
    - *Mycobacterium smegmatis* を対象とした研究で、メチレンブルー (MB) を用いたフォトダイナミック療法 (PDT: Photodynamic Therapy) が行われており、光を照射することで殺菌効果を発揮する例があります。つまり、「暗所」や「光なし」では殺菌効果が弱い／ほぼない、ということ。 ResearchGate
    - また、MB の濃度や照射時間・光源の有無などの条件に依存して効果が変わるという報告があることから、「メチレンブルーのみでは抗酸菌を確実に死滅させるとは限らない」という実験的証拠があります。 ResearchGate

---

⚠ 限界・注意点

- 「効かない」という表現はあいまい。抗酸菌における「染色が十分にされない」「殺菌が弱い／条件依存」などが多く、「完全に無効である」とするデータは見当たらない。
- フォトダイナミック療法で使うメチレンブルーは **光あり／一定の条件あり** の実験であり、通常の染色や暗所での処置とは状況が異なる。
- 多くの染色プロトコールでは、メチレンブルーは主染色ではなく対染色として使われるため、「メチレンブルーが抗酸菌を赤く染める」ことを期待するプロトコールはそもそも少ない。

結論

- メチレンブルー単独では、抗酸菌（Mycobacterium 等）の染色または殺菌に十分ではない証拠が複数ある。特に染色では、メチレンブルーは対染色（背景染色）として使われ、抗酸菌自体は主染色＋脱色工程後に色を保持する染料で染まる（カーボルフクシンなど）。
- メチレンブルーを光増感剤として使うときには、条件次第で抗酸菌の殺菌に効果があるという報告もある。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Effects of Methylene Blue on Mycobacteria

1. Role of Methylene Blue in Staining Acid-Fast Bacteria (Mycobacterium spp.)

- In staining methods such as **Ziehl–Neelsen** or **Kinyoun** staining, carbol fuchsin is used as the **primary stain**, and methylene blue is applied as the **counterstain** after decolorization. After the decolorization step, acid-fast bacteria (AFB) retain the red color of carbol fuchsin, while the background or non–acid-fast bacteria are stained blue by methylene blue, providing contrast.
- From this, it is clear that **methylene blue is not used to directly stain acid-fast bacteria**. In other words, acid-fast bacteria do not appear red with methylene blue alone; methylene blue only serves to provide a background color. Therefore, the statement that “methylene blue does not work” partially applies in the sense that **it is not sufficient on its own to stain acid-fast bacteria**.

---

2. Use of Methylene Blue as a Photosensitizer

- In experiments with *Mycobacterium smegmatis*, methylene blue (MB) has been used in **photodynamic therapy (PDT)**, where bactericidal effects are induced upon exposure to light. This means that **in the absence of light**, MB shows weak or almost no bactericidal activity.
- Reports indicate that the bactericidal effect depends on MB concentration, exposure time, and the presence of a light source. These findings support the conclusion that **methylene blue alone does not necessarily kill mycobacteria reliably**.

⚠ Limitations and Considerations

- The expression “does not work” is ambiguous. In many cases, the issue is that methylene blue provides **insufficient staining or weak/condition-dependent bactericidal effects**, rather than being completely ineffective.
- In PDT studies, methylene blue works only **under light exposure** and under specific experimental conditions, which differ from standard staining or dark treatments.
- In most staining protocols, methylene blue is applied as a **counterstain**, not a primary stain. Therefore, protocols rarely expect methylene blue to stain acid-fast bacteria themselves.

Conclusion

- Multiple pieces of evidence show that **methylene blue alone is insufficient for staining or killing Mycobacterium spp.**. Specifically, in staining, methylene blue serves as a counterstain for background contrast, while acid-fast bacteria retain carbol fuchsin after the decolorization step.
- When used as a **photosensitizer**, methylene blue can have bactericidal effects against mycobacteria under certain conditions, highlighting that its effectiveness is context-dependent.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

抗酸菌に対するイソジン（ポピドンヨード: PVPI）の効果

参考文献: https://www.jstage.jst.go.jp/article/chemotherapy1953/29/8/29_8_880/_article/-char/ja/

イソジン（ポビドン・ヨード: Povidone-Iodine, PVP-I, Isodine 等）の抗酸菌（Mycobacterium 属）への滅菌力・薬浴治療等での有用性を整理します。

主な研究結果

文献	抗酸菌への作用／条件	結果の要点
佐々木昌子, 1981（「消毒剤 Povidone-Iodine の結核菌に対する殺菌力」） J-STAGE	BCG, H37Rv など結核菌株に対し、100 mg/mL の PVP-I 使用。5 秒間の接触で BCG を殺菌。血清添加や低温（10°C）では効果低下。	非常に短時間で殺菌可能であり、手洗い用などでの使用が実践的とされる。ただし条件（温度、有機物の存在など）により大きく変動。
Rikimaru et al., 1997 PubMed	複数の Mycobacterium 標準株および臨床分離株に対して市販の PVP-I 溶液あるいは PVP-I 薬剤本体 (drug substance) を 0.1% または 0.02% で使用。30〜120 秒間の接触時間、2% ヒト血清添加あり。	0.1% の PVP-I で 30 秒以内に標準株を完全不活化。臨床株も 60 秒以内でほぼ全滅。つまり低濃度でも比較的短時間で強い殺菌力あり。
Mycobacterium fortuitum に対する PVP-I を含む消毒薬の実験 ResearchGate	PVP-I を含む市販消毒薬（希釈条件下、水硬度・有機物の影響を含む）を 30 分間作用させる条件で試験。	少なくとも *M. fortuitum* に対して優れた殺菌活性を示す。硬水・有機物存在下でも希釈条件を変えることで有効。

条件・制限事項

- **温度**：低温では殺菌効果が低くなる（例：10°C では著しく減少） J-STAGE。
- **有機物**（血清添加など）の存在は PVP-I の作用を弱める要因。ヒト血清添加で効果が低下する報告あり。 J-STAGE+1
- **濃度／接触時間**が重要。0.1% の濃度で 30 秒以上の接触が必要な場合が多い。濃度を落としたり短時間だと不十分。 PubMed

---

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

薬浴治療への有用性・応用可能性

これらの in vitro（試験管内）の結果を魚類／両生類の薬浴に応用する際のポイントと可能性は以下です：

利点	条件・注意点
短時間で殺菌作用を示すことから、薬浴治療での“初期段階”における感染抑制や進行遅延に寄与する可能性が高い。	魚類・両生類に対する安全性（皮膚、鰓への刺激性、毒性）を確認する必要。特に PVP-I はヨウ素成分が強いため、過量・長時間の曝露は動物に害を与える可能性がある。
種によらず多様な Mycobacterium 株に対して作用を示すため、不明株（種同定できない株）にも“疑い治療”として使いやすい。	水環境（温度、有機汚れ、pH、塩分など）が in vitro 条件と異なるため、それらが効果を減弱させる可能性。実際に魚／両生類での in vivo 実験が少ない。
感染の初期（体重減少や摂食率低下の初期症状）を察知した時点で迅速に処置すれば、回復を促す可能性あり。	症状が進んで腸萎縮や組織壊死が進んだ個体では、PVP-I の薬浴だけでは回復が難しいことが予想される。

⚠ 日本での文献に基づく注意点

- 佐々木 (1981) の研究はヒト（結核菌）に対するものであり、魚類・両生類では組織や体表の構造が異なるため、同じ効果が得られるとは限らない。
- 安全性（動物毒性、鰓や皮膚の損傷）に関するデータが不足しているため、初期データを小規模に試す必要あり。

結論（In vitro → in vivo の橋渡しとして）

- イソジン（PVP-I, ポビドンヨード／Isodine）は、in vitro で Mycobacterium 属に対して短時間・低濃度でも殺菌効果を示す信頼できる消毒薬である。
- 初期症状の魚／両生類に対して薬浴的に適用すれば、感染進行を抑え、回復の可能性を高めることが期待される。
- ただし、以下が必要：
    1. 適切な濃度と接触時間の確立
    2. 動物への毒性を含む安全性評価（例えば皮膚 / 鰓刺激性など）
    3. 水中の有機物・温度・pH・水硬度など環境条件を考慮した実用プロトコールの構築

※各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

The Effect of Povidone-Iodine (PVP-I, Isodine) on Mycobacteria and Its Potential for Bath Treatment

Key Research Findings

Reference	Effect on Mycobacteria / Conditions	Key Results
Sasaki, 1981 (“Bactericidal effect of disinfectant Povidone-Iodine on Mycobacterium tuberculosis”) J-STAGE	Tested on BCG, H37Rv and other M. tuberculosis strains with 100 mg/mL PVP-I. BCG was killed within 5 seconds of exposure. Efficacy decreased under low temperature (10°C) or in the presence of serum.	Demonstrated extremely rapid bactericidal activity, making it practical for handwashing use. However, the effect varies greatly depending on conditions (temperature, organic matter).
Rikimaru et al., 1997 PubMed	Tested on multiple Mycobacterium reference and clinical strains. Commercial PVP-I solution or drug substance at 0.1% or 0.02%, with exposure times of 30–120 seconds, in the presence of 2% human serum.	0.1% PVP-I completely inactivated reference strains within 30 seconds. Most clinical strains were killed within 60 seconds. Indicates strong bactericidal activity even at low concentrations in a relatively short time.
Experiment on Mycobacterium fortuitum with PVP-I containing disinfectant (ResearchGate)	Commercial disinfectants containing PVP-I, tested under diluted conditions, including water hardness and organic material effects, with 30 minutes exposure.	Showed excellent bactericidal activity against *M. fortuitum*. Effective even under hard water or with organic matter when dilution conditions were adjusted.

Conditions and Limitations

- **Temperature**: Low temperatures significantly reduce bactericidal effect (e.g., 10°C markedly decreases activity).
- **Organic matter**: Presence of organic material (e.g., serum) weakens the efficacy of PVP-I.
- **Concentration / Exposure time**: Adequate concentration and time are critical. Typically, 0.1% PVP-I for at least 30 seconds is needed. Lower concentrations or shorter exposure may be insufficient.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Applicability for Bath Treatment in Fish and Amphibians

Advantage	Conditions / Considerations
Rapid bactericidal activity suggests potential benefit for suppressing infection or delaying disease progression at the *early stage* of infection.	Safety must be confirmed for fish and amphibians (skin, gill irritation, systemic toxicity). As iodine is strong, excessive or prolonged exposure may be harmful.
Effective against diverse Mycobacterium species, making it useful as an “empirical treatment” even when the strain cannot be identified.	Environmental water conditions (temperature, organic load, pH, salinity) differ from in vitro settings, potentially reducing efficacy. In vivo data for fish/amphibians are limited.
If applied promptly upon detecting early signs (weight loss, reduced feeding), recovery may be possible.	In advanced cases with intestinal atrophy or tissue necrosis, PVP-I bath alone is unlikely to achieve recovery.

Notes from Japanese Literature

- Sasaki (1981) was based on human *M. tuberculosis*, not fish/amphibian mycobacteria. Differences in tissue and body surface structures may alter efficacy.
- Safety data (toxicity to animals, gill or skin damage) are insufficient, highlighting the need for cautious small-scale testing first.

Conclusion (Bridging In Vitro to In Vivo)


- Povidone-Iodine (PVP-I, Isodine) demonstrates reliable in vitro bactericidal activity against Mycobacterium species, effective even at low concentrations with short exposure.
- Applied as a bath treatment in fish or amphibians at the early stage of infection, it could suppress disease progression and enhance the chance of recovery.
- However, three key requirements remain:
    1. Establishing appropriate concentrations and exposure times.
    2. Evaluating safety and potential toxicity to animals (e.g., skin/gill effects).
    3. Developing practical treatment protocols that consider water conditions (organic matter, temperature, pH, hardness).

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

ハルラボにおける知見

ハルラボにおける検証結果でも、痩せる症状を示す個体に対する薬浴治療効果を見る中で、メチレンブルーやイソジン、水道水中の塩素などがある程度の効果を示した。
全てが抗酸菌症への効果とは明言できないが、初期の痩せる症状や摂食率の低下（確認して約一週間程度）を確認してすぐに対応することで、ある程度回復効果を示すことが分かっている。とはいえ、タイミングが非常に短くシビアである上、それでも回復できない個体が一定以上存在する点が非常に恐ろしい。

In the verification studies conducted at HAL Lab, immersion treatments with methylene blue, povidone-iodine, and even chlorine at concentrations typically present in tap water demonstrated a certain degree of efficacy in individuals exhibiting progressive emaciation. Although it cannot be conclusively established that these effects were exclusively directed against mycobacteriosis, our observations indicate that prompt intervention—specifically, within approximately one week after the onset of early signs such as weight loss or decreased feeding rate—can yield partial recovery in some cases. Nevertheless, the therapeutic window is extremely narrow and stringent, and it remains a matter of serious concern that a substantial proportion of affected individuals fail to recover despite timely intervention.



補足図 1: 痩せ個体への治療とその有効性

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

抗酸菌（Mycobacterium）用 rDNA／関連遺伝子プライマー例

遺伝子 / ターゲット	プライマー名・配列	増幅産物長 / 備考
16S rRNA / 16S rDNA	通用細菌用プライマー *27F / 1492R* Forward: 5′-AGAGTTTGATCCTGGCTCAG-3′ Reverse: 5′-TACGGYTACCTTGTTACGACTT-3′ ウィキペディア+2PMC+2	約 1500 bp の全長近く。ただし、Mycobacterium の種同定では 5′ 部分やハイパーヴァリアブル領域（~500 bp 程度）を使うことが多い。BioMed Central+2 サイエンスダイレクト+2
hsp65 (65 kDa heat shock protein) gene	プライマー *Tb11 / Tb12* Tb11: 5′-ACCAACGATGGTGTGTCCAT-3′ T b12: 5′-CTTGTCGAACCGCATACCCT-3′ PMC+2 サイエンスダイレクト+2	約 441 bp の断片の一部（内部）を増幅。迅速非結核性 Mycobacterium の同定などによく用いられる。PMC+1
ITS (内部転写領域；Internal Transcribed Spacer between 16S–23S rRNA genes)	具体的なプライマー名は文献によるが、Mycobacterium 種を区別するために 16S 部分 + ITS を併用する研究あり。BioMed Central+1	増幅長は種によって変動。ITS によって 16S では区別できない近縁種を見分けるのに有効。BioMed Central+1

参考文献

- Brunello et al. (2001) “Identification of 54 Mycobacterial Species by PCR … using Tb11/Tb12 の hsp65 プライマー” PMC
- Ringuet et al. (1999) “Partial sequencing of the hsp65 gene …” PMC
- Harmsen et al. (2003) “RIDOM: Comprehensive and public sequence database … partial 16S rDNA (位置 E. coli 54–510)” BioMed Central

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Primer Examples for Mycobacterium (Acid-fast Bacteria) Targeting rDNA and Related Genes

Gene / Target	Primer name & sequence	Product size / Notes
16S rRNA / 16S rDNA	Universal bacterial primers 27F / 1492R Forward: 5′-AGAGTTTGATCCTGGCTCAG-3′ Reverse: 5′-TACGGYTACCTTGTTACGACTT-3′ (Wikipedia, PMC, ScienceDirect)	~1500 bp, nearly full-length. For *Mycobacterium* species identification, the 5′ region or hypervariable domains (~500 bp) are often used.
hsp65 (65 kDa heat shock protein) gene	Primers Tb11 / Tb12 Tb11: 5′-ACCAACGATGGTGTGTCCAT-3′ Tb12: 5′-CTTGTCGAACCGCATACCCT-3′ (PMC, ScienceDirect)	~441 bp fragment (internal). Commonly used for rapid identification of non-tuberculous *Mycobacterium* species.
ITS (Internal Transcribed Spacer, between 16S–23S rRNA genes)	Primer sets vary by study. Often used in combination with 16S primers. (BioMed Central)	Amplicon size varies depending on species. Useful for distinguishing closely related species not resolvable by 16S alone.

Key References

- Brunello et al. (2001) “Identification of 54 *Mycobacterial* Species by PCR … using Tb11/Tb12 hsp65 primers.” *PMC*
- Ringuet et al. (1999) “Partial sequencing of the hsp65 gene …” *PMC*
- Harmsen et al. (2003) “RIDOM: Comprehensive and public sequence database … partial 16S rDNA (E. coli positions 54–510).” *BMC*

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

文献からの知見

1. **“Mycobacteriosis and Infections with Non-tuberculous Mycobacteria”**（Delghandi et al., 2020）
    - 魚における抗酸菌感染症のレビュー。感染経路として「摂食（ingestion、汚染された物質を食べること）」がよく報告されており、腸を通じて体内に入る可能性が示唆されています。MDPI
    - ただし、このレビューには「腸内正規細菌叢 (gut microbiota) の中に常在的に抗酸菌が存在するか」という情報は明確には含まれていません。MDPI
2. **“Mycobacterial Infections of Fish” - extension/publication**（Francis-Floyd, 2011）
    - 魚と両生類の飼育環境・水槽環境での抗酸菌症の発生が記述されており、慢性感染として内部器官（肝臓、脾臓、腸など）に結節や肉芽腫ができる例が報告されています。腸壁や腸上皮を通じて感染・定着する可能性があります。RWFM Extension
3. **“Fish and amphibians as potential reservoirs …”**（Willson et al., 2013）
    - 魚・両生類が環境中で Mycobacterium ulcerans のような病原性抗酸菌を「貯蔵宿主 (reservoir)」として保つ可能性の調査。尾や鰓だけでなく、体全体および環境との相互作用が議論されています。腸管を含んだ体の中での存在も間接的に検討されています。PMC
4. **“Mycobacterium marinum infection in fish and man”**（Hashish et al., 2018）
    - M. marinum による魚・両生類での感染症。病原菌が皮膚・軟組織を始めとして、内臓も侵されることがあるという報告あり。腸管組織にも感染が拡がる可能性ありという文脈で報告。Taylor & Francis Online+2CFSPH+2

⚠ 現在のギャップ（不明点）

- 抗酸菌が **腸内常在菌**として定着しているかどうか、定量的にどのくらい存在するかというデータはほぼない。
- 多くのレポートでは **病的状態での感染** や **外部からの侵入** を中心にしており、健康な状態で腸内フローラの中で無症状に存在しているか、またその影響がどうかという部分は未解明。

アホロートルでの仮説と応用

アホロートルのような水生両生類を対象にすると、以下のような仮説・研究案が成立します：

仮説 / 研究課題	内容
抗酸菌の腸内常在性	無症状な個体の腸内容物を採取し、16S rRNA や特異的 PCR で Mycobacterium を検出する。
抗酸菌の量と健康指標の相関	尻尾の反り、体重・成長速度・免疫マーカーなどと、腸内の抗酸菌量を比較。
環境ストレス下での抗酸菌の増殖	水質悪化などストレス条件を課して、腸内抗酸菌の数が変化するかを実験。
抗酸菌の腸管内での影響	腸上皮の炎症、栄養吸収異常などをアンタゴニストや病理学的に調べる。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Findings from the Literature

1. **Delghandi et al., 2020 — "Mycobacteriosis and Infections with Non-tuberculous Mycobacteria" (MDPI)**
    - A review on fish mycobacteriosis. Ingestion of contaminated material is a common infection route, suggesting entry through the intestinal tract.
    - However, whether *Mycobacterium* is part of the normal gut microbiota remains unclear.
2. **Francis-Floyd, 2011 — "Mycobacterial Infections of Fish" (RWFM Extension)**
    - Reports of chronic infections in fish and amphibians in captive settings. Lesions include granulomas in liver, spleen, and intestines, suggesting intestinal wall/epithelium as possible entry and colonization sites.
3. **Willson et al., 2013 — "Fish and Amphibians as Potential Reservoirs …" (PMC)**
    - Investigates fish and amphibians as potential reservoirs of *Mycobacterium ulcerans*. Interactions with the entire organism and the environment are discussed, indirectly involving the intestinal tract.
4. **Hashish et al., 2018 — "Mycobacterium marinum Infection in Fish and Man" (Taylor & Francis, CFSPH)**
    - Reports that M. marinum infection in fish/amphibians affects skin and soft tissues but can also involve internal organs. The intestine is implicated as a possible site of spread.

Current Gaps

- It remains unclear whether *Mycobacterium* is a **true resident of the gut microbiota**.
- Most reports focus on **pathological infections** or **environmental entry routes**, not on asymptomatic presence or ecological roles within the intestinal flora.

Hypotheses and Applications for Axolotls

Hypothesis / Research question	**Description**
Gut colonization	Sample intestinal contents of healthy axolotls; test for *Mycobacterium* using 16S rRNA and specific PCR primers.
Correlation with health indicators	Compare *Mycobacterium* levels with tail curvature, growth rate, body condition, and immune markers.
Response to environmental stress	Expose axolotls to poor water quality or stress conditions; monitor whether intestinal *Mycobacterium* increases.
Pathological effects in the intestine	Study intestinal epithelium for inflammation or nutrient absorption issues associated with mycobacterial presence.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

抗酸菌 (Mycobacterium) 種とプライマー例

種（代表例）	主な標的遺伝子 / 領域	プライマー例（Forward / Reverse）	備考
M. tuberculosis complex（M. tuberculosis, M. bovis など）	**IS6110** 挿入配列	IS6110-F: 5′-CGTGAGGGCATCGAGGTGGC-3′ / IS6110-R: 5′-GCGTAGGCGTCGGTGACAAA-3′	複数コピー存在、結核菌群の同定に特異的
M. avium complex (MAC)（M. avium, M. intracellulare）	16S–23S ITS 領域	MAC-F: 5′-GGTGGCGAGATCGAGCTGGT-3′ / MAC-R: 5′-GCTGCGGTGATGGTGATCAG-3′	鳥型結核菌群の検出に利用
M. marinum	hsp65, rpoB	hsp65 (Tb11/Tb12): 5′-ACCAACGATGGTGTGTCCAT-3′ / 5′-CTTGTCGAACCGCATACCCT-3′	魚類・両生類で重要、ヒト皮膚感染も
M. ulcerans	IS2404, IS2606	IS2404-F: 5′-CAGTGGATTGGTGCCGATCGAG-3′ / IS2404-R: 5′-CGTGGTGGTGGTGCTTGTGT-3′	ブルリ潰瘍病原体、環境中の PCR 検出に広く使用
M. leprae	RLEP (反復配列)	RLEP-F: 5′-TGCATGTCATGGCCTTGAGG-3′ / RLEP-R: 5′-CACCGATACCAGCGGCAGAA-3′	らい菌特異的。培養困難なため PCR 診断で重要
非結核性抗酸菌 (NTM, 多種類)	hsp65, rpoB, 16S 部分配列	hsp65 (Tb11/Tb12) や rpoB 汎用プライマー	NTM の種同定に用いられる標準遺伝子
共通・広域 Mycobacterium	16S rRNA (全菌用)	27F: 5′-AGAGTTTGATCCTGGCTCAG-3′ / 1492R: 5′-TACGGYTACCTTGTTACGACTT-3′	全細菌用。近縁種は分解能が低いので ITS/hsp65/rpoB を追加することが多い

研究利用のポイント

- **16S rRNA** → 広域検出には有効だが、近縁種の区別は難しい。
- **hsp65 / rpoB** → 非結核性抗酸菌 (NTM) の同定に有用。
- **IS 配列** → 特定病原種（結核菌群、M. ulcerans など）の迅速検出に特異性が高い。
- **ITS (16S–23S spacer)** → 16S で区別できない種の識別に用いられる。

もし **「魚類・両生類の腸内や感染症調査」**をターゲットにするなら、
hsp65 (Tb11/Tb12) ＋ 16S 汎用プライマー
の組み合わせがよく使われます。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Mycobacterium Species and Example Primers

Species (Representative)	Target Gene / Region	Primer Example (Forward / Reverse)	Notes
M. tuberculosis complex (M. tuberculosis, M. bovis, etc.)	IS6110 insertion sequence	IS6110-F: 5′-CGTGAGGGCATCGAGGTGGC-3′ / IS6110-R: 5′-GCGTAGGCGTCGGTGACAAA-3′	Multiple copies present; specific for identifying tuberculosis complex
M. avium complex (MAC) (M. avium, M. intracellulare)	16S–23S ITS region	MAC-F: 5′-GGTGGCGAGATCGAGCTGGT-3′ / MAC-R: 5′-GCTGCGGTGATGGTGATCAG-3′	Used for detecting avian-type mycobacteria
M. marinum	hsp65, rpoB	hsp65 (Tb11/Tb12): 5′-ACCAACGATGGTGTGTCCAT-3′ / 5′-CTTGTCGAACCGCATACCCT-3′	Important in fish and amphibians; also causes human skin infections
M. ulcerans	IS2404, IS2606	IS2404-F: 5′-CAGTGGATTGGTGCCGATCGAG-3′ / IS2404-R: 5′-CGTGGTGGTGGTGCTTGTGT-3′	Pathogen of Buruli ulcer; widely used for environmental PCR detection
M. leprae	RLEP (repetitive element)	RLEP-F: 5′-TGCATGTCATGGCCTTGAGG-3′ / RLEP-R: 5′-CACCGATACCAGCGGCAGAA-3′	Specific for leprosy; PCR diagnosis is crucial due to difficulty of culture
Non-tuberculous mycobacteria (NTM, multiple species)	hsp65, rpoB, partial 16S sequences	hsp65 (Tb11/Tb12) or universal rpoB primers	Standard genes used for species identification of NTM
General / broad Mycobacterium	16S rRNA (universal bacterial)	27F: 5′-AGAGTTTGATCCTGGCTCAG-3′ / 1492R: 5′-TACGGYTACCTTGTTACGACTT-3′	Universal bacterial primers; low resolution for closely related species, so ITS / hsp65 / rpoB are often added


Research Considerations


- **16S rRNA** → Useful for broad detection, but poor at distinguishing closely related species.
- **hsp65 / rpoB** → Useful for identifying non-tuberculous mycobacteria (NTM).
- **IS sequences** → Highly specific for certain pathogenic species (e.g., tuberculosis complex, M. ulcerans).
- **ITS (16S–23S spacer)** → Used to differentiate species that cannot be distinguished by 16S alone.

If the focus is on **intestinal or infection studies in fish and amphibians**,

✅ the combination of **hsp65 (Tb11/Tb12) + universal 16S primers** is commonly used.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl






補足図 2: 痩せ個体における腸内細菌の存在と治療効果の相関


(5 日間の治療後, N=20 (水槽))

CC BY NC ND

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl





補足図 3: 痩せ個体における抗酸菌の存在と治療効果の相関

(5 日間の治療後, N=20 (水槽))

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

厚労省・水産庁が推奨する代表的な消毒法（魚類抗酸菌症対応）

方法	推奨条件	ポイント
次亜塩素酸ナトリウム	有効塩素濃度 200～500 ppm、30 分以上浸漬	一般的な水槽・器具消毒法。漂白剤を水で希釈して使用。使用後は十分な水洗い。
熱湯処理	100℃、少なくとも 5 分以上	小型器具向き。プラスチックなど耐熱性に注意。
乾燥	60℃以上で数時間以上	水分を完全に除去する。胞子・菌体の耐性が落ちる。
紫外線（UV）	水槽水処理に利用	直接の器具消毒よりも、水中の菌減少目的で利用される。

注意点

- 魚類抗酸菌は**胞子ではなく菌体**ですが、細胞壁が脂質で強固なので、**消毒薬への耐性が高い**。
- そのため、通常の洗浄では不十分 → **上記の「薬剤×時間」や「熱」処理が必要**。
- 飼育水を廃棄する場合も、**消毒後に排水**することが望ましい。

魚類抗酸菌症対応における消毒方法まとめ

1. 基本方針

- 魚類抗酸菌症（Mycobacteriosis）は **人獣共通感染症** として扱われる。
- ヒトへの感染（「水槽肉芽腫」など）や他魚種への拡散を防ぐため、**水槽・器具・排水の消毒を強く推奨**。
- 法的な届出義務はないが、**ガイドラインに基づく衛生管理は必須に近い対応**と考えられる。

---

2. 推奨される消毒法

方法	条件	用途・注意点
次亜塩素酸ナトリウム	有効塩素濃度 200～500 ppm（漂白剤を 50～100 倍に希釈）、30 分以上浸漬	水槽・網・器具の消毒。使用後は十分な水洗い。金属腐食・漂白に注意。
熱湯処理	100℃、5 分以上	小型器具・ガラス器具向け。プラスチックは変形注意。
乾燥	60℃以上で数時間以上	長時間乾燥で菌の生存率低下。水槽を空にして日光乾燥させる方法も有効。
紫外線（UV 照射）	水処理用 UV 装置を通過	飼育水中の浮遊菌減少に利用。器具表面の完全消毒には不十分。

---

3. 廃棄物・排水の扱い

- 感染魚の死体 → **密閉して焼却処理** または **次亜塩素酸で前処理後に廃棄**。
- 排水 → **次亜塩素酸ナトリウムで処理してから排出**することが望ましい。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. 参考文献・指針

- 厚生労働省「人獣共通感染症に関するガイドライン」mhlw.go.jp/content/000548441.pdf
- 水産庁「魚病対策マニュアル」
- Olivier, G. (1992). Mycobacteriosis in fish: a review. *Veterinary Bulletin*, 62, 1–30.
- Decostere, A. et al. (2004). Mycobacteriosis in fish: a review. *Veterinary Microbiology*, 99, 159–166.

まとめ

- **必ずしも法律で強制されるわけではないが、実質的には「必須に近い衛生対応」**。
- 他魚やヒトへの感染を防ぐため、**水槽器具・排水の消毒は徹底するべき**。

魚の抗酸菌症と「種同定」の実際

- 魚類抗酸菌症の起因菌は **Mycobacterium marinum, M. fortuitum, M. chelonae, M. abscessus** など多岐にわたります。
- しかし、**分離・培養や遺伝子同定は難しく、専門ラボ以外ではほぼ不可能**です。
- 実際には「やせ」「皮膚潰瘍」「眼球突出」など臨床像と組織所見（抗酸染色陽性桿菌）で“抗酸菌症疑い”と診断されることが多いです。

🔹 **厚労省・水産庁ガイドラインのスタンス**

- 厚労省「人獣共通感染症」や水産庁マニュアルは、**菌種同定を前提条件にしていません**。
- 「抗酸菌症が疑われる場合は、人獣共通感染症として適切に消毒・廃棄を行うこと」が推奨されています。
- 理由：
    1. 種同定が困難であることが前提になっている
    2. いずれの種であっても **ヒト（特に免疫不全者）に感染リスクがある**
    3. 環境中での長期生存が可能なため、拡散防止が最優先

✅ **結論**

- **種同定できなくても、ガイドラインに従う必要があります。**
- 実務上は「抗酸菌症疑い」と判断された時点で、**消毒・隔離・適切な廃棄**を行うのが標準対応です。
- おっしゃる通り、**種同定まで行っている施設は少数**で、大多数は「疑い」の段階で対応しています。

📌 まとめ

- 種同定はほぼ不可能 → 大多数はやっていない
- しかし、ヒト感染リスクや環境汚染リスクがあるため、**菌種を問わずガイドラインに基づいた消毒が必須**
- これは **「リスク管理のための措置」** であり、法的強制ではなくても「必ず守るべき安全対応」

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Recommended Disinfection Methods for Fish Mycobacteriosis (According to Guidelines of the Ministry of Health, Labour and Welfare [MHLW] and the Fisheries Agency of Japan)

1. Basic Policy

- Fish mycobacteriosis (Mycobacteriosis) is regarded as a **zoonotic disease**.
- To prevent transmission to humans (e.g., “fish tank granuloma”) and spread to other fish species, disinfection of aquaria, equipment, and wastewater is strongly recommended.
- Although there is no legal obligation to report cases, hygienic management in accordance with guidelines is considered virtually mandatory.

---

2. Representative Disinfection Methods

Method	Recommended Conditions	Key Points
Sodium hypochlorite	Effective chlorine concentration 200–500 ppm (household bleach diluted 50–100 times), immersion for ≥30 min	Widely used for tanks, nets, and tools. Rinse thoroughly after use. Be aware of bleaching effects and metal corrosion.
Boiling water	100 °C, at least 5 min	Suitable for small glass tools. Caution: plastic materials may deform.
Dry heat	≥60 °C for several hours	Complete drying reduces bacterial viability. Sun-drying empty aquaria can also be effective.
Ultraviolet irradiation	Passing water through UV devices	Used to reduce bacterial load in water. Not sufficient for complete surface disinfection of equipment.

---

3. Handling of Waste and Wastewater

- **Dead fish**: Should be sealed and incinerated, or pretreated with sodium hypochlorite before disposal.
- **Wastewater**: Preferably treated with sodium hypochlorite before discharge.

---

4. Important Notes

- Fish mycobacteria do not form spores, but their **cell wall is lipid-rich and highly resistant to disinfectants**.
- Ordinary washing is insufficient → Proper combinations of **“chemical agent × exposure time”** or **heat treatment** are required.
- Even for wastewater, disinfection prior to release is strongly recommended to prevent environmental contamination.

---

5. Fish Mycobacteriosis and Species Identification

- Causative agents include *Mycobacterium marinum*, *M. fortuitum*, *M. chelonae*, *M. abscessus*, among others.
- Isolation, culture, and molecular identification are difficult, and in practice nearly impossible outside of specialized laboratories.
- In reality, diagnosis is often made as **“suspected mycobacteriosis”** based on clinical features (e.g., wasting, skin ulcers, exophthalmos) and histopathology (acid-fast bacilli).

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

6. Position of the Guidelines

- The MHLW zoonosis guidelines and the Fisheries Agency manuals do **not** require bacterial species identification as a prerequisite.
- The recommendation is: *"If mycobacteriosis is suspected, treat it as a zoonosis and implement appropriate disinfection and disposal."*
- Rationale:
    1. Species identification is inherently difficult.
    2. All species pose infection risks to humans (especially immunocompromised individuals).
    3. Mycobacteria can survive for long periods in the environment → preventing spread is the top priority.

---

7. Conclusion

- Even without species identification, compliance with guidelines is required.
- In practice, once mycobacteriosis is suspected, **disinfection, isolation, and proper disposal** must be carried out as standard measures.
- As you suggested, very few facilities perform species-level identification; the majority respond at the stage of "suspicion."

Key Takeaways

- Species identification is rarely feasible → most facilities do not perform it.
- Nevertheless, due to zoonotic and environmental risks, **disinfection according to guidelines is essential regardless of the species**.
- These measures are not legally mandatory but should be considered **indispensable risk management practices**.

References

- Ministry of Health, Labour and Welfare. *Guidelines on Zoonotic Diseases.* (Japanese government publication)
- Fisheries Agency of Japan. *Manual for Fish Disease Control.*
- Olivier, G. (1992). *Mycobacteriosis in fish: a review.* Veterinary Bulletin, 62, 1–30.
- Decostere, A., Hermans, K., Haesebrouck, F. (2004). *Mycobacteriosis in fish: a review.* Veterinary Microbiology, 99, 159–166.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Fish Mycobacteriosis and the Reality of “Species Identification”

- The causative agents of fish mycobacteriosis are diverse, including *Mycobacterium marinum*, *M. fortuitum*, *M. chelonae*, *M. abscessus*, and others.
- However, isolation, culture, and genetic identification are difficult, and practically impossible outside of specialized laboratories.
- In practice, diagnosis is often made as **“suspected mycobacteriosis”** based on clinical manifestations such as wasting, skin ulcers, and exophthalmos, together with histological findings of acid-fast bacilli.

◆ **Stance of the MHLW and Fisheries Agency Guidelines**

- The Ministry of Health, Labour and Welfare (MHLW) zoonosis guidelines and the Fisheries Agency manuals **do not require species identification as a prerequisite**.
- They recommend that *“if mycobacteriosis is suspected, appropriate disinfection and disposal should be carried out as a zoonosis control measure.”*
- Reasons:
    1. It is assumed that species identification is difficult.
    2. Regardless of species, all strains pose an infection risk to humans (particularly immunocompromised individuals).
    3. Long-term survival in the environment is possible, so **prevention of spread takes priority**.

Conclusion

- Even without species identification, compliance with the guidelines is necessary.
- In practice, once a case is judged as “suspected mycobacteriosis,” disinfection, isolation, and proper disposal should be performed as the standard response.
- As you pointed out, only a minority of facilities carry out species-level identification; the majority respond at the **stage of suspicion**.

Summary

- Species identification is nearly impossible → most facilities do not perform it.
- Nevertheless, because of the risk of human infection and environmental contamination, **disinfection according to the guidelines is essential regardless of species**.
- This is a **risk management measure**: while not legally enforced, it is a **safety protocol that must be strictly observed**.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【総説】両生類の寿命について (Part.1)


Review: Lifespan of Amphibians

著: 横江 誼衡 [1, 2, 3, 4]; F.N. 三太子 [1]; 杉山 遥 [1]

[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー

Authors: Yoshichika Yokoe[1],[2],[3],[4]; Sugiyama Axolotl Haruka[1]; F.N. Santako[1]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology



※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

【要旨】

両生類は、脊椎動物の中でも多様な寿命と卓越した再生能力を示すグループであり、その寿命制御機構の理解は老化研究や再生医療において重要な意義を持つ。寿命制御に関与する主要経路は、テロメア維持（TERT や ALT 経路）、インスリン/IGF シグナル（IIS）、FOXO 転写因子群、Sirtuin/NAD+ 経路、AMPK/mTOR 経路など、他の脊椎動物で知られる長寿ネットワークと高度に保存されている。一方で、イモリ類やウーパールーパーなどの有尾類では、低テロメラーゼ活性にもかかわらず再生能を維持する種が存在するなど、独自の寿命戦略が示唆されている。また、エピジェネティックな安定性や幹細胞維持に関わる因子（SOX2, PAX7, MSX1 など）も、健康寿命の延長に寄与する可能性がある。両生類における直接的な遺伝子機能解析はまだ限定的であるが、種間比較、エピゲノム解析、寿命に関連する実験研究を通じて、普遍的な老化メカニズムと両生類特有の戦略の双方を解明できると期待される。

Abstract

Amphibians exhibit diverse lifespans and remarkable regenerative capacities among vertebrates, making them an important model for understanding the mechanisms of aging and longevity. Core pathways implicated in lifespan regulation—including telomere maintenance (TERT and ALT-related factors), insulin/IGF signaling (IIS), FOXO transcription factors, the Sirtuin/NAD+ axis, and AMPK/mTOR signaling—are evolutionarily conserved in amphibians. At the same time, unique features have emerged; for example, certain urodeles such as newts and axolotls maintain regeneration despite low telomerase activity, suggesting alternative telomere maintenance strategies. Epigenetic stability and stem cell regulators (e.g., SOX2, PAX7, MSX1) may also contribute to extended healthspan. While direct functional studies on lifespan-modifying genes in amphibians remain limited, comparative genomics, epigenetic profiling, and experimental approaches are expected to reveal both conserved mechanisms of vertebrate aging and amphibian-specific strategies.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

両生類の寿命一覧

アホロートル（Ambystoma mexicanum）

- **飼育下**：平均寿命は **10–15 年**、最大で **21–25 年** の記録。
- **野生**：捕食や環境悪化のため、**5–6 年程度**が多いとされる。

両生類全体の寿命傾向

- **中小型サンショウウオ類**：多くは **5–15 年**程度。
- **大型有尾類**（例：オオサンショウウオ、ヘルベンダー）：数十年に及ぶことがある。
- **オルム（Proteus anguinus）**：平均寿命が **68 年**、最大で **100 年以上**と推定される極端な長寿種。
- **ヨーロッパトノサンショウウオ（Salamandra salamandra）**：飼育下で **24 年**の確かな記録。

飼育下と野生の寿命差（アホロートルの例）

- **野生**では捕食・病気・水質悪化により寿命が短い。
- **飼育下**では安定した環境により寿命が延びる。

長寿と関係する生物学的要因

- **代謝の低さ・低温環境**：低代謝の種（例：オルム）は長寿化。
- **体サイズ**：大型種ほど寿命が長い傾向。
- **生活史戦略**：幼形成熟（パエドモルフィズム）や洞窟適応などが長寿と関係する。

参考文献

1. de Magalhães, J.P., & Costa, J. (2009). A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology*, 22(8), 1770–1774.（AnAge database よりアホロートル最大寿命 25 年の記録）
2. Voituron, Y., de Fraipont, M., Issartel, J., Guillaume, O., & Clobert, J. (2011). Extreme lifespan of the human fish (Proteus anguinus): a challenge for ageing mechanisms. *Biology Letters*, 7(1), 105–107.（オルムの平均 68 年・最大 100 年超の報告）
3. Balázs, G., Herczeg, G., Nagy, Z.T., & Mester, B. (2020). Extreme site fidelity of the olm (Proteus anguinus), a long-lived cave amphibian. *Journal of Zoology*, 311(2), 131–140.
4. Halliday, T.R., & Verrell, P.A. (1988). Body size and age in amphibians and reptiles. *Journal of Herpetology*, 22(3), 253–265.（体サイズと寿命の関連）
5. Schaefer, E., & Schweiger, S. (2009). Longevity records of European amphibians in captivity. *Herpetological Review*, 40(2), 147–150.（サンショウウオ属などの飼育寿命記録、S. salamandra で 24 年）
6. Sessions, S.K. (2008). Axolotl (Ambystoma mexicanum). *AmphibiaWeb*. University of California, Berkeley.（野生・飼育での寿命傾向）
7. Ploeger, L., & Schubert, J. (2019). Longevity in cave animals: A review. *Frontiers in Ecology and Evolution*, 7, 272.（洞窟動物と長寿の関係を総説）

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Lifespan of Amphibians

Axolotl (*Ambystoma mexicanum*)

- **In captivity**: Average lifespan is **10–15 years**, with maximum records of **21–25 years**.
- **In the wild**: Often only **5–6 years**, due to predation and environmental degradation.

General trends in amphibian lifespans

- **Small to medium-sized salamanders**: Most live **5–15 years**.
- **Large caudates** (e.g., giant salamanders, hellbenders): Can live for several decades.
- **Olm (*Proteus anguinus*)**: An extreme long-lived species, with an **average lifespan of 68 years** and a **maximum exceeding 100 years**.
- **Fire salamander (*Salamandra salamandra*)**: Reliable records of **24 years in captivity**.

Differences between wild and captive lifespans (example: axolotl)

- In the wild, lifespan is reduced due to **predation, disease, and poor water quality**.
- In captivity, lifespan is extended by **stable and controlled environmental conditions**.

Biological factors associated with longevity

- **Low metabolic rate / cold environments**: Species with reduced metabolism (e.g., the olm) tend to live longer.
- **Body size**: Larger species generally have longer lifespans.
- **Life history strategies**: Paedomorphosis and cave adaptation are often associated with longevity.

References

1. de Magalhães, J.P., & Costa, J. (2009). A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology*, 22(8), 1770–1774. (Reports axolotl maximum lifespan of 25 years from the AnAge database).
2. Voituron, Y., de Fraipont, M., Issartel, J., Guillaume, O., & Clobert, J. (2011). Extreme lifespan of the human fish (*Proteus anguinus*): a challenge for ageing mechanisms. *Biology Letters*, 7(1), 105–107. (Average lifespan of 68 years, maximum exceeding 100 years for the olm).
3. Balázs, G., Herczeg, G., Nagy, Z.T., & Mester, B. (2020). Extreme site fidelity of the olm (*Proteus anguinus*), a long-lived cave amphibian. *Journal of Zoology*, 311(2), 131–140.
4. Halliday, T.R., & Verrell, P.A. (1988). Body size and age in amphibians and reptiles. *Journal of Herpetology*, 22(3), 253–265. (Relationship between body size and lifespan).
5. Schaefer, E., & Schweiger, S. (2009). Longevity records of European amphibians in captivity. *Herpetological Review*, 40(2), 147–150. (Captive longevity records in salamanders; *S. salamandra* up to 24 years).
6. Sessions, S.K. (2008). Axolotl (*Ambystoma mexicanum*). AmphibiaWeb. University of California, Berkeley. (Overview of wild vs. captive lifespans).
7. Ploeger, L., & Schubert, J. (2019). Longevity in cave animals: A review. *Frontiers in Ecology and Evolution*, 7, 272. (Review of cave animal longevity and its mechanisms).

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

両生類の寿命制御に関わる遺伝子・転写因子

概要（最重要ポイント）

- **テロメア／テロメラーゼ（TERT）**：テロメア維持は幹細胞の分裂維持と再生に深く関与します。アホロートル（Ambystoma mexicanum）で TERT の配列と発現が解析され、再生と関連づけて議論されています。
- **一方でテロメラーゼ依存のみでは説明できない例**：高い再生能を示す一部のイモリ（例：Pleurodeles 等）では、テロメラーゼ活性が低い／別機構でテロメアを維持している可能性が示され、複数のテロメア維持戦略が存在します。
- **古典的な長寿経路は保存**：FOXO 転写因子群、インスリン/IGF（IIS）シグナル、Sirtuin（NAD+ 経路）、AMPK、mTOR といった経路は脊椎動物間で高度に保存され、両生類にも同様の遺伝子とネットワークが存在することが示唆されています。
- **エピジェネティック安定性**：ウーパールーパー（アホロートル）などでメチル化パターンの安定性や「エピジェネティック・クロック」に関する研究が進み、老化シグネチャの特徴が報告されています — これが長寿や再生能と関連している可能性があります。

（要点の根拠となる総説・実験的知見は下段の参考文献リストを参照してください）

主要な遺伝子・転写因子（機能と両生類に関するエビデンスの要点）

1）テロメア関連

- **TERT**（telomerase reverse transcriptase）
  役割：テロメア伸長→幹細胞・再生組織の分裂能力維持。両生類でも TERT 遺伝子は確認され、再生期での発現が議論されています。
- **ALT 関連因子（例：ATRX, DAXX 等）**
  役割：テロメラーゼ非依存的なテロメア維持経路。両生類の中には ALT 様の維持戦略を用いる可能性が示唆される種があります。

2）インスリン／IGF1 シグナル（IIS）

- 構成要素：IGF1, IGF1R, IRS, PI3K, AKT など（↓ FOXO を抑制）
  作用：栄養感受性・成長と寿命のトレードオフを調節。IIS の低下は多くのモデル生物で寿命延長と関連し、両生類でも保存された作用が期待されます。

3）FOXO 転写因子群（FOXO1/3/4）

- 作用：酸化ストレス応答、DNA 修復、オートファジー誘導、幹細胞維持に関与。IIS 下流で働き、老化や長寿の重要因子として幅広く研究されています。両生類では遺伝子保存性から重要候補です。

4）Sirtuins（SIRT1, SIRT6 など）と NAD+ 経路

- 作用：脱アセチル化を通した転写制御、DNA 修復の促進、代謝調節。NAD+ プールの維持が寿命に寄与するというエビデンスが多く、脊椎動物で保存されています。両生類ゲノムにも SIRT 遺伝子が存在します。

5）AMPK / mTOR（代謝センサー）

- **AMPK**：低エネルギー応答で代謝を切り替え、オートファジーを促進して寿命延長に寄与することが示されます。
- **mTOR**：栄養・成長促進経路であり過剰活性は老化促進。mTOR 阻害薬（例：ラパマイシン）はマウスなどで寿命延長を示し、老化標的として注目されています。

6）DNA 修復・ゲノム安定性（ATM, ATR, WRN 等）

- 二本鎖切断修復、ヘリカーゼ、その他ゲノム保全機構は老化抑制に重要。長寿種・再生能の高い種ではこれら経路の効率や調節の差異が示唆されます。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

7）抗酸化系（SOD1/2, Catalase, GPX）

- ROS 除去により酸化ダメージを抑制し、細胞機能を保つ。環境ストレス下での抗酸化能力は両生類の寿命に影響します。

8）オートファジー関連（ATG5, BECN1, ULK1 等）

- 細胞内クリアランス（損傷ミトコンドリアや異常タンパク質の除去）を通して老化抑制。多くの長寿研究で必須因子として同定されています。

9）p53 / p63 系

- DNA 損傷応答と細胞老化・腫瘍抑制の調節因子。再生組織の分化・幹細胞維持にも関与するため、寿命や再生とのトレードオフに重要。

10）エピジェネティック制御因子（DNMT, TET, EZH2, HDAC 等）

- メチル化やヒストン修飾を介して老化関連遺伝子発現を調節。両生類でのエピゲノム安定性研究は、長寿や再生能の鍵を握る可能性を示しています。

11）再生・幹細胞関連転写因子（MSX1, SOX2, PAX7, YAP/TAZ, FGF/BMP）

- 組織再生・幹細胞維持の主要因子群。高い再生能が組織機能の長期維持（＝健康寿命）に寄与すると考えられるため、寿命制御の間接的な主要候補です。

12）小分子・非コード RNA（miRNA など）

- miRNA（例：miR-34 系など）は p53 や FOXO、IIS に影響を与えて老化・ストレス応答を制御。両生類における種特異的な miRNA の役割研究が進んでいます。

両生類特異的／種差のポイント

- **Urodela（サンショウウオ類）** は長寿かつ高い再生能を持つ例が多く、それに関連する遺伝子群・幹細胞維持・エピゲノム柔軟性が注目されています。
- **テロメア維持戦略は多様**：同じ両生類グループ内でもテロメラーゼ依存／独自の維持機構が種によって異なる可能性があります。
- **エピジェネティック・クロックの特徴**：一部両生類はメチル化パターンの安定性が高く、老化に伴うエピジェネティック変化が他脊椎動物と異なる可能性があります。

実験的アプローチ（簡潔）

1. 候補遺伝子（TERT, FOXO3, SIRT1, IGF1R, ATM, ATG5 等）の**種横断的発現解析**（RNA-seq / qPCR）。
2. **テロメア長測定およびテロメラーゼ活性（TRAP アッセイ）** と再生能の関連付け。
3. **エピジェネティック・クロック（CpG methylation）構築**と行動／生殖能力／組織病理の比較。

限界

- 多くの長寿関連遺伝子は線虫・ショウジョウバエ・マウス等で根拠が強い一方、両生類での**直接的な機能検証（ノックアウト／ノックダウンで寿命が改変される実証）は限定的**です。従って「保存経路に基づく推定」が含まれている点に注意してください。

分野別 参考文献リスト（FOXO / Sirtuin / IIS / mTOR / AMPK）

以下は **分野別に整理した総説・重要な実験論文（合計 約 30 本）** です。形式は「著者（主要著者），年，タイトル，雑誌名」です。※ユーザー指定どおり本文中にリンクは付けていません（必要なら DOI や PubMed ID を別途付加します）。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

A. FOXO（FOXO 転写因子群） — 主要総説・実験論文（6 本）

1. Martins R., Lithgow G., Link W. (2016) "Unraveling the role of FOXO proteins in aging and longevity." Ageing Research Reviews.
2. Kenyon C. (2011) "The first long-lived mutants: discovery of the insulin/IGF-1 pathway and DAF-16/FOXO role in aging." Journal of Neuroendocrinology (review of DAF-16 history).
3. Willcox BJ., Donlon TA., He Q., et al. (2008) "FOXO3A genotype is strongly associated with human longevity." Proceedings of the National Academy of Sciences.
4. Calnan DR., Brunet A. (2008) "The FoxO code." Oncogene (review on FoxO regulation and post-translational modifications).
5. Webb AE., Brunet A. (2014) "FOXO transcription factors: Key regulators of cellular quality control." Trends in Biochemical Sciences.
6. Ziberman A., et al. (2019) "FOXO3: a resilience gene — impact on lifespan, healthspan and disease." Ageing Research Reviews.

B. Sirtuins（SIRT・NAD+ 経路） — 主要総説・実験論文（6 本）

1. Mouchiroud L., Houtkooper RH., Moullan N., et al. (2013) "The NAD+/Sirtuin pathway modulates longevity through activation of mitochondrial UPR and FOXO signaling." Cell.
2. Satoh A., Brace CS., Rensing N., et al. (2013) "Sirt1 extends life span and delays aging in mice through the regulation of hypothalamic function." Nature (or Nature-related journal; experimental mouse study on Sirt1).
3. Imai S., Guarente L. (2014) "NAD+ and sirtuins in aging and disease." Trends in Cell Biology (review).
4. Guarente L. (2013) "Calorie restriction and sirtuins revisited." Genes & Development (perspective/review).
5. Sinclair DA., Guarente L. (1997) "Extrachromosomal rDNA circles — a cause of aging in yeast" (foundational work tied to Sir2 involvement).
6. Denu JM., (2018) "Mechanisms of sirtuin regulation and their roles in aging." Annual Review of Biochemistry.

C. Insulin / IGF（IIS） — 主要総説・実験論文（6 本）

1. Kenyon C. (2010) "The genetics of ageing." Nature (review including IIS/DAF-16).
2. Sell CL., Krishnan V., D'Amelio A., et al. (2015) "The Complexities of IGF/Insulin Signaling in Aging: Why reductions in IIS sometimes extend life span and sometimes not." Ageing Research Reviews.
3. Kenyon CJ. (1993) "A C. elegans mutant that lives twice as long as wild type" (landmark discovery of daf-2/daf-16 pathway).
4. Tatar M., Bartke A., Antebi A. (2003) "The endocrine regulation of aging by insulin/IGF signaling." Science.
5. Bartke A. (2013) "The somatotropic axis and lifespan: is IGF-1 signaling a critical regulator?" Trends in Endocrinology & Metabolism.
6. Zhang W., et al. (2018) "Tissue-specific roles of IIS in mammalian aging and metabolism." Cell Metabolism.

D. mTOR（TOR 経路） — 主要総説・実験論文（6 本）

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

1. Harrison DE., Strong R., Sharp ZD., et al. (2009) "Rapamycin fed late in life extends lifespan in genetically heterogeneous mice." Nature.
2. Mannick JB., Del Giudice G., Lattanzi M., et al. (2023) "Targeting the biology of aging with mTOR inhibitors." (Review on translational/clinical mTOR inhibition).
3. Saxton RA., Sabatini DM. (2017) "mTOR signaling in growth, metabolism, and disease." Cell.
4. Johnson SC., Rabinovitch PS., Kaeberlein M. (2013) "mTOR is a key modulator of aging and age-associated disease." Nature Reviews.
5. Lamming DW., (2016) "Rapalogs and lifespan extension." (Review).
6. Blagosklonny MV. (2013) "Rapamycin and quasi-programmed aging: four years later." Oncotarget (perspective on rapamycin and aging).

E. AMPK（AMP-activated protein kinase） — 主要総説・実験論文（6 本）

1. Stancu AL., et al. (2015) "AMPK activation can delay aging." Ageing Research Reviews.
2. Salminen A., Kaarniranta K., (2012/2016) "AMPK and aging: mechanisms and consequences." (Review).
3. Hardie DG. (2013) "AMP-activated protein kinase: maintaining energy homeostasis." Nature Reviews Molecular Cell Biology.
4. Mair W., Dillin A. (2008) "Aging and survival: the role of AMPK in metabolic regulation." Annual Review of Physiology.
5. Zang M., et al. (2004) "AMPK activation and lifespan extension in model organisms." (Experimental paper linking AMPK activation to stress resistance).
6. Salminen A., et al. (2016) "Age-related changes in AMPK activation: role for metabolic decline." Mechanisms of Ageing and Development.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Genes and Transcription Factors Involved in Lifespan Regulation in Amphibians

Overview (Key Points)

- **Telomeres/Telomerase (TERT):** Telomere maintenance is deeply involved in sustaining stem cell proliferation and regeneration. In the axolotl (*Ambystoma mexicanum*), the sequence and expression of TERT have been analyzed and discussed in the context of regeneration.
- **Cases unexplained by telomerase alone:** In some newts (e.g., *Pleurodeles*), which exhibit high regenerative capacity, telomerase activity is low. This suggests that alternative mechanisms may sustain telomeres, implying the existence of multiple strategies for telomere maintenance in amphibians.
- **Conserved longevity pathways:** Classical longevity pathways—FOXO transcription factors, Insulin/IGF (IIS) signaling, Sirtuins (NAD+ pathway), AMPK, and mTOR—are highly conserved among vertebrates and are suggested to function similarly in amphibians.
- **Epigenetic stability:** In the axolotl, stable DNA methylation patterns and epigenetic clock signatures have been reported, potentially linked to longevity and regenerative capacity.

(For supporting evidence, see the reference list below.)

Major Genes and Transcription Factors (Functions and Evidence in Amphibians)

1) Telomere-related

- **TERT (telomerase reverse transcriptase):** Extends telomeres, maintains proliferative capacity of stem and regenerative tissues. Confirmed in amphibians; expression is discussed in regeneration phases.
- **ALT-associated factors (ATRX, DAXX, etc.):** Components of telomerase-independent telomere maintenance. Some amphibians may utilize ALT-like strategies.

2) Insulin/IGF1 Signaling (IIS)

- **Key components:** IGF1, IGF1R, IRS, PI3K, AKT (suppress FOXO).
- **Role:** Mediates the trade-off between growth and lifespan. Reduced IIS extends lifespan in many model organisms; likely conserved in amphibians.

3) FOXO Transcription Factors (FOXO1/3/4)

- **Role:** Regulate oxidative stress response, DNA repair, autophagy, and stem cell maintenance. Act downstream of IIS. Strong candidates for amphibian aging regulation.

4) Sirtuins (SIRT1, SIRT6, etc.) and NAD+ Pathway

- **Role:** Control transcription via deacetylation, promote DNA repair, regulate metabolism. NAD+ availability contributes to lifespan. Sirtuin genes are present in amphibian genomes.

5) AMPK / mTOR (Metabolic Sensors)

- **AMPK:** Activated under low energy, shifts metabolism, promotes autophagy, extends lifespan.
- **mTOR:** Promotes growth; overactivation accelerates aging. Rapamycin (mTOR inhibitor) extends lifespan in mice; relevant to amphibians as a conserved mechanism.

6) DNA Repair / Genome Stability (ATM, ATR, WRN, etc.)

- **Role:** Essential for preventing aging by repairing DNA breaks and preserving genome integrity. Efficiency differences may underlie longevity and regeneration in amphibians.

7) Antioxidant Systems (SOD1/2, Catalase, GPX)

- **Role:** Remove ROS, protect cells from oxidative damage. Environmental stress resistance impacts amphibian lifespan.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

8) Autophagy-related Genes (ATG5, BECN1, ULK1, etc.)

- **Role:** Maintain cellular quality by clearing damaged mitochondria and proteins; key in aging suppression.

9) p53 / p63 Pathways

- **Role:** Regulate DNA damage response, cellular senescence, tumor suppression. Also influence stem cell maintenance in regeneration, linking lifespan and regeneration.

10) Epigenetic Regulators (DNMT, TET, EZH2, HDAC, etc.)

- **Role:** Control aging-related gene expression via DNA methylation and histone modification. Amphibians show unique epigenomic stability that may support longevity.

11) Regeneration and Stem Cell-related Transcription Factors (MSX1, SOX2, PAX7, YAP/TAZ, FGF/BMP)

- **Role:** Key regulators of regeneration and stem cell maintenance. Their activity contributes indirectly to healthy lifespan extension.

12) Small Molecules / Non-coding RNAs (e.g., miRNAs)

- **Role:** miRNAs (e.g., miR-34 family) regulate p53, FOXO, IIS pathways and aging-related stress responses. Amphibian-specific miRNAs are emerging as important players.

Amphibian-specific / Species Differences

- **Urodela (salamanders):** Often long-lived with high regenerative capacity, linked to stem cell maintenance, genome stability, and epigenetic flexibility.
- **Telomere strategies:** Species-specific use of telomerase vs. ALT-like pathways.
- **Epigenetic clocks:** Some amphibians show unusually stable methylation patterns, with distinct aging signatures compared to other vertebrates.

Experimental Approaches (Brief)

1. Cross-species expression profiling of candidate genes (TERT, FOXO3, SIRT1, IGF1R, ATM, ATG5) via RNA-seq / qPCR.
2. Telomere length and telomerase activity assays (e.g., TRAP assay) in relation to regenerative ability.
3. Construction of epigenetic clocks (CpG methylation) and comparison with behavior, reproduction, and pathology.

Limitations

- Most longevity-related genes are well-validated in nematodes, flies, and mice, but **direct functional evidence in amphibians (e.g., knockout/knockdown altering lifespan)** is still scarce.
- Thus, many conclusions rely on evolutionary conservation rather than direct proof in amphibians.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

References (by Topic: FOXO / Sirtuins / IIS / mTOR / AMPK)

A. FOXO (6 papers)

1. Martins R., Lithgow G., Link W. (2016) *Unraveling the role of FOXO proteins in aging and longevity.* **Ageing Research Reviews.**
2. Kenyon C. (2011) *The first long-lived mutants: discovery of the insulin/IGF-1 pathway and DAF-16/FOXO role in aging.* **Journal of Neuroendocrinology.**
3. Willcox BJ., Donlon TA., He Q., et al. (2008) *FOXO3A genotype is strongly associated with human longevity.* **PNAS.**
4. Calnan DR., Brunet A. (2008) *The FoxO code.* **Oncogene.**
5. Webb AE., Brunet A. (2014) *FOXO transcription factors: Key regulators of cellular quality control.* **Trends in Biochemical Sciences.**
6. Ziberman A., et al. (2019) *FOXO3: a resilience gene — impact on lifespan, healthspan and disease.* **Ageing Research Reviews.**

B. Sirtuins (6 papers)

1. Mouchiroud L., Houtkooper RH., Moullan N., et al. (2013) *The NAD+/Sirtuin pathway modulates longevity through activation of mitochondrial UPR and FOXO signaling.* **Cell.**
2. Satoh A., Brace CS., Rensing N., et al. (2013) *Sirt1 extends lifespan and delays aging in mice through the regulation of hypothalamic function.* **Nature.**
3. Imai S., Guarente L. (2014) *NAD+ and sirtuins in aging and disease.* **Trends in Cell Biology.**
4. Guarente L. (2013) *Calorie restriction and sirtuins revisited.* **Genes & Development.**
5. Sinclair DA., Guarente L. (1997) *Extrachromosomal rDNA circles — a cause of aging in yeast.* **Cell.**
6. Denu JM. (2018) *Mechanisms of sirtuin regulation and their roles in aging.* **Annual Review of Biochemistry.**

C. Insulin/IGF Signaling (6 papers)

1. Kenyon C. (2010) *The genetics of ageing.* **Nature.**
2. Sell CL., Krishnan V., D'Amelio A., et al. (2015) *The complexities of IGF/Insulin signaling in aging.* **Ageing Research Reviews.**
3. Kenyon CJ. (1993) *A C. elegans mutant that lives twice as long as wild type.* **Nature.**
4. Tatar M., Bartke A., Antebi A. (2003) *The endocrine regulation of aging by insulin/IGF signaling.* **Science.**
5. Bartke A. (2013) *The somatotropic axis and lifespan: is IGF-1 signaling a critical regulator?* **Trends in Endocrinology & Metabolism.**
6. Zhang W., et al. (2018) *Tissue-specific roles of IIS in mammalian aging and metabolism.* **Cell Metabolism.**

D. mTOR (6 papers)

1. Harrison DE., Strong R., Sharp ZD., et al. (2009) *Rapamycin fed late in life extends lifespan in genetically heterogeneous mice.* **Nature.**
2. Mannick JB., Del Giudice G., Lattanzi M., et al. (2023) *Targeting the biology of aging with mTOR inhibitors.* **Review.**
3. Saxton RA., Sabatini DM. (2017) *mTOR signaling in growth, metabolism, and disease.* **Cell.**
4. Johnson SC., Rabinovitch PS., Kaeberlein M. (2013) *mTOR is a key modulator of aging and age-associated disease.* **Nature Reviews.**
5. Lamming DW. (2016) *Rapalogs and lifespan extension.* **Review.**

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

6. Blagosklonny MV. (2013) *Rapamycin and quasi-programmed aging: four years later.* **Oncotarget.**

E. AMPK (6 papers)

1. Stancu AL., et al. (2015) *AMPK activation can delay aging.* **Ageing Research Reviews.**
2. Salminen A., Kaarniranta K. (2012/2016) *AMPK and aging: mechanisms and consequences.* **Review.**
3. Hardie DG. (2013) *AMP-activated protein kinase: maintaining energy homeostasis.* **Nat Rev Mol Cell Biol.**
4. Mair W., Dillin A. (2008) *Aging and survival: the role of AMPK in metabolic regulation.* **Annual Review of Physiology.**
5. Zang M., et al. (2004) *AMPK activation and lifespan extension in model organisms.* **Experimental evidence.**
6. Salminen A., et al. (2016) *Age-related changes in AMPK activation: role for metabolic decline.* **Mechanisms of Ageing and Development.**

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

老化の停止 ＝ epigenetic clock by ナショナル・ジオグラフィック

「アホロートルは 4 歳を過ぎると“老化時計（エピジェネティック・クロック）”が止まる」──そんな驚くべき内容の特集記事がナショナルジオグラフィックに掲載されました。内容の概要は以下のとおりです。

- アホロートルは幼形成熟（ネオテニー）を維持し、再生能力・病気耐性に優れています。
- 一部の研究によると、4 歳以降、DNA メチル化に基づく「エピジェネティック・クロック」が停止し、老化指標がほとんど進まなくなる可能性が示唆されました。
- この現象は“若さの維持”というユニークなアンチエイジング戦略として、人間の老化研究や再生医療に大きなヒントを提供するかもしれません。

（※この記事はプレプリント掲載研究に基づく内容であり、査読付き論文ではありません。）

元論文（プレプリント）の内容概要と出典

- **プレプリントタイトル（推定）**：おそらく「Axolotl epigenetic clocks offer insights into the nature of ...」
- **要旨内容**：
  - アホロートルは優れた再生能力と長寿性、老化関連疾患への耐性を持つ。
  - DNA メチル化パターンを用いてエピジェネティック・クロックを構築すると、4 歳以降に老化速度が著しく鈍化／停止する特徴が観察された。
  - この結果は、「老化プロセスの緩やかさ」が再生能力や寿命と関連する可能性を示唆している。
- **出典**：
  - プレプリント投稿：2024 年 9 月
  - 出典情報：タイトル「Axolotl epigenetic clocks offer insights into the nature of ...」、bioRxiv に掲載（バージョン v1: 2024 年 9 月 10 日; v2: 10 月 1 日） BioRxiv+1

参考文献リスト

1. 「Axolotl epigenetic clocks offer insights into the nature of ...」, bioRxiv プレプリント, 2024 年 9 月（Ver.1）および 10 月（Ver.2）投稿 BioRxiv+1
2. National Geographic 特集記事「Axolotls stop aging after four years — could studying them help humans live longer?」, 2024 年 10 月 National Geographic
3. HowStuffWorks 記事「Axolotl life expectancy is roughly halved in the wild」, 2025 年 7 月 — 飼育下で 10–15 年、野生で約 5 年という寿命データ HowStuffWorks
4. A–Z Animals ウェブ記事「The Axolotl Lifespan: Captive vs. Wild」, 2025 年 7 月 — 上記と同様の寿命差記述 A-Z Animals

---

まとめ（表形式）

項目	内容
プレス（記事）	ナショナルジオグラフィック「4 歳を過ぎると老化時計が止まる可能性」
基盤となる研究	エピジェネティック・クロック分析によるエビデンス（bioRxiv プレプリント, 2024 年）
飼育 vs 野生の寿命	飼育下：10–15 年（最大 25 年）、野生：平均 5 年程度（HowStuffWorks、A–Z Animals）
参考文献	上記 4 件を参照

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

アホロートル（*Ambystoma mexicanum*）の「エピジェネティック・クロック（DNA メチル化に基づく生物学的年齢指標）が若年期（およそ 4 歳）以降ほぼ進行しない＝“止まる／非常に遅くなる”可能性」まとめ

1) プレス報道の要点（要約）

複数のメディアが、アホロートル（*Ambystoma mexicanum*）の「エピジェネティック・クロック（DNA メチル化に基づく生物学的年齢指標）が若年期（およそ 4 歳）以降ほぼ進行しない＝“止まる／非常に遅くなる”可能性」を報じました。報道は、この現象がアホロートルの長寿性・高い再生力（negligible senescence）と関連するかもしれない、という驚きと期待を強調しています。主要な報道例としては National Geographic の特集記事があり、研究の共同執筆者の一人である Steve Horvath らのコメントも取り上げられています。National GeographicX (formerly Twitter)

（短評）報道は「インパクトの大きい仮説」を強調する一方で、元データがプレプリントで査読前である点や、エピジェネティック・クロックの解釈の難しさについての慎重な説明は必ずしも詳細ではありません。National GeographicThe Company of Biologists Journals

---

2) 元論文（プレプリント）の要約 —《Axolotl epigenetic clocks offer insights into the nature of negligible senescence》

（著者の一部：Y. Haluza, J. A. Zoller, A. T. Lu, H. E. Walters, M. Lachnit, R. T. Brooke, N. Park, M. H. Yun, S. Horvath 他 ／ bioRxiv, 2024.09.09; DOI: 10.1101/2024.09.09.611397）。BioRxiv

目的

アホロートル全生涯にわたる DNA メチル化（メチローム）プロファイルを解析し、メチレーションに基づく「エピジェネティック・クロック」を構築して、アホロートルが示す“老化（epigenetic ageing）”の特徴を調べること。特に「再生能」と「エピジェネティック年齢」の関係を検証することが主眼。

サンプル／データ（要点）

- 組織横断的（pan-tissue）および組織別の DNA メチレーションデータを、幼体から高齢個体までのサンプルで収集（論文内に詳細なサンプル数・年齢レンジ・組織種の表あり）。BioRxiv
- メチル化測定は CpG 単位のプロファイリングを使用（プレプリント本文に用いた実験法の詳細および解析パイプラインの記述あり）。BioRxiv

解析手法（要旨）

- CpG データを用いて機械学習的アプローチ（例：penalized regression / elastic net 等）でエピジェネティック・クロックモデル（年齢予測モデル）を構築。組織特異型と汎用（pan-tissue）型のクロックを作成。BioRxiv
- メチレーション変化が局在するゲノム領域（例：Polycomb Repressive Complex 2 — PRC2 標的領域）や再生時のメチレーション動態についても解析。BioRxiv

主要結果

1. **二相性（biphasic）のクロック**：構築したクロックは「早期の年齢（幼年期・若年期）には年齢を予測するが、ある時点（およそ 4 歳）以降は予測精度が著しく低下する」──つまりクロックが“止まる/鈍化する”ように見える。BioRxiv
2. **メチロームの安定性**：全般的にアホロートルのメチロームは生涯を通じて比較的安定しており、特に PRC2 標的サイトでの加齢に伴うメチレーション変化が他の脊椎動物に比べて小さい。BioRxiv
3. **再生時の局所的若返り（rejuvenation）イベント**：一部の組織において、断片再生後に局所的なメチレーションの“若返り”が観察され、再生プロセスがエピジェネティックな若返りを引き起こす可能性を示唆。BioRxiv

解釈

- アホロートルは「初期には進行性のエピジェネティック老化を示すが、その後メチロームの変化がほとんど止まる」ように見える — これが「negligible senescence（無視できる老化）」の分子的特徴の一端である可能性。再

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

生プロセスとエピジェネティック制御の相互作用が、老化の遅延に寄与している可能性が示唆された。BioRxiv

図・データの要旨（読者が確認すべき点）

- 年齢 vs 予測年齢プロット（クロックの biphasic 性を示す散布図／残差プロット）。BioRxiv
- 組織別のメチレーション変化ヒートマップと PRC2 標的領域の安定性を示す図。BioRxiv
- 再生前後での局所 CpG の若返りを示す局所的プロファイル（再生組織の時系列データ）。BioRxiv

---

3) 批評・限界点・研究者・メディアの反応

批評・限界

- **査読前のプレプリント**である点：結果は興味深いが、査読による方法論・解析の厳密性確認が今後必要。BioRxiv
- **クロックの解釈の難しさ**：エピジェネティック・クロックが「老化そのもの」を直接測るのか、ある種の発達的プログラムや細胞組成の変化を反映しているのかは論争がある。したがって「クロックが止まる＝老化が止まる」と単純に解釈することは危険。専門家解説でも注意が促されています。The Company of Biologists JournalsBioRxiv
- **サンプル数・年齢範囲・環境要因**：飼育個体に偏る、野外個体のサンプルが少ない、あるいは環境（飼育条件）でメチレーションが変わりうる点は注意が必要。これらは論文本文で議論されています。BioRxiv

研究者・メディアの反応

- Steve Horvath をはじめ複数の研究者が本成果に興味を示し、メディア（National Geographic 等）が広く取り上げたことで注目度が高まりました。ただし専門家は「さらなる独立検証」と「生理学的老化指標（寿命・機能低下）の対応付け」を求めています。National GeographicX (formerly Twitter)The Company of Biologists Journals

---

4) 参考文献（主要ソース）

以下は本要約で用いた主要情報源です（出典表示は web.run の返却 ID を併記します）。必要であれば、これらの完全な書誌情報（著者一覧・doi・PDF）を APA 等の形式で整えます。

1. Haluza Y., Zoller J. A., Lu A. T., Walters H. E., Lachnit M., Lowe R., Haghani A., Brooke R. T., Park N., Yun M. H., Horvath S. (2024) *Axolotl epigenetic clocks offer insights into the nature of negligible senescence.* bioRxiv preprint. DOI: 10.1101/2024.09.09.611397. BioRxiv
2. National Geographic (2024) *How axolotls' secret to pausing aging could change medicine.* (feature article; Oct 18, 2024). National Geographic
3. Development (The Company of Biologists) (2025) *In preprints: an epigenetic clock that does not tick.* — commentary discussing the preprint and interpretational issues. The Company of Biologists Journals
4. ScienceCast / ResearchHub / social-science summaries and discussion pages summarizing the bioRxiv preprint. ScienceCastResearchHub
5. Related discussion and media reposts (tweets by authors/researchers, Reddit threads) capturing community reaction. X (formerly Twitter)Reddit

Summary: Epigenetic Clock “Stops” in Axolotl Aging

1) Press Coverage (Summary)

Several media outlets reported that the axolotl (*Ambystoma mexicanum*) shows a remarkable phenomenon: its **epigenetic clock (DNA methylation–based biological age indicator) appears to nearly stop or slow**

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

dramatically after around four years of age. Reports emphasized the link between this phenomenon, axolotl longevity, and its extraordinary regenerative capacity, suggesting that this may represent a form of negligible senescence.

One of the main features appeared in **National Geographic**, which also cited co-author Steve Horvath's comments.

(Note: While the story highlights a striking hypothesis, it is based on a preprint that has not yet undergone peer review. Media coverage tends to emphasize the dramatic aspect, while technical caveats and interpretational complexity are less detailed.)

---

2) Preprint Summary — *Axolotl epigenetic clocks offer insights into the nature of negligible senescence*

(Authors include: Y. Haluza, J.A. Zoller, A.T. Lu, H.E. Walters, M. Lachnit, R.T. Brooke, N. Park, M.H. Yun, S. Horvath, et al. / bioRxiv, Sept 9, 2024; DOI: 10.1101/2024.09.09.611397)

Objective

To analyze DNA methylation (methylome) profiles across the entire lifespan of the axolotl and build methylation-based **epigenetic clocks**, in order to characterize patterns of epigenetic aging and to investigate the relationship with regeneration.

Samples & Data

- Pan-tissue and tissue-specific DNA methylation datasets covering individuals from juvenile to aged stages.
- CpG-level profiling with detailed methodological descriptions provided in the preprint.

Methods (Outline)

- Constructed age-prediction models (epigenetic clocks) using CpG data and machine learning approaches (e.g., penalized regression/elastic net).
- Built both pan-tissue and tissue-specific clocks.
- Analyzed genomic regions enriched for age-related methylation changes (e.g., Polycomb Repressive Complex 2 [PRC2] targets) and examined regeneration-associated dynamics.

Key Results

1. **Biphasic clock dynamics**: The clocks predict chronological age during early life but lose predictive accuracy around **4 years of age**, suggesting that the epigenetic clock "stops" or slows dramatically.
2. **Methylome stability**: The axolotl methylome remains relatively stable across life, with smaller age-related changes at PRC2 target sites compared with other vertebrates.
3. **Regeneration-linked rejuvenation**: Some tissues displayed local "rejuvenation" events in CpG methylation following regeneration, hinting at epigenetic resetting during tissue regrowth.

Interpretation

- Axolotls exhibit **progressive epigenetic aging in early life**, followed by a near-halting of methylation changes later — consistent with negligible senescence.
- Regenerative processes may interact with epigenetic control to slow or prevent age-associated decline.
-

Figures to Note

- Age vs. predicted age scatterplots showing biphasic dynamics.
- Heatmaps of tissue-specific methylation stability and PRC2-targeted CpGs.
- Time-series profiles showing rejuvenation during regeneration.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

3) Critique, Limitations, and Community Response

Limitations

- **Preprint status**: Findings are preliminary and require peer review.
- **Interpretation of clocks**: Epigenetic clocks may reflect developmental programs or cell composition shifts rather than “aging” per se; thus, equating “clock stoppage” with “aging stoppage” is problematic.
- **Sample scope**: Data largely from captive individuals; wild axolotl methylomes are underrepresented. Environmental effects (e.g., rearing conditions) could influence results.

Responses

- Researchers including Steve Horvath have expressed excitement at the findings, and coverage in **National Geographic** and other outlets boosted attention.
- Experts urge caution, calling for **independent replication** and linking epigenetic results with physiological markers of aging (lifespan, decline in function).

---

4) References (Key Sources)

1. Haluza Y., Zoller J.A., Lu A.T., Walters H.E., Lachnit M., Lowe R., Haghani A., Brooke R.T., Park N., Yun M.H., Horvath S. (2024). *Axolotl epigenetic clocks offer insights into the nature of negligible senescence.* bioRxiv preprint. DOI: 10.1101/2024.09.09.611397.
2. **National Geographic** (2024). *How axolotls' secret to pausing aging could change medicine.* Feature article, Oct 18, 2024.
3. **Development (The Company of Biologists)** (2025). *In preprints: an epigenetic clock that does not tick.* Commentary on the bioRxiv preprint.
4. ScienceCast / ResearchHub community summaries and discussions of the preprint.
5. Social media discussions and reposts (e.g., X/Twitter, Reddit), reflecting broader scientific and public reactions.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

ハルラボの知見


1). 300 グループにアホロートルの寿命と飼育情報を聞き取りし、その結果を統計処理してみた。

結果 → https://researchmap.jp/HAL_lab_Axolotl/published_papers/51108485

相関係数（抜粋）

	餌頻度(回/week)	飼い主年齢層(代)	寿命(年)	体長(cm)	体調不良回数
餌頻度		-0.04	-0.07	-0.08	**0.42**
飼い主年齢層	-0.04		0.08	0.09	0.11
寿命	-0.07	0.08		0.01	**0.44**
体長	-0.08	0.09	0.01		-0.06
体調不良回数	0.42	0.11	**0.44**	-0.06	


「餌頻度」と「体調不良回数」（r=0.42）、
「寿命」と「体調不良回数」（r=0.44）
などで比較的強めの正の相関が見られた。



補足図 1: 飼育にまつわる情報の相関関係を示すヒートマップ

- 餌頻度（回/week）→ **Feed frequency (times/week)**
- 飼い主の年齢層（代）→ **Owner age group (decade)**
- 寿命（年）→ **Lifespan (years)**
- 体長（cm）→ **Body length (cm)**
- 体調不良回数 → **Number of illness episodes**

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl



補足図 2: 寿命と体調不良回数の関係を示す散布図


- **X 軸**: Lifespan (years) → 寿命（年）
- **Y 軸**: Number of illness episodes → 体調不良回数
- **タイトル**: Scatter plot of Lifespan vs Number of illness episodes → 寿命と体調不良回数の散布図

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

この結果を寿命で補正すると…？

方法①: 年あたり体調不良率（正規化）

各個体の「体調不良回数 ÷ 寿命」により、1 年あたりの体調不良率を算出した。
この指標を用いることで、寿命の長短に左右されない健全性の比較が可能となる。



- X 軸: **Lifespan (years)** → 寿命（年）
- Y 軸: **Illness episodes per year** → 年あたり体調不良回数
- タイトル: **Scatter plot (normalized by lifespan)** → 寿命補正後の体調不良率散布図

- 寿命が長い個体でも「生きていた年数に比べてどのくらい体調不良が多いか」を直接比較できる。
- 仮に寿命が短い個体でも、1 年あたりの体調不良率が高ければ「相対的に不健康だった」ことを示唆する。
- 逆に寿命が長くても年あたりの体調不良率が低ければ「健康に長生きした」と解釈できる。

注意点

年あたりに換算しているため、「寿命が短いから体調不良回数も少ない」効果をうまく補正できるが、
年ごとの変動（例：老齢期に偏って起こる体調不良）は平均化されてしまう。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

方法①: 年あたり体調不良率（正規化）の結果から読み取れること

1. **縦軸（Illness episodes per year）**が「寿命の長さに依存しない健全性の指標」になっている。
   → 短命だから不調が少なく見える、という単純な影響を除外できている。
2. **横軸（Lifespan years）**が 2〜10 年まで分布しているが、縦軸の分布には強い傾向は見られない。
   → つまり**「寿命の長い個体だから必ずしも年あたりの不調が多い／少ない」とは言えない**。
3. 1 年あたり **0〜7 回程度**と幅広い。
   → 個体差が大きく、飼育条件や個体要因（餌、環境、遺伝など）の影響が強いことを示唆する。

解釈のまとめ

- この正規化により「寿命の長短に依存しない体調不良の頻度」を比較できる。
- 結果として、**寿命の長さとは独立に、健康状態の個体差が大きい**ことが明らかになっている。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

方法②: 回帰分析による補正（残差プロット）

体調不良回数 ~ 寿命 の回帰分析を行い、その残差を算出した。

この残差は「寿命の影響を取り除いた体調不良の多さ」を意味する。



- X 軸: **Lifespan (years)** → 寿命（年）
- Y 軸: **Residual illness episodes** → 体調不良回数の残差
- タイトル: **Scatter plot (residuals after regression)** → 回帰補正後の残差散布図

回帰モデルの要約（抜粋）

- **$R^2 = 0.192$**（寿命が体調不良回数の約 19%を説明している）
- **$p < 0.001$**（統計的に有意な関係が認められる）

- **やっていること**

「体調不良回数」を「寿命」で回帰し、寿命によって説明される部分を取り除いて残差を算出。

- **意味すること**
  - 残差が正（＋）なら「寿命に比べて体調不良が多い個体」、残差が負（−）なら「寿命に比べて体調不良が少ない個体」を意味する。
  - つまり、「寿命に見合った体調不良の回数」からのずれを定量化できる。
  - $R^2 = 0.192$ → 寿命が体調不良回数の約 19% を説明しているに過ぎず、それ以外の要因（飼育環境、餌、遺伝的要因など）が大きいことを示唆する。

注意点

回帰モデルは線形を仮定しているため、もし「寿命が極端に長い個体で急に体調不良が増える」といった非線形パターンがあれば、この方法では捉えきれない。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

方法②: 回帰分析による補正（残差プロット）の結果から読み取れること

図の意味

- **横軸（Lifespan years）**: 個体の寿命。
- **縦軸（Residual illness episodes）**: 回帰分析によって寿命の影響を取り除いた「体調不良回数の残差」。
- 赤の破線（0 ライン）: 「寿命から予測される体調不良回数」と一致するライン。

解釈

1. **残差が 0 付近**
   → 寿命の長さから見て「妥当」な体調不良回数を示す個体。
2. **残差が正（＋）に大きい個体**
   → 「寿命に比べて体調不良が多すぎる」個体。
   → 長生きしていても、不調の頻度が過剰であることを意味する。
3. **残差が負（−）に大きい個体**
   → 「寿命に比べて体調不良が少ない」個体。
   → 健康的に長生きした、もしくは短命でも体調不良は少なかった可能性を示唆。
4. **全体的な傾向**
   - **プロットは 0 を中心に広がっており、寿命だけでは体調不良回数を完全には説明できない。**
   - **実際に先の回帰モデルの結果では $R^2=0.192$ であり、寿命は体調不良回数の約 19%しか説明していない。**
   - **つまり、残りの大部分は 飼育環境・餌・遺伝的背景など、寿命以外の要因に依存している。**

まとめ


- この残差プロットは「寿命を揃えて補正した後に、どの個体が相対的に健康／不健康だったか」を示す。
- 今後はこの残差を用いて **群比較（餌の種類や飼育条件ごと）** を行うことで、「寿命の長短に依存しない健康度」を条件間で検証できる。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

寿命カテゴリ別

短命群・中間群・長命群で体調不良回数の分布を比較。

→「寿命が長い個体は、単純な回数としては体調不良も増えやすいのか」を確認できる。



寿命カテゴリ（三分位）別の体調不良回数

Boxplot: Lifespan categories vs Illness episodes

- **X-axis**: Lifespan category → 寿命カテゴリ（短命群・中間群・長命群）
- **Y-axis**: Number of illness episodes → 体調不良回数
- **Title**: Boxplot: Lifespan categories vs Illness episodes

  → 箱ひげ図: 寿命カテゴリ別の体調不良回数

傾向

- 寿命が「短命群 → 中間群 → 長命群」となるにつれて、体調不良回数の中央値が増える傾向がある。
- これは「寿命が長いと、当然その分だけ体調不良の機会も増える」ことを反映している。

解釈


- **寿命そのものが体調不良の「総回数」に強く影響している。**
- ただし、長命群の中にも体調不良が少ない個体、短命群でも体調不良が多い個体が存在しており、**寿命だけでは個体の健康状態を完全には説明できない**。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

餌頻度カテゴリ別

餌やり頻度の低・中・高で体調不良回数の分布を比較。

→「餌を与える頻度が体調不良の多さに関与しているか」を確認できる。



餌頻度カテゴリ（三分位）別の体調不良回数

Boxplot: Feeding frequency categories vs Illness episodes

- **X-axis**: Feeding frequency category → 餌頻度カテゴリ（低頻度群・中頻度群・高頻度群）
- **Y-axis**: Number of illness episodes → 体調不良回数
- **Title**: Boxplot: Feeding frequency categories vs Illness episodes

→ 箱ひげ図: 餌頻度カテゴリ別の体調不良回数


傾向

- 餌やり頻度が「低頻度群 → 中頻度群 → 高頻度群」となるにつれて、体調不良回数の中央値がやや増える傾向がある。
- 分布のばらつきも大きく、特に高頻度群では外れ値（極端に体調不良回数が多い個体）が目立つ。

解釈

- **餌を与える頻度が高いほど体調不良の機会が増える可能性を示唆する。**
- しかし「餌頻度が高い＝不健康」という単純な因果関係とは限らず、餌の種類、飼育密度、環境条件などの交絡要因が関与している可能性が高い。
- 餌頻度は一因子であり、総合的な飼育管理条件と併せて解釈する必要がある。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl



補足図 3: 成長率と寿命の相関

結果・考察

1. 成長率の傾向

- **正常型**や**成長不全 (PG)** は成長率が低く、特に PG は著しく低値。
- 一方、**肥大成長 (OG)** 条件（糖質摂取、高脂肪食、コラーゲン摂取、FGF/FGFR 処理など）は軒並み成長率が高い（15～20 程度）。
- 特に「OG + FGF + FGFR (mix)」「OG + hGF 摂取」などは成長率が最も高く、通常の成長を大きく上回っている。

👉 **介入により成長促進は明らかに可能である**ことが分かる。

---

2. 寿命の傾向

- **正常型**が最も長寿（平均 3.5 年程度）。
- **成長不全 (PG)** も比較的長寿で、正常型よりやや短いが 2.5～3 年程度。
- **肥大成長 (OG)** 条件では寿命が明確に短縮し、いずれも 2 年以下。
- 特に**高脂肪食**や **FGF 関連処理**では寿命が 1～1.5 年と大きく減少している。

👉 **成長促進は寿命の短縮を伴う可能性が高い**。

---

3. 成長と寿命の相関

- 「成長率が高い群ほど寿命が短い」という**負の相関関係**が見られる。
- 正常群：成長率は低いが長寿
- 肥大成長群：成長率は高いが短命
- 成長不全群：成長率は低いが比較的長寿

👉 成長の速さや過剰促進が、長期的には生存コスト（代謝負担や疾患リスク）につながっている可能性。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. 生物学的解釈

1. **エネルギーバランスの破綻**
   高成長＝高代謝負荷 → 酸化ストレスや細胞老化の促進。
2. **シグナル伝達過剰**
   FGF/FGFR 刺激のような強い成長促進シグナルが、長期的には腫瘍化や老化関連経路を活性化し、寿命を縮める可能性。
3. **「寿命と成長のトレードオフ」**
   生物一般に見られる原則（速く育つ種は寿命が短い）と一致。

まとめ

- **成長率と寿命の間には明確な負の相関がある。**
- 過剰な成長促進（高脂肪食や FGF 刺激など）は、短期的には体サイズ増加に有効だが、寿命を大幅に縮めるリスクがある。
- 正常型や成長不全型は成長は控えめだが寿命は長い →「適度な成長が長寿と健康維持に有利」という示唆。


本文中の肥大成長(OG)は健康的な育ち方を意味しておらず、明らかに規格外の成長度のものを指します。

つまり本文で実施している試験は、肥大成長する個体に、他の成長因子をプラスしています。

また、遺伝的な部分は、自然突然変異のものを掛け合わせています。
ANK1/ANKHD1 活性型 homo については、この変異個体が肥大成長してるという意味です。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

散布図（成長率 vs 寿命）**にして相関係数を可視化



結果

成長率と寿命の間には強い負の相関が認められた（相関係数 Pearson's r = -0.84）。

すなわち、成長が速い個体ほど寿命が短い傾向が明確に示された。

- 横軸（Growth Rate, Ratio）：成長率を示す。右に行くほど成長が速い。
- 縦軸（Lifespan, Years）：寿命を示す。上に行くほど寿命が長い。
- 結果として「成長率が高い群ほど寿命が短い」という**負の相関**が可視化されています。
  - **Normal**（正常型）は成長率が低めだが寿命は最長。
  - **Growth Defect (PG)** も成長は遅いが比較的長寿。
  - **Hypertrophy (OG) 系列**は成長率が高いが寿命は短縮。特に「OG+FGF+FGFR」は極端に短命。

この散布図から、**「急速な成長は寿命の犠牲を伴う」という生物学的トレードオフ**がより明確に見て取れます。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【総説】魚類・両生類の化学物質に対する耐性

Review: Cognition and Behavior in Fish and Amphibians

著: 杉山 遥[1]; 竜岡 竜騎[5], 阿保郎 太朗[6], 横江 誼衡[1, 2, 3, 4]; F.N. 三太子[1]


[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー,

[5.] トラフサンショウウオラボ, [6.]ハルラボ・ファーム

Authors: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]; F.N. Santako[1]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ,

[4] SciMountechnology, [5] Ambys Lab, [6] Hal lab farm



※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

要旨

アホロートルを含む両生類の上皮は粘膜によって保護されているが、有機溶剤をはじめとする化学薬品に対しては脆弱である。そのため、環境汚染は彼らの生息域に深刻な影響を及ぼし得る。特に飼育水に皮膚腐食性の高い有機溶剤が混入した場合、その影響は顕著である。また、飼育容器のプラスチックから溶出する添加材（いわゆる環境ホルモン）の作用も無視できない場合がある。本総説では、両生類および魚類における耐薬品性に関する知見を整理し、彼らの成育環境を守る重要性について論じる。

Abstract

The epithelium of amphibians, including axolotls, is protected by mucus but remains highly vulnerable to chemical agents such as organic solvents. Consequently, environmental pollution can exert profound effects on their habitats. In particular, exposure to corrosive organic solvents in rearing water leads to severe impacts, while leaching of plastic additives (so-called endocrine disruptors) from containers may also play a non-negligible role. This review summarizes current knowledge on chemical resistance in amphibians and fish, emphasizing the importance of safeguarding their developmental environments.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

両生類の薬品耐性

両生類や魚類は陸上動物に比べて **環境中の化学物質（農薬、重金属、環境ホルモンなど）に曝露されやすい** 生態系におり、それに対する耐性や感受性は広く研究されている。

---

1. 一般的な特徴

- **皮膚・鰓からの直接吸収**
  両生類は皮膚透過性が高く、魚類は鰓を介して水中の化学物質を容易に取り込みます。そのため化学物質曝露の影響を受けやすい一方で、独自の解毒機構を発達させています。
- **代謝酵素系（Phase I/II 酵素）**
  - **シトクロム P450 (CYP)**：有機汚染物質や農薬の酸化代謝を担う。
  - **グルタチオン S-トランスフェラーゼ (GST)、UDP-グルクロン酸転移酵素 (UGT)**：抱合反応によって水溶性を高め、排出を促す。
  - 種によって誘導能や発現レベルが異なり、耐性の差を生みます。

---

2. 魚類における耐性と感受性

- **PCB・ダイオキシン類**
  AHR（アリル炭化水素受容体）を介した毒性に感受性がある一方、**Fundulus heteroclitus（ミジンコハゼ科）**のように、汚染河口域で AHR 経路に変異が生じ、強い耐性を獲得した個体群が知られています。
- **重金属（カドミウム・水銀・鉛）**
  魚類はメタロチオネイン（MT）を発現して重金属をキレートし、細胞内毒性を緩和します。種ごとの発現能差が耐性に関与。
- **農薬・殺虫剤（有機リン、ピレスロイド）**
  コリンエステラーゼ阻害作用に感受性が高い種が多いが、コイ科やティラピアの一部では分解酵素の発達で耐性を示す例あり。

---

3. 両生類における化学物質影響

- **環境ホルモン（内分泌かく乱物質）**
  - アトラジン（除草剤）：カエルの性分化に影響し、雄の雌化や生殖腺異常を誘発する報告（Xenopus laevis）。
  - ビスフェノール A、ノニルフェノール：甲状腺ホルモンシグナルに干渉し、変態や発達に影響。
- **重金属・酸性雨関連物質**
  幼生は酸性化や金属イオンに弱く、成長遅延や奇形が報告される。特に皮膚呼吸依存度が高い幼生期は感受性が強い。
- **耐性事例**
  特定の農薬や重金属に長期間さらされてきた地域個体群では、代謝酵素や輸送体の発現が恒常的に高くなり、一定の耐性が形成される。

---

4. 共通する特徴と進化的意義

- **高い可塑性**：水質の変動環境に適応するため、代謝酵素やストレス応答遺伝子の誘導能力が進化的に発達。
- **発生段階による感受性の差**：卵・幼生は透過性が高く、成体より感受性が強い。
- **個体群レベルの進化的耐性**：汚染地域の集団では世代を経て耐性変異が固定化する例がある。

まとめ

両生類・魚類は **解毒酵素系や金属結合タンパク**によってある程度の化学物質耐性を持つが、環境ホルモンや水質汚染には脆弱な面が多い。特に **発生期の影響は顕著**で、生態系や種の存続に直結する。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Chemical Resistance in Amphibians

Amphibians and fish inhabit ecosystems where they are more frequently exposed to environmental chemicals (e.g., pesticides, heavy metals, endocrine disruptors) compared to terrestrial animals. Their tolerance and sensitivity to such substances have been extensively studied.

---

1. General Characteristics

- **Direct absorption through skin and gills**
  Amphibians have highly permeable skin, while fish readily take up chemicals through the gills. This makes them particularly vulnerable to chemical exposure, yet they have also evolved unique detoxification mechanisms.
- **Metabolic enzyme systems (Phase I/II enzymes)**
  - **Cytochrome P450 (CYP):** Catalyzes oxidative metabolism of organic pollutants and pesticides.
  - **Glutathione S-transferases (GST), UDP-glucuronosyltransferases (UGT):** Enhance water solubility of xenobiotics via conjugation, facilitating excretion.
  - Species differences in inducibility and expression levels of these enzymes contribute to variability in chemical resistance.

---

2. Resistance and Sensitivity in Fish

- **PCBs and dioxins**
  Toxicity is mediated by the aryl hydrocarbon receptor (AHR). Notably, populations of *Fundulus heteroclitus* inhabiting polluted estuaries have evolved mutations in the AHR pathway, conferring high resistance.
- **Heavy metals (cadmium, mercury, lead)**
  Fish express metallothioneins (MT), which chelate metals and mitigate intracellular toxicity. Species-specific differences in MT expression influence resistance.
- **Pesticides and insecticides (organophosphates, pyrethroids)**
  Many fish species are highly sensitive due to cholinesterase inhibition, although some cyprinids and tilapias exhibit resistance through enhanced detoxifying enzymes.

---

3. Chemical Effects in Amphibians

- **Endocrine-disrupting chemicals (EDCs)**
  - **Atrazine (herbicide):** Alters sex differentiation in frogs, inducing feminization and gonadal abnormalities (e.g., *Xenopus laevis*).
  - **Bisphenol A, nonylphenol:** Interfere with thyroid hormone signaling, affecting metamorphosis and development.
- **Heavy metals and acid rain-related compounds**
  Larvae are particularly vulnerable, with reports of delayed growth and malformations. Their strong reliance on cutaneous respiration increases sensitivity.
- **Cases of resistance**
  In populations chronically exposed to pesticides or heavy metals, constitutively elevated expression of metabolic enzymes and transporters has been observed, conferring adaptive resistance.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. Shared Features and Evolutionary Significance

- **High plasticity:** To cope with fluctuating aquatic environments, amphibians and fish have evolved inducible metabolic and stress-response genes.
- **Developmental stage–dependent sensitivity:** Eggs and larvae exhibit greater sensitivity than adults due to higher permeability.
- **Population-level evolutionary resistance:** In polluted habitats, resistant mutations can become fixed over generations.

Conclusion

Amphibians and fish possess a degree of chemical resistance through detoxifying enzyme systems and metal-binding proteins. However, they remain highly vulnerable to endocrine disruptors and aquatic pollution. The pronounced sensitivity during developmental stages has direct implications for ecosystem stability and species survival.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

薬品の毒性に関する用語: IC_{50}＆LD_{50}

あらゆる脊椎動物に関わる「IC_{50}」「LD_{50}」 について、まず用語説明→測定法／統計→種間差の要因→実務的な解釈・外挿法→規制・代替法までを整理した詳細まとめです。学術レビューやリスク評価資料にそのまま使えるよう、専門用語の定義や計算法式、注意点も含めて書いてあります。

1. 用語の定義（基本）

- **IC_{50}（半数阻害濃度：Inhibitory Concentration 50%）**
  ある生物学的反応（酵素活性、受容体結合、細胞生存率など）を **50%抑制する薬物や化学物質の濃度**。通常は in vitro アッセイ（細胞、酵素、受容体結合試験）で求められることが多く、単位は濃度（例：μM、nM、$mg \cdot L^{-1}$）。**時間依存性・アッセイ条件依存性**が強い点に注意。
- **LD_{50}（半数致死量：Lethal Dose 50%）**
  一定の投与条件（単回投与など）で **投与群の 50%が死亡する用量**。通常は in vivo（実際の動物）試験の指標で、単位は体重当たり（例：$mg \cdot kg^{-1}$）。急性毒性の標準指標としてよく使われる。
- **LC_{50}（半数致死濃度：Lethal Concentration 50%）**
  水棲生物などで用いられる。一定時間（水中暴露 24/48/96 h 等）における**水中濃度**で 50%が死亡する値（$mg \cdot L^{-1}$）。
- **EC_{50}（半数有効濃度）・NOAEL/LOAEL・BMD（benchmark dose）** 等は LD_{50}/IC_{50}の周辺で使われる関連指標。

2. IC_{50}（測定法・解析・注意点）

測定法

- 一般は複数濃度での反応（%抑制や%生存など）を測定し、**濃度—応答曲線** を作成する。
- 応答は**非線形回帰（Hill 式など）**でフィッティングして IC_{50} を推定する。

代表的モデル式（Hill 方程式）

E([I])=Emin+Emax−Emin1+(IC50[I])nE([I]) = E_{min} + ¥frac{E_{max}-E_{min}}{1 + ¥left(¥frac{IC_{50}}{[I]}¥right)^n}E([I])=Emin+1+([I]IC50)nEmax−Emin

- E([I])E([I])E([I])：濃度[I]での応答
- nnn：Hill 係数（曲線の傾き）
- Emin,EmaxE_{min}, E_{max}Emin,Emax：最小・最大応答値

Ki（阻害定数）との関係（Cheng–Prusoff 式）

競合阻害の場合、IC_{50}は検査条件（基質濃度 [S], Km）に依存する：

Ki=IC501+[S]KmK_i = ¥frac{IC_{50}}{1 + ¥frac{[S]}{K_m}}Ki=1+Km[S]IC50

→ IC_{50} を Ki に変換すれば競合阻害の本質的親和性を評価できる（ただし前提が重要）。

注意点（IC_{50}の解釈で重要）

- **アッセイ条件依存性**：基質濃度、反応時間、細胞数、代謝活性、pH、溶媒（DMSO など）で大きく変わる。
- **単位と時間**：短時間 IC_{50} と長時間（24–72 h）の IC_{50} は異なる。
- **Hill 係数**で作用様式（協同性など）を示唆できるが、証明は別途必要。
- **IC_{50} ≠ 安全濃度**：in vitro 値は in vivo 動態（吸収・分布・代謝・排泄）を考慮しない。

3. LD_{50}（測定法・解析・注意点）

測定法（伝統的）

- 動物に単回投与（経口、皮下、静注など）し、一定期間（通常 14 日）死亡率を観察。複数用量群で死亡率を求める。
- 死亡率データを**対数用量—応答変換**（probit 法など）でフィッティングし LD_{50} を推定。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

代表的解析手法

- **Probit 法**（Mortality → probit に変換し線形回帰）：古典的で多くの規格が採用。
- **Logit 回帰（ロジスティック）**：尤度法でパラメータを推定する方法。
- **Spearman-Karber 法**：単純で非線形フィッティングの代替。
- **信頼区間（95% CI）** を必ず報告するのが望ましい。

単位・条件

- LD_{50} は通常 **mg·kg⁻¹ bw**（体重当たり）で報告。
- 投与経路（経口、経皮、吸入、静注など）で LD_{50} が大きく変わる。
- **時間軸**（観察日数）を明記（24 h, 48 h, 14 d 等）。

注意点（LD_{50}の解釈）

- **種・年齢・性別・状態（病気や栄養状態）** によりばらつく。
- **製剤・溶媒・添加剤** の影響（溶媒が毒性を増減させる）を受ける。
- LD_{50} はあくまで **急性致死性の指標**。慢性毒性、生殖毒性、発がん性、神経毒性など別評価が必要。
- 動物福祉観点から LD_{50}試験は近年縮小・代替が推進されている（OECD 指針の改訂等）。

4. 脊椎動物間での差（なぜ種ごとに異なるか）

- **薬物動態（ADME）の違い**
  - 吸収経路（皮膚透過 vs 鰓 vs 肺 vs 消化管）
  - 分布（体脂肪率、結合蛋白）
  - 代謝（CYP 酵素ファミリー、GST、UGT、SULT などの種類・発現量）
  - 排泄（腎機能・胆汁排泄等）
- **薬力学的差**
  - 標的（受容体・酵素）の配列差・発現差 → 親和性や感受性が変わる
- **生理的差**
  - 体温（変温動物 vs 恒温動物）や代謝速度
  - 呼吸様式（皮膚呼吸依存度の違い＝両生類幼生の高感受性）
- **生活史・生態的要因**
  - 水生 vs 陸生や食性の違い（摂取・曝露経路が異なる）
- **個体差**（年齢、性別、遺伝的背景）
- **環境要因**（水温、pH、溶存酸素、共存化学物質）

5. 種間外挿（異種→異種 / 動物→人 へ）の方法と限界

単純手法

- **体重当たり換算（mg·kg⁻¹）**：同一 mg·kg⁻¹ で等しく作用するという仮定は一般に不適切。
- **体表面積（BSA）換算（Km 法）**：ヒト換算でよく用いられる（動物種ごとに Km 値を用いる）。簡便だが個体差・代謝差を吸収できない。

より高度な手法

- **Allometric scaling（体重^b）**：代謝速度は体重のべき則に従う（b ≈ 0.75 等）。薬物動態パラメータ（CL, V）をスケールする手法。
- **PBPK（生体内動態モデル）**：種ごとの生理学的パラメータと化学物質の物性を組み合わせ、内部曝露（臓器濃度）を推定する最も理論的。外挿に有用だがパラメータ同定が必要。
- **不確実性（安全）係数**：規制評価で用いられる（例：動物→ヒトの不確実性係数 10、個体内差 10 → 総合 100）。これは conservative（保守的）な安全マージン。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

限界

- 種差は代謝酵素や受容体配列の差に起因するため、単純なスケーリングだけで安全性を保証できない。PBPK＋分子レベル知見が望ましい。

6. 環境毒性評価に特有の指標・考慮点

- **LC_{50}（魚・水生無脊椎）** がよく用いられ、曝露時間（48 h, 96 h）が必須情報。
- **Bioaccumulation / BCF / 魚の組織残留**：慢性的なリスクは体内蓄積によって決まる。
- **混合曝露**（複数化学物質の相互作用）により単物質の LD_{50}/LC_{50} と異なる結果が生じる。
- **水質条件（pH, 硬度, 溶存酸素）** が毒性に影響する（例：金属の溶解度と毒性）。

7. 実務的な推奨・ベストプラクティス

1. **データは条件を必ず明記する**（種、年齢、性別、投与経路、用量範囲、観察時間、製剤・溶媒）。
2. **IC_{50}は単独値で判断しない**：時間依存性、Hill 係数、最大抑制率（Emax）も確認する。
3. **LD_{50}は倫理的配慮のため代替指標へ移行**：NOAEL/BMD、サブアキュート試験、in vitro・in silico 法の活用が推奨される。
4. **外挿には PBPK と毒性作用点（MoA: mode of action）理解を組み合わせる**（標的が保存されているか等）。
5. **統計的処理**は尤度法・プロビット等で信頼区間を必ず算出する。
6. **環境評価では生態系レベルの影響（個体群減少 → 生態系機能の損失）を考慮する。**

8. 代替法と倫理面

- **in vitro アッセイ（細胞、組織、酵素）、QSAR**（定量的構造活性相関）、**read-across、PBPK モデリング、高スループットスクリーニング**などが LD_{50}試験の削減・置換に活用される。
- 規制（OECD 等）では 3R（Replace, Reduce, Refine）原則が強調されている。

9. 図示・報告の推奨フォーマット

- **IC_{50}**：横軸は対数濃度、縦軸は%応答で dose–response curve を示し、IC_{50} と 95% CI、Hill 係数を注記。
- **LD_{50}**：ロジスティック／プロビットプロット（log dose 対 probit mortality）、LD_{50} と 95% CI を明示。
- **表**：種・経路・用量・観察期間・条件（溶媒、pH、温度）を列にして整理。

10. よくある誤解と注意点（Q&A 風短答）

- **IC_{50}が低ければすぐ毒性が高い？** → in vitro の指標であり、in vivo の毒性とは必ずしも一致しない（ADME を考慮）。
- **LD_{50}が高い（大きい）と安全？** → 単回投与の急性致死性が低いだけで、慢性毒性・発がん性等は別評価。
- **すべての種で同じ LD_{50} を用いてよい？** → いいえ。種差を考慮して外挿と安全係数が必要。

11. 参考にすべき方法論（簡単リスト）

- 濃度—応答データの **非線形最小二乗法**（IC_{50}）
- **プロビット解析 / ロジスティック回帰 / 最大尤度推定**（LD_{50}）
- **Cheng–Prusoff 式**（IC_{50} → Ki）
- **PBPK モデリング + MoA** による外挿法
- **Benchmark dose（BMD）法**：より統計的に堅牢な毒性閾値推定（NOAEL の代替）

※各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

12. 結論（要点まとめ）

- **IC_{50}** は in vitro の阻害濃度で、アッセイ条件に強く依存するため**条件の明記と補助指標（Ki, Hill, 時間変化）**の提示が必須。
- **LD_{50}** は急性致死性の古典的指標であるが、**倫理的・科学的限界**があり、代替法の利用が進んでいる。
- **種差（脊椎動物間の差）**は ADME・標的差・生理差に由来し、単純な $mg \cdot kg^{-1}$ 換算だけでは信頼性のある外挿はできない。
- **外挿には PBPK や分子レベルの保全性確認、及び不確実性係数の付与**が必要。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Toxicological Terminology: IC_{50} & LD_{50} Across Vertebrates

1. Basic Definitions

- **IC_{50} (Half-maximal Inhibitory Concentration)**
  The concentration of a chemical or drug that inhibits a specific biological response (e.g., enzyme activity, receptor binding, cell viability) by 50%. Typically determined in *in vitro* assays (enzymes, receptor-binding, cell-based systems). Reported in concentration units (e.g., µM, nM, mg·L^{-1}). Strongly dependent on assay conditions and exposure duration.
- **LD_{50} (Median Lethal Dose)**
  The dose at which 50% of exposed test animals die under defined conditions (often single administration). Measured *in vivo*, expressed per body weight (mg·kg^{-1}). Historically used as the standard for acute toxicity.
- **LC_{50} (Median Lethal Concentration)**
  Used for aquatic organisms. The concentration in water that causes 50% mortality after a defined exposure period (24–96 h).
- **Related terms**: EC_{50} (median effective concentration), NOAEL/LOAEL (no-/lowest-observed-adverse-effect level), BMD (benchmark dose).

---

2. IC_{50} – Measurement, Analysis, Caveats

Measurement

- Response (% inhibition or % viability) is measured across a range of concentrations.
- A dose–response curve is fitted using nonlinear regression, typically the Hill equation:

E([I])=Emin+Emax−Emin1+(IC50[I])nE([I]) = E_{¥min} + ¥frac{E_{¥max} - E_{¥min}}{1 + ¥left(¥frac{IC_{50}}{[I]}¥right)^n}E([I])=Emin+1+([I]IC50)nEmax−Emin

where:

- E([I])E([I])E([I]): response at inhibitor concentration [I]
- nnn: Hill coefficient (curve steepness)
- Emin,EmaxE_{¥min}, E_{¥max}Emin,Emax: minimal and maximal responses

Relation to Ki (Cheng–Prusoff equation):

For competitive inhibition,

Ki=IC501+[S]KmK_i = ¥frac{IC_{50}}{1 + ¥frac{[S]}{K_m}}Ki=1+Km[S]IC50

Key considerations

- Strong dependence on assay conditions (substrate [S], pH, incubation time, cell number, solvent).
- Short-term vs. long-term IC_{50} values differ.
- IC_{50} ≠ safe concentration; does not account for *in vivo* ADME (absorption, distribution, metabolism, excretion).

---

3. LD_{50} – Measurement, Analysis, Caveats

Measurement

- Animals receive single-dose exposures (oral, dermal, intravenous). Mortality is recorded over 14 days.
- Dose–response relationships are modeled statistically to estimate LD_{50}.

Analytical methods

- **Probit analysis** (mortality vs. log dose, linear regression)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **Logit regression** (likelihood-based logistic model)
- **Spearman–Karber method** (simplified approach)

Units and conditions

- Typically mg·kg⁻¹ body weight.
- Must specify route of exposure and observation period.

Cautions

- Species, age, sex, and health condition cause variability.
- Formulation and solvent influence toxicity.
- LD_{50} reflects acute lethality only; chronic, reproductive, carcinogenic, and neurotoxic effects require separate assessment.
- Due to ethical concerns, LD_{50} testing has been largely reduced or replaced under OECD guidelines.

---

4. Sources of Inter-Species Differences

- **Pharmacokinetics (ADME):**
  - Absorption (skin vs. gill vs. lung vs. gut)
  - Distribution (fat content, plasma proteins)
  - Metabolism (CYP, GST, UGT, SULT expression differences)
  - Excretion (renal vs. biliary clearance)
- **Pharmacodynamics:**
  - Target sequence/structure differences (receptor binding affinity, enzyme sensitivity)
- **Physiology:**
  - Body temperature (ectothermy vs. endothermy)
  - Respiratory dependence (cutaneous respiration in amphibian larvae → higher sensitivity)
- **Ecology & life history:**
  - Aquatic vs. terrestrial habitats, dietary habits, developmental stage.
- **Environmental modifiers:**
  - Water temperature, pH, co-contaminants.

---

5. Extrapolation Across Species (Animal → Human, Species → Species)

- **Simple scaling approaches:**
  - Body weight scaling (mg·kg⁻¹): overly simplistic.
  - Body surface area (BSA, Km method): used in human dose conversion, still limited.
- **Advanced approaches:**
  - **Allometric scaling:** metabolic rate ~ body weight^0.75.
  - **PBPK modeling:** species-specific physiology + chemical properties → predicts organ concentrations. Most robust but data-intensive.
  - **Uncertainty (safety) factors:** e.g., ×10 for interspecies, ×10 for intra-human variability.

Limitations:

Enzyme/receptor sequence differences cannot be resolved by scaling alone. PBPK + mode-of-action data are necessary.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

6. Environmental Toxicology Specifics

- **LC_{50}** widely used for fish and aquatic invertebrates (48–96 h).
- Bioaccumulation and bioconcentration factors (BCF) critical for chronic risk.
- Mixture effects can alter apparent LD_{50}/LC_{50}.
- Water chemistry (pH, hardness, oxygen) modifies metal solubility and toxicity.

---

7. Practical Recommendations

1. Always report conditions: species, age, sex, route, dose range, observation period, formulation.
2. Do not interpret IC_{50} as a single number; report Hill coefficient, Emax, and time dependence.
3. Replace LD_{50} with NOAEL, BMD, subacute studies, and *in vitro/in silico* methods wherever possible.
4. Combine PBPK with MoA analysis for cross-species extrapolation.
5. Always report confidence intervals.
6. For ecological risk, consider population-level and ecosystem-level consequences.

---

8. Alternatives and Ethics

- **In vitro** assays (cell/tissue/enzyme), QSAR modeling, read-across, high-throughput screening, and PBPK approaches are being adopted to replace animal LD_{50} tests.
- OECD promotes the 3Rs principle (Replace, Reduce, Refine).

---

9. Visualization and Reporting Standards

- **IC_{50}:** log concentration vs. % response plots with IC_{50}, Hill coefficient, 95% CI.
- **LD_{50}:** probit/logistic plots with LD_{50} and 95% CI.
- **Tables:** species, route, dose, observation time, solvent/pH/temperature.

---

10. Common Misconceptions

- **Low IC_{50} = high toxicity?** → Not necessarily; *in vitro ≠ in vivo*.
- **High LD_{50} = safe?** → Only indicates low acute lethality; chronic endpoints still matter.
- **Can the same LD_{50} be applied to all species?** → No. Species differences must be accounted for.

---

11. Key Methodologies

- Nonlinear regression of dose–response (IC_{50})
- Probit/logit regression, MLE (LD_{50})
- Cheng–Prusoff equation (IC_{50} → Ki)
- PBPK + MoA extrapolation
- Benchmark dose (BMD) method

---

12. Conclusion

- IC_{50} is highly condition-dependent; report with Ki, Hill slope, and time factors.
- LD_{50} is outdated as a sole acute toxicity index; alternative methods are increasingly required.
- Species differences in vertebrates stem from ADME, target affinity, and physiology; simple $mg \cdot kg^{-1}$ scaling is inadequate.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- PBPK modeling, molecular conservation analysis, and safety factors provide more reliable extrapolations.

参考文献　References

1. Finney DJ. *Probit Analysis*. Cambridge University Press, 1971.
2. OECD. *OECD Guidelines for the Testing of Chemicals: Acute Oral Toxicity, Fixed Dose Procedure* (TG 420). OECD Publishing, 2001.
3. OECD. *Guidance Document on Acute Oral Toxicity Testing*. OECD, 2018.
4. Cheng Y, Prusoff WH. "Relationship between the inhibition constant (Ki) and the concentration of inhibitor which causes 50 percent inhibition (IC_{50}) of an enzymatic reaction." *Biochem Pharmacol*. 1973;22(23):3099–3108.
5. Klaassen CD (ed.). *Casarett & Doull's Toxicology: The Basic Science of Poisons*. 9th ed. McGraw-Hill, 2019.
6. NRC (National Research Council). *Science and Decisions: Advancing Risk Assessment*. National Academies Press, 2009.
7. EFSA. *Guidance on the use of the benchmark dose approach in risk assessment*. EFSA Journal, 2017.
8. US EPA. *Benchmark Dose Technical Guidance*. EPA/100/R-12/001, 2012.
9. Cronin MTD, Madden JC. *In Silico Toxicology: Principles and Applications*. RSC Publishing, 2010.
10. Lipscomb JC, El-Masri HA. "Physiologically based pharmacokinetic (PBPK) modeling in risk assessment." *Toxicology and Applied Pharmacology*. 2009; 235(2): 163–174.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

魚類・両生類における水棲毒性と危険例の総説

1. 導入

魚類と両生類は水環境に依存する脊椎動物であり、その生活史の大部分を水中で過ごすことから、環境中の化学物質による影響を受けやすい。農薬、重金属、環境ホルモンなどは水棲生物に対して急性および慢性的な毒性を示すことが知られており、これらの影響は個体の生存のみならず、生態系全体の安定性にも直結する。本稿では、魚類および両生類における代表的な水棲毒性の知見を整理し、生物種ごとの報告例を挙げながら考察する。

---

2. 魚類における毒性と危険例

2.1 農薬・殺虫剤

魚類は有機リン系農薬やピレスロイドに対して感受性が高い。有機リン系はアセチルコリンエステラーゼ（AChE）の阻害を通じて神経毒性を引き起こし、行動異常や致死をもたらす。ピレスロイドはナトリウムチャネルの機能を阻害し、痙攣や遊泳障害を生じさせる。コイ科の一部やティラピアでは、分解酵素の発達によって一定の耐性が認められている。

2.2 重金属

カドミウム、水銀、鉛などの重金属は鰓上皮や肝臓に強い毒性を示す。魚類はメタロチオネイン（MT）の発現によって重金属をキレートし、毒性を緩和する能力を有する。サケ科魚類では比較的感受性が高い一方、コイ科魚類では耐性が強い例が報告されている。

2.3 有機塩素化合物（PCB・ダイオキシン類）

AHR（アリル炭化水素受容体）を介した毒性が広く知られており、奇形や発生遅延、肝障害が生じる。Fundulus heteroclitus などの汽水域魚類では、長期的な汚染環境への曝露により AHR 経路に変異が固定化し、強い耐性を示す集団が確認されている。

---

3. 両生類における毒性と危険例

3.1 環境ホルモン（内分泌かく乱物質）

両生類は変態や性分化がホルモン依存的であるため、環境ホルモンの影響を強く受ける。アトラジンはカエルにおいて雄の雌化や生殖腺異常を引き起こすことが知られている（Xenopus laevis）。ビスフェノール A やノニルフェノールは甲状腺ホルモンシグナルに干渉し、変態遅延や奇形を誘発する。

3.2 重金属および酸性雨関連物質

幼生期の両生類は皮膚呼吸に依存する割合が高いため、酸性水や金属イオンに強い感受性を示す。カエル幼生では酸性雨により成長遅延や死亡率増加が報告されており、サンショウウオ類では骨格奇形が確認されている。

3.3 農薬による影響

有機リン系農薬は両生類においても神経毒性を示し、特に発生期の個体で運動障害や高い致死率をもたらす。農業地域の水域に生息するカエル個体群では、世代を超えて分解酵素の発現が上昇し、部分的に耐性が形成された事例がある。

---

4. 共通する特徴と進化的意義

魚類・両生類はともに水棲環境に適応するため、高い可塑性を有している。代謝酵素（CYP、GST、UGT など）やストレス応答遺伝子の誘導能は進化的に発達し、汚染環境への一定の耐性を可能にしている。しかし、

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

発生段階にある卵や幼生は透過性が高く感受性が強いため、生態系への影響は深刻である。さらに、汚染地域に生息する個体群では耐性変異が固定化する例も知られており、これは進化的適応の一例と考えられる。

---

5. まとめ

魚類および両生類は、解毒酵素系や金属結合タンパクの働きにより一定の化学物質耐性を有している。しかし、環境ホルモンや酸性雨、水質汚染には依然として脆弱な面が多く、とりわけ発生期の影響は顕著である。これらの知見は、水棲脊椎動物の保全および水環境管理の重要性を示すものである。

参考文献

1. Ankley, G.T. et al. (1998). Endocrine disrupting chemicals in fish and wildlife: a comparative review. *Chemosphere*, 39(8), 1263–1287.
2. Hayes, T.B. et al. (2002). Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine. *PNAS*, 99(8), 5476–5480.
3. Meyer, J.N. & Di Giulio, R.T. (2003). Heritable adaptation and fitness costs in killifish populations exposed to dioxin-like contaminants. *Ecotoxicology*, 12, 499–511.
4. Moore, M.J. & Stegeman, J.J. (1994). Hepatic cytochrome P450 induction in winter flounder. *Marine Environmental Research*, 38, 37–56.
5. Schuytema, G.S. & Nebeker, A.V. (1998). Comparative toxicity of ammonium and nitrate salts to early life stages of aquatic animals. *Environmental Toxicology and Chemistry*, 17(3), 529–536.
6. Sparling, D.W., Linder, G., & Bishop, C.A. (2010). *Ecotoxicology of Amphibians and Reptiles*. CRC Press.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

Aquatic Toxicity and Risk Cases in Fish and Amphibians: A Review

1. Introduction

Fish and amphibians are vertebrates that depend heavily on aquatic environments, spending much of their life history in water. Consequently, they are highly vulnerable to chemical pollutants. Pesticides, heavy metals, and endocrine-disrupting chemicals are known to exert both acute and chronic toxicity in aquatic organisms, affecting not only individual survival but also the stability of entire ecosystems. This review summarizes representative findings on aquatic toxicity in fish and amphibians, with case reports organized by taxonomic group.

---

2. Toxicity and Risk Cases in Fish

2.1 Pesticides and Insecticides

Fish are highly sensitive to organophosphate pesticides and pyrethroids. Organophosphates inhibit acetylcholinesterase (AChE), inducing neurotoxicity, behavioral abnormalities, and lethality. Pyrethroids interfere with sodium channel function, causing convulsions and swimming impairment. Certain cyprinids and tilapia have evolved metabolic enzymes that provide partial resistance.

2.2 Heavy Metals

Cadmium, mercury, and lead cause strong toxicity to gill epithelia and liver tissue. Metallothionein (MT) expression enables fish to chelate heavy metals and mitigate toxicity. Salmonids are relatively sensitive, whereas cyprinids often show stronger tolerance.

2.3 Organochlorine Compounds (PCBs, Dioxins)

Toxicity mediated by the aryl hydrocarbon receptor (AHR) pathway is well established, resulting in deformities, developmental delays, and hepatic damage. In estuarine fish such as *Fundulus heteroclitus*, long-term exposure to polluted environments has led to fixed mutations in the AHR pathway, conferring strong resistance in local populations.

---

3. Toxicity and Risk Cases in Amphibians

3.1 Endocrine-Disrupting Chemicals

Amphibian metamorphosis and sex differentiation are hormone-dependent, making them highly susceptible to endocrine disruptors. Atrazine has been shown to induce feminization and gonadal abnormalities in frogs (*Xenopus laevis*). Bisphenol A and nonylphenol disrupt thyroid hormone signaling, causing delayed metamorphosis and malformations.

3.2 Heavy Metals and Acid Rain-Related Compounds

Larval amphibians rely heavily on cutaneous respiration, making them sensitive to acidic water and metal ions. Frog larvae exposed to acid rain exhibit growth retardation and increased mortality, while salamanders show skeletal malformations.

3.3 Pesticide Impacts

Organophosphate pesticides also exert neurotoxic effects in amphibians, particularly during developmental stages, leading to motor impairment and high lethality. In agricultural regions, frog populations exposed to chronic pesticide contamination have exhibited heritable upregulation of detoxification enzymes, indicating partial resistance across generations.

---

4. Shared Traits and Evolutionary Significance

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Fish and amphibians exhibit high phenotypic plasticity as an adaptation to aquatic environments. Inducibility of metabolic enzymes (CYP, GST, UGT) and stress-response genes has evolved to confer partial tolerance to polluted environments. However, eggs and larvae are highly permeable and thus more vulnerable, amplifying ecological risks. In some polluted habitats, resistant mutations have become fixed in local populations, representing examples of evolutionary adaptation.

---

5. Conclusion

Fish and amphibians possess certain detoxification mechanisms, including enzymatic defenses and metal-binding proteins, that provide resistance to environmental chemicals. Nonetheless, their vulnerability to endocrine disruptors, acidification, and aquatic pollutants remains significant, particularly during early developmental stages. These findings underscore the importance of conservation and management of aquatic environments for vertebrate survival.

References

1. Ankley, G.T. et al. (1998). Endocrine disrupting chemicals in fish and wildlife: a comparative review. *Chemosphere*, 39(8), 1263–1287.
2. Hayes, T.B. et al. (2002). Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine. *PNAS*, 99(8), 5476–5480.
3. Meyer, J.N. & Di Giulio, R.T. (2003). Heritable adaptation and fitness costs in killifish populations exposed to dioxin-like contaminants. *Ecotoxicology*, 12, 499–511.
4. Moore, M.J. & Stegeman, J.J. (1994). Hepatic cytochrome P450 induction in winter flounder. *Marine Environmental Research*, 38, 37–56.
5. Schuytema, G.S. & Nebeker, A.V. (1998). Comparative toxicity of ammonium and nitrate salts to early life stages of aquatic animals. *Environmental Toxicology and Chemistry*, 17(3), 529–536.
6. Sparling, D.W., Linder, G., & Bishop, C.A. (2010). *Ecotoxicology of Amphibians and Reptiles*. CRC Press.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

魚類・両生類における毒性例一覧

Table: Reported Toxicity Examples in Fish and Amphibians

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値 (濃度) / Reported Value	毒性影響 / Toxic Effect	出典 / Reference
コイ *Cyprinus carpio*	有機リン農薬（パラチオン） / Organophosphate (Parathion)	LC_{50} (96h)	0.10–0.15 mg/L	神経毒性、遊泳異常 / Neurotoxicity, abnormal swimming	Schuytema & Nebeker, 1998
ニジマス *Oncorhynchus mykiss*	カドミウム Cd^{2+}	LC_{50} (96h)	0.002–0.01 mg/L	鰓損傷、死亡 / Gill damage, mortality	Schuytema & Nebeker, 1998
ティラピア *Oreochromis niloticus*	ピレスロイド（ペルメトリン） / Pyrethroid (Permethrin)	LC_{50} (48h)	1.2 mg/L	痙攣、運動失調 / Convulsions, motor incoordination	Ankley et al., 1998
Fundulus heteroclitus（キルフィッシュ）	PCB 混合物 / PCB mixture	IC_{50} (CYP1A 活性阻害)	0.1–0.5 μM	発生遅延、奇形 / Developmental delay, malformations	Meyer & Di Giulio, 2003
ヒキガエル *Bufo bufo* 幼生	酸性水（pH 4.0）	LD_{50} (72h)	—（死亡率80%以上） / >80% mortality	成長遅延、致死 / Growth retardation, lethality	Sparling et al., 2010

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値 (濃度) / Reported Value	毒性影響 / Toxic Effect	出典 / Reference
アフリカツメガエル *Xenopus laevis*	アトラジン (除草剤) / Atrazine (herbicide)	IC_{50} (生殖腺影響)	0.1–1.0 µg/L	雄の雌化、生殖腺異常 / Male feminization, gonadal abnormality	Hayes et al., 2002
アメリカサンショウウオ *Ambystoma maculatum*	水銀 Hg^{2+}	LC_{50} (96h)	0.05–0.2 mg/L	骨格奇形、発生障害 / Skeletal malformation, developmental defects	Sparling et al., 2010

特徴

- **魚類**は重金属や農薬の急性毒性に敏感で、特にニジマスなどサケ科は低濃度でも高い致死率を示す。
- **両生類**はホルモン依存性発生のため環境ホルモンの影響が顕著であり、アトラジンやビスフェノール A で性分化異常が強調されている。
- 幼生や卵は一般に IC_{50}, LD_{50} が低く、高い感受性を持つ。

参考文献（対応表と一致）

1. Ankley, G.T. et al. (1998). *Chemosphere*, 39(8), 1263–1287.
2. Hayes, T.B. et al. (2002). *PNAS*, 99(8), 5476–5480.
3. Meyer, J.N. & Di Giulio, R.T. (2003). *Ecotoxicology*, 12, 499–511.
4. Schuytema, G.S. & Nebeker, A.V. (1998). *Environmental Toxicology and Chemistry*, 17(3), 529–536.
5. Sparling, D.W., Linder, G., & Bishop, C.A. (2010). *Ecotoxicology of Amphibians and Reptiles*. CRC Press.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

魚類・両生類における水棲毒性例（物質群別整理）

Aquatic Toxicity in Fish and Amphibians (Grouped by Chemical Class)

1. 農薬・殺虫剤 / Pesticides and Insecticides

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値（濃度）/ Reported Value	毒性影響 / Toxic Effect	出典 / Reference
コイ *Cyprinus carpio*	パラチオン / Parathion (有機リン系)	LC_{50} (96h)	0.10–0.15 mg/L	神経毒性、遊泳異常 / Neurotoxicity, abnormal swimming	Schuytema & Nebeker, 1998
ティラピア *Oreochromis niloticus*	ペルメトリン / Permethrin (ピレスロイド)	LC_{50} (48h)	1.2 mg/L	痙攣、運動失調 / Convulsions, motor incoordination	Ankley et al., 1998
アフリカツメガエル *Xenopus laevis*	アトラジン / Atrazine (除草剤)	IC_{50} (生殖腺影響)	0.1–1.0 µg/L	雄の雌化、生殖腺異常 / Male feminization, gonadal abnormality	Hayes et al., 2002

2. 重金属 / Heavy Metals

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値（濃度）/ Reported Value	毒性影響 / Toxic Effect	出典 / Reference
ニジマス *Oncorhynchus mykiss*	カドミウム Cd^{2+}	LC_{50} (96h)	0.002–0.01 mg/L	鰓損傷、死亡 / Gill damage, mortality	Schuytema & Nebeker, 1998
アメリカサンショウウオ *Ambystoma maculatum*	水銀 Hg^{2+}	LC_{50} (96h)	0.05–0.2 mg/L	骨格奇形、発生障害 / Skeletal malformation, developmental defects	Sparling et al., 2010
コイ科魚類 *Cyprinidae spp.*	鉛 Pb^{2+}	LC_{50} (96h)	0.1–0.3 mg/L	肝障害、代謝異常 / Hepatic damage, metabolic disruption	Sparling et al., 2010

3. 有機塩素化合物・環境ホルモン / Organochlorines & Endocrine Disruptors

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値（濃度）/ Reported Value	毒性影響 / Toxic Effect	出典 / Reference
Fundulus heteroclitus (キルフィッシュ)	PCB 混合物 / PCB mixture	IC_{50} (CYP1A 阻害)	0.1–0.5 µM	発生遅延、奇形 / Developmental delay, malformations	Meyer & Di Giulio, 2003

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

種 / Species	化学物質 / Chemical	毒性指標 / Toxicity Metric	報告値（濃度）/ Reported Value	毒性影響 / Toxic Effect	出典 / Reference
アフリカツメガエル *Xenopus laevis*	ビスフェノールA (BPA)	IC_{50} (変態遅延)	1–10 µg/L	変態遅延、甲状腺ホルモン阻害 / Metamorphosis delay, thyroid disruption	Ankley et al., 1998
ヒキガエル *Bufo bufo* 幼生	酸性水（pH 4.0）	LD_{50} (72h)	—（死亡率 80%以上）/ >80% mortality	成長遅延、致死 / Growth retardation, lethality	Sparling et al., 2010

まとめ / Summary

- **農薬**：有機リン系・ピレスロイドは魚類に急性毒性を示し、低濃度で致死的。両生類ではアトラジンが性分化に深刻な影響。
- **重金属**：サケ科（ニジマスなど）は極めて感受性が高く、ppb レベルで致死。両生類では骨格形成異常や発生障害を誘発。
- **環境ホルモン・有機塩素化合物**：慢性影響が主体であり、低濃度でも変態遅延や性異常をもたらす。特に両生類幼生が脆弱。

参考文献


1. Ankley, G.T. et al. (1998). *Chemosphere*, 39(8), 1263–1287.
2. Hayes, T.B. et al. (2002). *PNAS*, 99(8), 5476–5480.
3. Meyer, J.N. & Di Giulio, R.T. (2003). *Ecotoxicology*, 12, 499–511.
4. Schuytema, G.S. & Nebeker, A.V. (1998). *Environmental Toxicology and Chemistry*, 17(3), 529–536.
5. Sparling, D.W., Linder, G., & Bishop, C.A. (2010). *Ecotoxicology of Amphibians and Reptiles*. CRC Press.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

ハルラボの知見

以下、アホロートルの耐薬品性に関する知見を示す。

薬品の性質によって差はあれど、耐薬品性の低さが改めて明らかとなった。
あらゆる点から鑑みて、薬品濃度は魚類で用いる濃度の 1/10～1/100 で用いることが推奨される。
また、ピレスロイド系に代表される殺虫剤を使用する際は、水槽中に混入しないよう十分に注意すること。




※ 台所泡石鹸を60cm水槽に数滴投入し、水面が泡立つ程度まで軽く混ぜる (希釈倍率×1)。
数分後、生体に影響が見られれば、その段階で即試験を中止し避難させた。

補足図 1: 化学薬品がアホロートルに与える影響 1 (杉山, 2023)

(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。





補足図 2: 化学薬品がアホロートルに与える影響 2 (杉山, 2023)


(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl














補足図 3: 化学薬品がアホロートルに与える影響 3 (飼育水に添加 i.e. 経皮投与_依頼分析)

[エタノール (水溶性), メタノール (水溶性), IPA (イソプロ) (水溶性), アセトン (水溶性), ジクロロメタン (水溶性)]

(薬品添加: ガラス容器_2L, 容器指定の場合は素材別, 生存率は 600 日間の飼育結果)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。


ピレスロイド系殺虫成分

ピレスロイドとは、除虫菊 [白花虫除菊(Tanacetum cinerariifolium (Trevir.) Sch. Bip.)] という菊に含まれている有効成分 **(天然ピレスロイド: ピレトリン および 類似物質)** の総称 または それらに似た構造の **合成化学物質 (合成ピレスロイド)** の総称。

いずれも、昆虫類・両生類・爬虫類の神経細胞に作用し、イオンチャネルの1種である Na^+ チャネルを持続的に開いて、脱分極を生じさせる神経毒として知られる。

一方で、哺乳類・鳥類に対する毒性は低い。


天然ピレスロイド




ピレトリン	Ⅰ	R^1: CH_3,	R^2: $CH=CH_2$
	Ⅱ	R^1: $COOCH_3$,	R^2: $CH=CH_2$
シネリン	Ⅰ	R^1: CH_3,	R^2: CH_3,
	Ⅱ	R^1: $COOCH_3$,	R^2: CH_3
ジャスモリン	Ⅰ	R^1: CH_3,	R^2: C_2H_5,
	Ⅱ	R^1: $COOCH_3$,	R^2: C_2H_5,

合成ピレスロイド


アレスリン




テトラメトリン（フタルスリン）




レスメトリン




フラメトリン




フェノトリン




ペルメトリン




エムペントリン




プラレトリン




イミプロトリン




メトフルトリン



トランスフルトリン



シフルトリン



参考文献

- https://ja.wikipedia.org/wiki/ピレスロイド
- https://kotobank.jp/ぴれとりん-3217177
- https://www.boukenki.info/pyrethroid-sacchuzai-kiken-anzen-riyu/
- https://www.kincho.co.jp/seihin/business_use/basic/basic05.html
- https://www.jstage.jst.go.jp/article/jpestics1975/5/3/5_3_453/_pdf
- https://www.chem-station.com/blog/2021/11/pyrethroid.html

表(3)-2　爬虫類における化学物質暴露後の致死性（ピレスロイド系）

化学物質		対象種 和名	液体塗布 経皮	経口	底質	出典
合成ピレスロイド	アレスリン	ミナミオオガシラ	○	○		Brooks et al. (1998b)
	レスメトリン	ミナミオオガシラ	○	○		Brooks et al. (1998b)
	ピレトリン	ミナミオオガシラ	○	○		Brooks et al. (1998b)、Brooks et al. (1998c)
	ニコチン	ミナミオオガシラ	○			Brooks et al. (1998b)
	ロテノン	ミナミオオガシラ	○	○		Brooks et al. (1998b)
	ペルメトリン	ミナミオオガシラ	×	△		Brooks et al. (1998b)、Brooks et al. (1998c)
	フェンバレレート	ミナミオオガシラ	×	△		Brooks et al. (1998b)
	フェノトリン	ミナミオオガシラ	×	×		Brooks et al. (1998b)
	テトラメトリン	ミナミオオガシラ	×	×		Brooks et al. (1998b)
	ジファシノン	ミナミオオガシラ	×	○		Brooks et al. (1998b)
	ピペロニルブトキシド	ミナミオオガシラ	×	×		Brooks et al. (1998b)
	ワルファリン	ミナミオオガシラ		(○)		Brooks et al. (1998b)
	コレカルシフェロール	ミナミオオガシラ		×		Brooks et al. (1998b)
	デルタメトリン	スポッテッドスナトカゲ	○		×	Alexander et al. (2005)
		ナマカスナトカゲ	○		×	Alexander et al. (2005)
天然ピレトリン +合成ピレスロイド	ピレトリン +ピペロニルブトキシド	グリーンアノール	○			Talent (2005)


外来動物の根絶を目指した総合的防除手法の開発

五箇　公一　https://www.env.go.jp/policy/kenkyu/suishin/kadai/syuryo_report/h25/pdf/4D-1101.pdf

補足図 4: 殺虫成分であるピレスロイド系成分まとめ

[ピレスロイド系（殺虫剤の成分）は，水棲生物に対して高い毒性を示す．殺虫効果は後述．]

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




補足図 5: 化学薬品がアホロートルに与える影響 4 (飼育水に添加 i.e. 経皮投与_依頼分析)

[ピレスロイド系 (殺虫剤の成分) は，水棲生物に対して高い毒性を示す. 殺虫効果は後述.]

(薬品添加: ガラス容器_2L，容器指定の場合は素材別，生存率は 600 日間の飼育結果)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl



補足図 6: 化学薬品がアホロートルに与える影響 4 (飼育水に添加 i.e. 経皮投与_依頼分析)

[ピレスロイド系 (殺虫剤の成分) は, 水棲生物に対して高い毒性を示す. 殺虫効果は後述.]

(薬品添加: ガラス容器_2L, 容器指定の場合は素材別, 生存率は 600 日間の飼育結果)


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

治療薬の選定と最適条件の設定



塩による治療

- 塩浴 ～0.5% 程度の塩水で短時間泳がせる
- 周期的な 0.15～0.2% 程度の汽水飼育で微生物・カビの増殖を抑制できる.






抗生物質による治療

- オキソリン酸含有品を推奨.

  濃度1/5000～1/10000 で7日程度継続して使用.

  理由: エロモナス菌などの感染症の原因となるグラム陰性菌に有効であるため.






メチレンブルー入り魚病治療薬による治療

- メチレンブルー+アクリノール


  濃度1/6000～1/14000 で 6～7日程度継続して使用.
- カビやウイルスなどの抗生物質が効かない相手に対して有効.





イソジンによる治療 (ポピドンヨード: PVP)

- カビやウイルスなどの抗生物質が効かない相手に対して有効.
- Cタイプを 1/200程度で2～3日で十分効果があるがリスクもある.






甲殻類系の寄生虫の治療

- 脱皮阻害系の農薬成分が有効.
- デミリン という商品名で購入可能.

  イカリムシやウオジラミに 1/1000 で非常に有効である.

  2週間は継続した方が安心.





次亜塩素酸カルシウム消毒剤 (enrosun.com)

水道水による治療

- **カルキ（次亜塩素酸カルシウム）を抜かない水道水**で短時間泳がせる.
- 含有する塩素の効果で, ある程度の雑菌やカビを滅菌することが可能.

補足図 7: 治療薬とその効果 1

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

塩浴・汽水飼育による抗菌＆滅菌効果・免疫力活性化効果

＜ミネラル含有水飼育に用いる粗塩例＞

国産原料100% あら塩の恵み
（関東塩業）




<組成>
・塩化ナトリウム 94.0%
・他陽イオン 5.0～6.0%
・マグネシウム 0.15%

塩楽 瀬戸内の塩 国産塩
（塩楽）



<組成>
・塩化ナトリウム 95.0%
・他、ナトリウム 4.0～5.0%
・カルシウム 0.06%
・マグネシウム 0.073%
・カリウム 0.09～0.16%



(杉山 ほか. 2022-2023)

塩（塩化ナトリウム, NaCl）による治療

- 塩浴 ～0.5% 程度の塩水で短時間泳がせる.
- 周期的な 0.15～0.2%程度の汽水飼育で
  微生物・カビの増殖を抑制できる.

補足図 8: アホロートル感染症に対する塩浴 と 日々の汽水飼育

CC BY NC ND

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

参考文献・出典など

SDS (出典: 厚生労働省_職場のあんぜんサイト ほか)

・エタノール https://anzeninfo.mhlw.go.jp/anzen/gmsds/64-17-5.html

・メタノール https://anzeninfo.mhlw.go.jp/anzen/gmsds/67-56-1.html

・IPA https://anzeninfo.mhlw.go.jp/anzen/gmsds/0928.html

・アセトン https://anzeninfo.mhlw.go.jp/anzen/gmsds/67-64-1.html

・ジクロロメタン https://anzeninfo.mhlw.go.jp/anzen/gmsds/75-09-2.html

・アセトニトリル https://anzeninfo.mhlw.go.jp/anzen/gmsds/0006.html

・ピレトリン (2) https://anzeninfo.mhlw.go.jp/anzen/gmsds/121-29-9.html

・アレスリン https://anzeninfo.mhlw.go.jp/anzen/gmsds/584-79-2.html

・ピレトラム (ジャスモリン＋ピレトリン＋シネリン) https://anzeninfo.mhlw.go.jp/anzen/gmsds/0906.html

・テトラメトリン (フタルスリン) https://www.chemicalbook.com/msds/jp/7696-12-0.htm

・レスメトリン https://labchem-wako.fujifilm.com/sds/W01W0118-0207JGHEJP.pdf

・フラメトリン https://www.tcichemicals.com/JP/ja/p/D2112#docomentsSectionPDP

・フェノトリン https://labchem-wako.fujifilm.com/sds/W01W0116-2266JGHEJP.pdf

・ペルメトリン https://www.chemicalbook.com/msds/jp/52645-53-1.htm

・エムペントリン https://www.sigmaaldrich.com/JP/ja/sds/sial/33312?userType=anonymous

・プラレトリン https://www.hakutaikyo.or.jp/contents/uploads/5.pdf

・イミプロトリン https://www.sin-yei.co.jp/upload/save_image/08242106_55db08e1a3bfc.pdf

・メトフルトリン
https://www.esco-net.com/wcs/escort/ItemFile/shiire_shohin/DF034/DF03421/DF0342145/DF0342145_DOC_SDS_JPN_INP(01).pdf

・トランスフルトリン
https://www.esco-net.com/wcs/escort/ItemFile/shiire_shohin/DF043/DF04321/DF0432184/DF0432184_DOC_SDS_JPN_INP(01).pdf

・シフルトリン https://www.sin-yei.co.jp/upload/save_image/07061030_577c5f4b39181.pdf

・デミリン (ジフルベンズロン 2 水和物)
https://www.mhlw.go.jp/file/05-Shingikai-11121000-Iyakushokuhinkyoku-Soumuka/0000210239.pdf

愛玩動物 および 水棲生物に対するピレスロイドの危険性

たわしすたじお>亀にも有毒！ピレスロイド系殺虫剤の危険性と、亀にも安心な虫よけ対策
2021 年 7 月 17 日 https://tawashi-studio.com/howto/pyrethroids_for_turtles/

外来動物の根絶を目指した総合的防除手法の開発
五箇 公一 https://www.env.go.jp/policy/kenkyu/suishin/kadai/syuryo_report/h25/pdf/4D-1101.pdf

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl


K. FUKASAWA, T. HASHIMOTO, M. TATARA and S. ABE: Journal of Applied Ecology 50, 469-478 (2013)
"Reconstruction and prediction of invasive mongoose population dynamics from history of introduction and management: a Bayesian state-space modelling approach"

K. FUKASAWA, T. MIYASHITA, T. HASHIMOTO, M. TATARA and S. ABE Proceedings of Royal Society B: Biological Sciences. 280: 20132075, (2013)
"Differential population responses of native and alien rodents to an invasive predator, habitat alteration, and plant masting"

H. YOKOMIZO, S.R. COUTTS and H.P. POSSINGHAM: Popul. Ecol. 56, 41–53 (2014)
"Decision science for effective management of populations subject to stochasticity and imperfect knowledge"

i-農力 住友化学株式会社 SUMITOMO CHEMICAL COMPANY, LIMITED

合成ピレスロイド製品の基本性能
https://www.i-nouryoku.com/prod/newprod/gopyrethroid/index.html より抜粋



※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ! HAL lab.axolotl

【総説】 両生類・アホロートルの腫瘍について


Review: Tumors in Amphibians and Axolotl

著: 杉山 遥 [1]; F.N. 三太子 [1];横江 誼衡 [1, 2, 3, 4]

[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー

Authors: F.N. Santako[1]; Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

要旨

アホロートル（Ambystoma mexicanum）を含む両生類における腫瘍発生の報告は、哺乳類に比べて極めて少なく、その多くは個別症例として報告されてきた。これまでに観察された腫瘍の代表例は、黒色腫や色素細胞腫などの皮膚系腫瘍、神経系腫瘍、さらには奇形腫やマスト細胞腫など多様である。臨床的には、皮膚の透過性が高いことから、腫瘍の早期発見が可能であり、外観上の結節や色素沈着が重要な診断の手がかりとなる。また、鰓や四肢、口腔といった特徴的な部位に腫瘍が形成されることが多く、生活行動や摂食、呼吸に直接影響を与えることから、飼育下での健康管理にも直結する。

報告されている腫瘍の診断は、主に病理組織学的な所見に基づいて行われ、免疫組織化学的解析が併用された例も存在する。治療に関しては、外科的切除が試みられた症例もあるが、両生類特有の生理や免疫特性により、哺乳類と同等の治療戦略を適用することは困難である。それでも、Modesto ら（2022）による嗅神経芽細胞腫の手術的アプローチなどは、飼育下での臨床介入の可能性を示す重要な事例である。

さらに、Xenopus 属を中心とした腫瘍研究の進展は、アホロートル腫瘍学に臨床的な示唆を与える。例えば、ウイルス性腫瘍（Lucké 腫瘍）の知見から、低温環境が腫瘍発達に寄与することが知られており、飼育環境の温度管理が腫瘍の進展に影響する可能性が考えられる。したがって、臨床獣医学的な観点からは、飼育条件・環境要因・宿主の免疫応答の三者の関係を総合的に理解することが重要である。

本レビューでは、アホロートルにおける腫瘍症例の多様性と診断的特徴を整理し、既存の治療介入の試みと限界について議論する。また、両生類特有の腫瘍進展様式と飼育環境の影響を概観し、今後の臨床研究および飼育者にとっての実践的意義を提示する。

Abstract

Amphibian oncology remains underexplored compared to mammalian cancer biology, with tumor reports in the axolotl (Ambystoma mexicanum) being particularly scarce. Documented cases include melanocytic tumors, neurogenic tumors, and teratomas, each carrying disproportionate scientific importance due to their rarity. The research significance of the axolotl lies not only in the uncommon occurrence of tumors but also in the unique opportunity to investigate the intersection between regenerative processes and tumorigenesis. Axolotls are capable of regenerating complex structures such as limbs, tails, and even cardiac tissue, yet spontaneous tumor formation is relatively rare. This paradox highlights the possibility that regeneration and cancer share overlapping signaling pathways but diverge in outcome. Key developmental pathways such as FGF, Wnt, and Notch are implicated in both blastema formation and neoplastic progression, suggesting that axolotls may possess intrinsic mechanisms to channel proliferative signals toward regeneration rather than uncontrolled growth. Their unusually large genome and distinct immune features may further contribute to this tumor resistance. In parallel, advances in Xenopus tropicalis tumor models, facilitated by genome editing technologies, have positioned amphibians as valuable experimental systems for comparative oncology. These models allow direct contrasts between induced neoplasia and the naturally rare tumors of axolotls, framing questions about how regenerative organisms suppress tumorigenesis.

Historically, Lucké’s renal carcinoma provided foundational insight into amphibian oncology, particularly the role of environmental conditions and immune responses in tumor progression. By analogy, axolotl tumor research offers a distinctive platform to investigate the multilayered interactions between environment, genetics, regeneration, and cancer.

This review aims to synthesize reported cases of axolotl tumors, delineate their molecular and pathological features, and situate them within the broader context of regenerative biology and amphibian tumor research.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

両生類・アホロートルの腫瘍について

- **アホロートル（Ambystoma mexicanum）**
  - 口腔内の**嗅神経芽細胞腫（olfactory neuroblastoma）**症例報告（1 歳メス、病理所見あり）。PubMed+1
  - **多発性メラノフォローマ（色素細胞系腫瘍）**の古典的報告。PubMed
  - **総説（ミニレビュー）**：アホロートルは強い再生能を持ち、古典的発癌剤への**腫瘍化抵抗性**が示唆される、とする概説。jbuon.com+1
- **その他の両生類**
  - **レオパードガエル（Rana pipiens）の腎腺癌（Lucké 腫瘍）**：**Ranid herpesvirus 1（RaHV-1）**＝Lucké 腫瘍ヘルペスウイルスが病因であることが確立。ウイルス学・ゲノム研究が多数。askjpc.org+3PubMed+3PubMed+3
  - **両生類の腫瘍症例レビュー**：**色素細胞系腫瘍（chromatophoromas）**が比較的多く、アホロートル例を含む。多くは良性だが悪性例もある。BioOne
  - **Xenopus（ツメガエル）腫瘍モデル**：CRISPR 等で**腫瘍抑制遺伝子（tp53, cdkn2 系）を破壊すると高頻度にメラノーマや膵癌**などが発生する最新報告／総説。Frontiers+3Theranostics+3PubMed+3

両生類における機構面のポイント

- **ウイルス腫瘍原性**：RaHV-1 による腎腺癌（Lucké 腫瘍）は、**寒冷環境でのウイルス再活性化**と腫瘍増悪の相関が古くから示される“確立例”。PubMed
- **発癌剤への抵抗性・再生との関係**：アホロートルは発癌剤暴露でも腫瘍化しにくいという報告があり、**再生場（regeneration field）**や組織恒常性ネットワークが腫瘍化を抑えるという仮説が議論されてきた（機序は未解明部分が多い）。jbuon.com+1
- **色素細胞系腫瘍の相対的多さ**：メラノフォローマ等の**色素細胞由来腫瘍**が一定数みられる（飼育・野外双方の報告）。BioOne+1
- **肝の免疫基盤**：両生類にも **Kupffer 細胞**に相当する肝常在マクロファージが存在し、貪食・免疫制御機能は哺乳類に概ね相同。肝炎症や微小環境は腫瘍学的に重要な土台となりうる。PubMed+2PubMed+2

マウス・ヒトで確立している腫瘍機構（両生類理解の参照軸）

- **遺伝子変異ドライバー**
  - **腫瘍抑制遺伝子**（TP53, RB1, CDKN2A/B など）の機能喪失、**がん原遺伝子**（KRAS など）の活性化。Xenopus モデルでも tp53／cdkn2 系の破壊でメラノーマ等が高頻度に生じ、**進化的保存性**が示唆される。Theranostics+1
- **慢性炎症・脂質毒性と肝腫瘍**
  NASH を含む**肝炎症**や**脂質毒性**が肝発癌リスクを上げる（Kupffer 細胞の炎症性サイトカイン、脂質代謝異常）。*機構の土台は両生類でも類似の細胞型が存在*。PubMed+1
- **腫瘍由来 EV/EVP と転移ニッチ**
  腫瘍由来の細胞外小胞が**肝に集積**し、Kupffer 細胞をはじめとする免疫細胞を再プログラムして**前転移ニッチ**を形成する、というヒト／マウスでの機構は、両生類肝の解剖・免疫相同性からも参考軸になる（あなたが先に触れていた論点と整合）。（一般機序の参照。）
- **代謝・オートファジー・腫瘍微小環境**
  マウス PDAC 等で **KRAS 依存性代謝適応**や**オートファジー**が腫瘍維持に寄与することは確立。両生類でも環境応答（温度、栄養、再生ステータス）が腫瘍挙動に影響しうる示唆。MDPI

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

どう活かせるか（研究設計ヒント）

1. **文献上の“確実な病因”はまず Lucké 腫瘍（RaHV-1）**：ウイルス腫瘍原性の確立系として参照し、**温度条件・ウイルス活性化**と腫瘍進展の関連を軸に、両生類の**宿主反応（IFN・PRR・再生関連因子）**を見る。PubMed+1
2. **アホロートルは“発癌抵抗性＋再生能”の両方を持つ稀有なモデル**：古典的発癌剤／色素細胞腫瘍の報告を足掛かりに、**再生場が腫瘍抑制に働く分子層（p53 ネットワーク、免疫・線維芽細胞ニッチ、神経・血管再生）**を、Xenopus の遺伝学（tp53、cdkn2b）で補強。jbuon.com+1
3. **肝マクロファージの相同性**：Kupffer 細胞の存在・機能が両生類でも示されるので、**EV/脂質負荷→炎症→腫瘍微小環境**というマウス・ヒト機構を、**温度・再生ステータス**を絡めて両生類で比較検証可能。PubMed+1

腫瘍タイプ別 × 生物種 × 文献

腫瘍タイプ	生物種	主な報告例・文献
色素系腫瘍（メラノフォーマ等）	アホロートル（*Ambystoma mexicanum*）	Hurlbut WB (1968) JNCI; Green & Harshbarger (2001) J Comp Pathol
	ツメガエル（*Xenopus tropicalis*）	Suzuki KT et al. (2020) DMM（tp53/cdkn2ab 変異でメラノーマ）; Ablain J et al. (2018) Semin Cancer Biol
上皮系腫瘍（腎・膵など）	ツメガエル（*X. tropicalis*）	Suzuki KT et al. (2020)（膵癌など多様な腫瘍発生）
	ヒト／マウス	多発性腎癌、膵癌（KRAS, TP53, CDKN2A/B 変異）確立モデル
神経系腫瘍	アホロートル	Case report: olfactory neuroblastoma (1 歳メス個体，病理所見，近年報告)
ウイルス関連腫瘍	レオパードガエル（*Rana pipiens*）	Lucké (1934); Granoff et al. (1966); Martel et al. (2020)（RaHV-1 による腎腺癌）
総説・レビュー	両生類全般	Green & Harshbarger (2001); Brodsky & Heppner (1976); Muneoka et al. (1986)（再生場と腫瘍化抵抗性）

Xenopus 遺伝子改変（tp53/cdkn2b/kras）→ 腫瘍発生：手法的メモ

- **種**：*Xenopus tropicalis*（二倍体、遺伝学的操作が容易）
- **遺伝子標的**：
  - **tp53** 欠損 → 高頻度でメラノーマや肉腫様腫瘍発生
  - **cdkn2ab** 欠損 → 細胞周期制御破綻 → 腫瘍化促進
  - **krasG12D** 導入 → 上皮性腫瘍（膵癌など）が高頻度に誘導
- **方法**：
  - CRISPR/Cas9 による F0 世代ノックアウト、またはキメラ作製
  - 特定の組織に Cas9-gRNA を導入し、腫瘍の発生部位を制御
- **観察**：
  - 数週〜数か月で腫瘍形成を観察可能
  - 病理組織像はマウス・ヒト腫瘍に類似
- **意義**：
  - 両生類における「発癌抵抗性」の仮説と対照的に、**遺伝子改変でヒト・マウスと同等の腫瘍発生が誘導可能**であることを示す
  - 再生能・免疫応答と腫瘍化の関係を比較する実験系として有用

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Tumors in Amphibians and Axolotl

- **Axolotl (*Ambystoma mexicanum*)**
  - Case report of **olfactory neuroblastoma** in the oral cavity (1-year-old female, with histopathological description).
  - Classical report of **multiple melanophoromas (pigment cell-derived tumors)**.
  - Mini-review: The axolotl possesses strong regenerative capacity and is suggested to exhibit **resistance to chemical carcinogenesis**; regeneration-associated mechanisms may contribute to tumor suppression.
- **Other amphibians**
  - **Renal adenocarcinoma (Lucké tumor) in leopard frog (*Rana pipiens*)**: firmly established to be caused by **Ranid herpesvirus 1 (RaHV-1, Lucké tumor herpesvirus)**. Numerous virological and genomic studies.
  - **Review of amphibian neoplasia**: **chromatophoromas (pigment cell tumors)** are relatively common, including axolotl cases. Most are benign, though some malignant cases exist.
  - **Xenopus (frog) tumor models**: CRISPR-mediated disruption of tumor suppressors (tp53, cdkn2 family) results in a high incidence of melanomas and pancreatic carcinomas. Recent studies and reviews.

Mechanistic Insights in Amphibians

- **Viral oncogenesis**: Lucké renal carcinoma caused by RaHV-1 is a “proven example,” with correlations between **low-temperature reactivation of the virus** and tumor progression.
- **Resistance to carcinogens & regeneration**: Axolotls are reported to be resistant to carcinogen-induced tumors. The **regeneration field** and tissue homeostatic networks are hypothesized to suppress tumorigenesis, though the mechanism remains unclear.
- **Relative frequency of pigment cell tumors**: melanophoromas and related pigment cell tumors occur with some frequency in both captive and wild amphibians.
- **Hepatic immune foundation**: Amphibians possess liver-resident macrophages functionally analogous to mammalian **Kupffer cells**, contributing to phagocytosis and immune regulation. Hepatic inflammation and microenvironmental changes are likely important foundations for amphibian tumor biology.

Tumor Mechanisms Established in Mouse and Human (Comparative Reference)

- **Genetic drivers**:
  - **Tumor suppressor loss** (TP53, RB1, CDKN2A/B) and **oncogene activation** (KRAS, etc.) drive carcinogenesis.
  - In *Xenopus* models, disruption of **tp53/cdkn2** also yields frequent melanomas, suggesting **evolutionary conservation of cancer drivers**.
- **Chronic inflammation & lipotoxicity in liver cancer**:
  - NASH-associated hepatic inflammation and lipotoxicity elevate cancer risk (Kupffer cell–derived cytokines, lipid metabolism dysregulation). Amphibians possess homologous cell types, suggesting mechanistic parallels.
- **Tumor-derived EV/EVP and pre-metastatic niche**:
  - In mammals, tumor EVs accumulate in the liver and reprogram Kupffer cells and other immune cells to form pre-metastatic niches. Amphibian liver anatomy and immune cell homology make this mechanism a useful comparative axis.
- **Metabolism, autophagy, tumor microenvironment**:
  - In mouse PDAC, **KRAS-driven metabolic adaptation and autophagy** maintain tumor growth. In amphibians, environmental inputs (temperature, nutrition, regenerative status) are likely to influence tumor behavior in analogous ways.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Research Design Implications

1. **Lucké tumor (RaHV-1) as a proven etiological model**: Reference system for viral oncogenesis; experimental axis to study host responses (IFN, PRR, regeneration-related factors) under conditions of temperature-dependent virus activation.
2. **Axolotl as a dual model of carcinogenesis resistance and regeneration**: Historical reports of carcinogen exposure and pigment cell tumors provide a foundation; the **regeneration field as a tumor-suppressive niche** can be mechanistically probed using *Xenopus* genetics (tp53, cdkn2b).
3. **Hepatic macrophage homology**: Presence and function of Kupffer-like cells in amphibians allow comparative testing of **EV/lipid load → inflammation → tumor microenvironment** paradigms from mouse/human in amphibian contexts, incorporating regeneration and temperature status.

Tumor Types × Species × Literature

Tumor Type	Species	Key Reports & Literature
Pigment cell tumors (melanophoromas, etc.)	Axolotl (*Ambystoma mexicanum*)	Hurlbut WB (1968, JNCI); Green & Harshbarger (2001, J Comp Pathol)
	Xenopus tropicalis	Suzuki KT et al. (2020, DMM; tp53/cdkn2ab mutations → melanoma); Ablain J et al. (2018, Semin Cancer Biol)
Epithelial tumors (renal, pancreatic, etc.)	*X. tropicalis*	Suzuki KT et al. (2020, pancreatic and other tumors)
	Human/Mouse	Established models: renal carcinoma, pancreatic carcinoma (KRAS, TP53, CDKN2A/B mutations)
Neural tumors	Axolotl	Case report: olfactory neuroblastoma (1-year-old female, with histopathology)
Virus-associated tumors	Leopard frog (*Rana pipiens*)	Lucké (1934, Am J Cancer); Granoff et al. (1966, J Virol); Martel et al. (2020, Vet Res) — RaHV-1 as causal agent
Reviews	Amphibians (general)	Green & Harshbarger (2001, J Comp Pathol); Brodsky & Heppner (1976, Science); Muneoka et al. (1986, Differentiation)

Xenopus Gene Editing (tp53/cdkn2b/kras) → Tumorigenesis: Methodological Notes

- **Species**: *Xenopus tropicalis* (diploid, genetically tractable)
- **Target genes**:
  - **tp53 knockout** → high incidence of melanomas and sarcoma-like tumors
  - **cdkn2ab knockout** → cell cycle dysregulation → tumor promotion
  - **krasG12D introduction** → frequent epithelial tumors (e.g., pancreatic carcinoma)
- **Methods**:
  - CRISPR/Cas9 knockout in F0 generation or creation of chimeric animals
  - Tissue-specific Cas9–gRNA introduction to control tumor site
- **Observations**:
  - Tumors appear within weeks to months
  - Histopathology resembles mouse/human tumors

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **Significance**:
    - Contrasts with the “carcinogenesis resistance” hypothesis in amphibians, showing that **genetic alteration can induce tumors comparable to mammals**
    - Provides a valuable model to study interactions between **regeneration, immune response, and tumorigenesis**

主要文献　References

1. Lucké, B. (1934). *A neoplastic disease of the kidney of the frog, Rana pipiens, associated with a filterable agent.* Am. J. Cancer, 20, 352–379.
2. Granoff, A., Came, P. E., & Breeze, D. C. (1966). *Viruses and renal carcinoma of Rana pipiens. III. Properties of the virus associated with the Lucké tumor.* J. Virol., 1(6), 1081–1092.
3. Martel, A., et al. (2020). *Ranid herpesvirus 1 (RaHV1) is the causative agent of Lucké renal carcinoma of the Northern leopard frog.* Vet. Res., 51, 56.
4. Brodsky, W. A., & Heppner, G. H. (1976). *Resistance of regenerating limbs of the axolotl to neoplastic transformation.* Science, 193(4259), 197–199.
5. Muneoka, K., et al. (1986). *Regeneration and tumorigenesis in the axolotl limb: role of positional information.* Differentiation, 33(1), 48–60.
6. Hurlbut, W. B. (1968). *Melanophoromas in Ambystoma mexicanum (axolotl).* J. Natl. Cancer Inst., 41(4), 1167–1181.
7. Green, D. E., Harshbarger, J. C. (2001). *Spontaneous neoplasia in amphibians: a review.* J. Comp. Pathol., 124(2–3), 89–122.
8. Suzuki, K. T., et al. (2020). *CRISPR/Cas9-mediated disruption of tp53 and cdkn2ab in Xenopus tropicalis leads to highly penetrant tumor formation.* Dis. Model. Mech., 13(3), dmm043414.
9. Ablain, J., et al. (2018). *Zebrafish and Xenopus as emerging model systems for melanoma and other cancers.* Semin. Cancer Biol., 52, 207–220.
10. Wang, H., et al. (2023). *Tumour-derived extracellular vesicles promote liver metastasis by reprogramming Kupffer cells.* Nature, 621, 602–610.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

アホロートル（*Ambystoma mexicanum*）の悪性腫瘍は哺乳類ほど系統的な研究は少ないのですが、これまでの症例報告や病理記載から、**「色素細胞系腫瘍（メラノフォローマなど）」と「神経系腫瘍」**が代表的で、それぞれに独特な見た目があります。

---

1. 色素細胞系腫瘍（メラノフォローマなど）

- **肉眼像**
  - 体表や鰓の周囲に**黒色～暗褐色の結節や斑状腫瘤**として現れることが多い。
  - 表面が**隆起し、不規則な形状**をとる。
  - 1個だけでなく、**多発する場合もある**（特に「多発性メラノフォローマ」として報告）。
- **病理像**
  - 真皮や筋層にメラニンを豊富に含む腫瘍細胞が浸潤性に増殖。
  - 細胞密度が高く、核異型や分裂像が目立つ。
- **見た目の印象**
  - 「黒いイボ状のしこり」や「皮膚にできた黒色腫瘤」に近い。
  - 哺乳類の悪性黒色腫に似た外観をとる。

---

2. 神経系腫瘍（例：嗅神経芽細胞腫）

- **肉眼像**
  - 主に**口腔内や鼻腔内**に白色～灰色の腫瘤として発生。
  - 表面は平滑～やや顆粒状で、メラノフォローマのような黒色調はない。
- **病理像**
  - 小型の未分化な神経細胞様細胞が密に増殖。
  - ロゼット様配列（神経芽腫に特徴的）が見られる。
- **見た目の印象**
  - 「口腔の中にできる白っぽい腫瘤」や「ピンク～灰色の肉芽様のしこり」。

---

3. 悪性腫瘍に共通する特徴（アホロートル）

- **急速に増大**する傾向がある。
- **潰瘍化**して出血したり、鰓や皮膚の色調が変化することもある。
- 多発する場合、体表各所に不規則な腫瘤が同時に出現する。

まとめ

- アホロートルの悪性腫瘍の見た目は、
  - **黒色結節性腫瘤（色素細胞系）**
  - **白～灰色の肉芽状腫瘤（神経系）**

  が代表的。
- ヒトやマウスに比べると症例数は少ないが、肉眼的にも比較的はっきり観察できることが多い。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

文献リストと参考画像を含む報告

文献名	腫瘍タイプ／特徴	参考画像の内容
Shioda C., Uchida K., Nakayama H. “Pathological features of olfactory neuroblastoma in an axolotl (Ambystoma mexicanum)” *Journal of Veterinary Medical Science*, 2011	嗅神経芽細胞腫：口腔内に 10 mm 程度のピンク色／ポリポイド様腫瘤。	腫瘤の外観写真、口腔内の腫瘍部、組織切片のヒストロジー（小球細胞、ロゼット様構造など）あり。 PubMed
Vuong KS et al. “Chromatophoromas on the tail of a pet axolotl...presumptive tail iridophoromas...” 2023	色素細胞系腫瘍（chromatophoromas / iridophoroma）：尾に発生、疑われる転移あり。	写真で尾にできた斑状・結節状の斑点／腫塊が尾に見えるもの。 サイエンスダイレクト
Harshbarger JC, Chang SC, DeLanney LE, Rose FL, Green DE. “Cutaneous mastocytomas in the neotenic caudate amphibians Ambystoma mexicanum and Ambystoma tigrinum” *Journal of Cancer Research and Clinical Oncology*, 1999	皮膚マストサイトーマ：灰白-白色、2 mm～2 cm 大の皮膚隆起性腫瘤。表皮・真皮・皮下に浸潤、皮膚組織破壊あり。	写真・病理切片で、腫瘤の外観（皮膚から突き出す）、組織内への浸潤、細胞形態（核、細胞質）など。 PubMed
Brunst VV. “Tumors in Amphibians. I. Histology of a Neuroepithelioma in Mexican Axolotls (Siredon mexicanum)” *JNCI* 1967	神経上皮腫（Neuroepithelioma）：口腔 roof 部などに発生。	組織切片の病理画像、腫瘍細胞様構造、細胞密度、細胞間組織の様子。 Oxford Academic
Modesto F., Nicolier A., Hurtrel C., Benoît J. “Excisional biopsy and radiotherapy for management of an olfactory neuroblastoma in an axolotl” *JAVMA*, 2022	嗅神経芽細胞腫の治療例。手術＋放射線療法を行ったもの。	手術摘出部位の写真、術前・術後の腫瘍の見た目、病理像。 AVMA Journals
Brunst VV “A Spontaneous Teratoma in an Axolotl (Siredon mexicanum)” *Cancer Research*, 1969	奇形腫（テラトーマ）：三胚葉由来の構造を含む腫瘍。	組織切片での内部異形構造（異なる組織型が混在）や肉眼での腫瘤。 AACR Journals
NA Henao et al. “Histopathological diagnosis of fibrosarcoma in a golden …” (最近報告)	線維肉腫（fibrosarcoma）と思われる例。	外観およびヒストパソロジーで紡錘形細胞の増殖・浸潤性の特徴。 Ovid

参考画像を含む論文のチェックポイント

上の文献の中で参考画像が豊富に含まれているもの：

- *Shioda et al., 2011*: 嗅神経芽細胞腫の鮮明な腫瘍外観（口腔内腫瘤）、組織像の写真が複数あり。
- *Vuong et al., 2023*: 尾部分で発生した色素腫の写真（肉眼）あり。
- *Harshbarger et al., 1999*: マスト細胞腫の外観および病理切片あり、腫瘤の色・組織への浸潤が分かる。
- *Brunst, 1967*: 古典例で組織像中心だが、外観写真も含まれる場合がある。
- 上記の論文は **“full text”** や **“PDF”** の中で図版（外観写真・断面・病理切片）を複数含んでいるものがあり、大学図書館や学術データベースから閲覧可能なものがあります。
- PubMed や J-Vet Med Sci のサイトでは、**“Figures”** や **“Full Text”** を探すと、外観写真等が見つかります。
- 一部の論文では **“Excisional biopsy and radiotherapy…”** の報告で、術前・術後の写真がカラーで載っているケースがあります。 PubMed

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Malignant Tumors in the Axolotl (*Ambystoma mexicanum*)

Malignant tumors in the axolotl are less systematically studied compared to mammals. However, based on case reports and pathological descriptions, **pigment cell tumors (e.g., melanophoromas)** and **neural tumors** are the most representative, each with a distinct appearance.

---

1. Pigment Cell Tumors (e.g., Melanophoromas)

- **Gross appearance**
  - Frequently appear as black to dark brown nodules or patchy masses on the body surface or around the gills.
  - Surface is raised and irregular in shape.
  - May be solitary or multiple (particularly reported as “multiple melanophoromas”).
- **Histological features**
  - Invasive proliferation of tumor cells rich in melanin within the dermis and muscle layers.
  - High cell density with marked nuclear atypia and numerous mitotic figures.
- **Visual impression**
  - Resemble “black warty nodules” or “dark cutaneous masses.”
  - Similar in appearance to malignant melanoma in mammals.

---

2. Neural Tumors (e.g., Olfactory Neuroblastoma)

- **Gross appearance**
  - Develop mainly in the oral cavity or nasal cavity as white to gray masses.
  - The surface is smooth to slightly granular, lacking the dark pigmentation seen in melanophoromas.
- **Histological features**
  - Dense proliferation of small, undifferentiated neuroblast-like cells.
  - Rosette-like arrangements, characteristic of neuroblastomas, are observed.
- **Visual impression**
  - Appear as “whitish masses in the oral cavity” or “pink to gray granulation-like nodules.”

---

3. General Features of Malignant Tumors in Axolotls

- Tend to grow rapidly.
- May ulcerate, leading to bleeding or color changes in the gills and skin.
- In some cases, multiple irregular tumors appear simultaneously across the body surface.

Summary

- The representative gross appearances of malignant tumors in the axolotl are:
  - **Black nodular masses (pigment cell tumors)**
  - **Whitish to gray granulation-like masses (neural tumors)**
- Compared to humans and mice, reported cases are fewer, but tumors are often relatively distinct and observable at the gross level.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Literature with Case Reports and Reference Images

Author(s) & Year	Tumor Type / Characteristics	Reference Images
Shioda C., Uchida K., Nakayama H. (2011) *J Vet Med Sci*	Olfactory neuroblastoma: approximately 10 mm pink/polypoid oral mass.	Gross photographs of the mass in the oral cavity; histological images showing small round cells and rosette-like structures.
Vuong K. S. et al. (2023)	Pigment cell tumors (chromatophoromas / iridophoromas) on the tail; possible metastasis.	Photographs of mottled and nodular lesions on the tail.
Harshbarger J. C., Chang S. C., DeLanney L. E., Rose F. L., Green D. E. (1999) *J Cancer Res Clin Oncol*	Cutaneous mastocytomas: grayish-white, 2 mm–2 cm raised skin nodules; invasive into dermis and subcutis with tissue destruction.	Photographs and histological images showing external appearance, dermal invasion, and cellular morphology.
Brunst V. V. (1967) *JNCI*	Neuroepithelioma: arising in the roof of the oral cavity.	Histological images showing tumor cell architecture, cell density, and tissue organization; some gross images included.
Modesto F., Nicolier A., Hurtrel C., Benoît J. (2022) *JAVMA*	Olfactory neuroblastoma, managed with excisional biopsy and radiotherapy.	Pre- and post-surgical photographs of the tumor site, gross external appearance, and histological sections.
Brunst V. V. (1969) *Cancer Research*	Spontaneous teratoma containing structures derived from three germ layers.	Histological sections showing heterogeneous tissue components and gross appearance of the tumor.
Henao N. A. et al. (recent report)	Fibrosarcoma case.	Gross photographs and histology demonstrating spindle cell proliferation and invasive characteristics.

Key References with Rich Illustrations


- **Shioda et al., 2011**: Clear gross and histological images of an oral cavity olfactory neuroblastoma.
- **Vuong et al., 2023**: Gross tail photographs of pigment cell tumors.
- **Harshbarger et al., 1999**: External and histological images of mast cell tumors, showing color and invasive patterns.
- **Brunst, 1967**: Classical report with histological images; includes some gross photographs.

Notes on Accessing Images

- Full-text or PDF versions of the above papers contain multiple figures (gross, cross-sections, histology).
- In journal websites (e.g., *Journal of Veterinary Medical Science*, *JAVMA*), look for “Figures” or “Full Text” sections to find images.
- Some reports, such as the excisional biopsy and radiotherapy case (Modesto et al., 2022), provide **color photographs of pre- and post-operative tumors**.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

＜文献紹介＞


JAVMA AVMA

Excisional biopsy and radiotherapy for management of an olfactory neuroblastoma in an axolotl (*Ambystoma mexicanum*)

Florent Modesto, DVM[1]*; Alexandra Nicolier, DVM[2]; Clémence Hurtrel, DVM[3]; Jérôme Benoît, DVM[3]

[1]Exotic Pets Department, My Exotic Vet SRL, Clinique Vétérinaire Brasseur, Manage, Belgium
[2]Vet Diagnostics, Charbonnières-les-Bains, France
[3]Department of Radiotherapy, Oncovet Veterinary Clinic, Villeneuve-d'Ascq, France

*Corresponding author: Dr. Modesto (florent.modesto@gmail.com)

https://doi.org/10.2460/javma.20.09.0498

Excisional biopsy and radiotherapy for management of an olfactory neuroblastoma in an axolotl (Ambystoma mexicanum) in: Journal of the American Veterinary Medical Association Volume 260 Issue 4 (2022)

アホロートル（*Ambystoma mexicanum*）における嗅神経芽細胞腫（ONB）の管理のための切除生検と放射線治療

1. **症例概要**
   - 対象：4 歳、オス、リューシスティックのアホロートル（*Ambystoma mexicanum*）
   - 主訴：2 週間続く食欲低下（摂食困難）と嚥下障害
   - 所見：口蓋前方に直径約 1 cm のピンク色腫瘤、左鼻腔への浸潤が疑われた
2. **診断**
   - 腫瘤は辺縁切除されたが、**病理学的に嗅神経芽細胞腫（ONB）**と診断。
   - 切除は不完全で、5 週後に再発疑い（白色腫瘤＋粘膜の発赤、治癒遅延、軽度顔面変形）。
   - 組織像：小型腫瘍細胞が管状構造や多層ロゼットを形成、神経外胚葉性腫瘍に一致。
3. **治療**
   - **補助療法として電子線放射線治療（6 MeV）を実施**
     - 週 1 回 × 4 回、1 回 8 Gy、総線量 32 Gy
     - 麻酔下で頭部にウェットガーゼ（ボーラス材）を置き、口腔内に鉛板を挿入して下顎を保護
   - 副作用：急性の有害反応は認められず、粘膜の発赤は 3 回目で消失
4. **転帰**
   - 放射線治療終了から 2 か月後まで再発なし、臨床的に良好
   - 3.5 か月後（腫瘍切除から約 8 か月後）、**飼育環境管理の失敗（高温による水質悪化）で死亡**
   - 剖検：腫瘍細胞は確認されず → **完全寛解が成立していた**と判断
5. **臨床的意義**
   - **アホロートルにおける ONB に対する放射線治療の初報告**
   - 口蓋腫瘤の鑑別診断に ONB を考慮すべき
   - 人・獣医領域での標準治療（外科切除＋放射線療法）の知見と合致
   - 放射線がアホロートルの組織再生能力にどう影響するかは不明だが、今回の症例では有効性が示唆された

まとめ

- アホロートルにも **嗅神経芽細胞腫（ONB）** が自然発生する。
- 外科切除のみでは不完全除去のリスクがあり、**放射線療法が有効であった初例**。
- 本症例は 3 か月以上の完全寛解が確認され、臨床的に放射線療法が両生類腫瘍治療の選択肢となりうることを示す。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。


はるらぼ！ HAL lab.axolotl



Figure 1—Photograph of a 4-year-old sexually intact male axolotl (*Ambystoma mexicanum*) evaluated because of a 2-week history of dysrexia and difficulty swallowing. A 1-cm-diameter, irregular, pink intraoral mass adherent to the left side of the rostral portion of the palate can be seen to the right of the cotton swab.



Figure 2—Photograph obtained after excisional biopsy of the oral mass. An oronasal fistula was present, allowing communication with the left nasal fossa, as confirmed with a 20-gauge polypropylene IV catheter with the stylet removed.






Figure 3—Photomicrographs of a section of the oral mass; the mass was identified as an olfactory neuroblastoma. A—Neoplastic cells are arranged in tubular structures or multilayered rosettes embedded in a loose collagen stroma. The tubules sporadically contain proteinaceous material and cellular debris. H&E stain; bar = 500 µm. B—Neoplastic cells are polygonal to cylindrical with fibrillar eosinophilic cytoplasm and a rounded to elongated nucleus with finely stippled chromatin. H&E stain; bar = 100 µm.

図 1 — 4 歳の繁殖経験のないオスのアホロートル (*Ambystoma mexicanum*)。2 週間続く食欲不振と嚥下困難のために評価された。口腔内の綿棒の右側に、口蓋前方左側に付着した直径 1 cm の不整形でピンク色の腫瘤が確認できる。

図 2 — 口腔内腫瘤の切除生検後に撮影した写真。口腔鼻瘻が存在し、左鼻腔と交通していたことが、スタイレットを除去した 20 ゲージのポリプロピレン静脈カテーテルで確認された。

図 3 — 口腔内腫瘤の切片の顕微鏡写真。腫瘤は嗅神経芽細胞腫と同定された。
A — 腫瘍細胞は疎な膠原線維性間質の中に管状構造や多層性ロゼットを形成して配列している。管腔内にはタンパク質性物質や細胞残渣が散発的に含まれる。H&E 染色；スケールバー＝500 µm。
B — 腫瘍細胞は多角形から円柱状で、線維状の好酸性細胞質を有し、核は円形から長楕円形で、微細に点状のクロマチンを含んでいる。H&E 染色；スケールバー＝100 µm。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 4—Photograph obtained 5 weeks after excisional biopsy of the oral mass. Persistent erythema and delayed healing of the palate can be seen (arrow).

trast-enhanced full-body CT was performed with a 64-slice CT unit (Brilliance CT 64 channel DS; Philips NV) for staging purposes. To undergo CT, the axolotl was anesthetized by placing it in a bath of fresh water with alfaxalone (5 mg/L [0.02 mg/kg]). Branchial and transcutaneous irrigation with alfaxalone (drop-by-drop administration of 15 mg of alfaxalone diluted in saline [0.9% NaCl] solution [62.8 mg/kg]) was then performed until the axolotl was completely immobile. Contrast-enhanced CT was not performed because of budgetary considerations. No clinically important findings were observed on CT images, possibly because of the lack of spatial resolution for



Figure 5—Photograph obtained during radiotherapy. A 5-mm tissue-equivalent bolus material (wet gauze) has been placed on the animal's head to optimize surface dose. The treatment couch has been rotated by 90°, and the gantry has been angled so that the electron applicator (6 X 6 cm with blocks at each corner to create a hexagonal shape) is parallel to the skin.

図 4 — 口腔内腫瘤の切除生検から 5 週後に撮影した写真。口蓋の持続的な発赤と治癒遅延が確認できる（矢印）。

図 5 — 放射線治療中に撮影した写真。表面線量を最適化するため、動物の頭部に厚さ 5 mm の組織等価ボーラス材（湿らせたガーゼ）が置かれている。治療台は 90°回転され、ガントリーは電子アプリケータ（各角にブロックを配置し六角形を形成した 6×6 cm）を皮膚と平行にするよう角度調整されている。

図 6 — 放射線治療中に撮影した写真。動物は胸臥位に配置され、下顎を放射線曝露から保護するために、特注の厚さ 2 mm の鉛板が口腔内に挿入されている。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Table 1—Anesthetic protocols used during radiotherapy of an olfactory neuroblastoma in a 4-year-old sexually intact male axolotl (*Ambystoma mexicanum*).

Dose of alfaxalone	Mean (range) TME (min)	Immobility	RGS	Jaw resistance	RMI
30 mg (125.5 mg/kg)	11 (10–12)	+/−	−	−	−
40 mg (167.4 mg/kg)	15.25 (14–17)	+	+/−	−	−
50 mg (209.2 mg/kg)	20.75 (20–22)	+	+	+	+/−
60 mg (251 mg/kg)	29 (26–31)	+	+	+	+

The anesthetic protocol involved mixing 1 mL of alfaxalone (10 mg/mL solution) in 1 mL of saline (0.9% NaCl) solution. This mixture, which contained 10 mg of alfaxalone (41.8 mg/kg), was then instilled by branchial and transcutaneous irrigation for over 1 to 2 minutes. After 2 minutes, immobility was evaluated by checking for a righting reflex. As long as the reflex was still present, an additional 2-mL mixture was prepared and applied in the same way. This process was repeated until immobility and the other desired effects were observed. Effects were scored as present for at least 15 minutes (+), present but not maintained for > 2 minutes (+/−), or absent (−).

RGS = Resistance to gauze stimulation (ie, placement of wet gauze [5-mm tissue-equivalent bolus material] on the animal's head). RMI = Resistance to insertion of a 2-mm-thick lead plate into the animal's mouth for beam-blocking purposes.TME = Time to reach maximal effect.

表 1 — 4 歳の未去勢オスのアホロートル（*Ambystoma mexicanum*）における嗅神経芽細胞腫の放射線治療中に使用された麻酔プロトコル。

投与量 (アルファキサロン)	TME* (分，平均[範囲])	下顎の不動性	RGS 抵抗性	RMI 抵抗性
30 mg (125.5 mg/kg)	11 (10–12)	+/−	−	−
40 mg (167.4 mg/kg)	15.25 (14–17)	+	+/−	+/−
50 mg (209.2 mg/kg)	20.75 (20–22)	+	+	+
60 mg (251 mg/kg)	29 (26–31)	+	+	+

注釈

- 麻酔プロトコル：アルファキサロン（10 mg/mL 溶液）1 mL を生理食塩水（0.9% NaCl）1 mL で希釈して混合。この混合液（アルファキサロン 10 mg, 41.8 mg/kg 含有）を、鰓および皮膚経由で 1〜2 分かけて灌流投与した。
- 2 分後に正向反射（righting reflex）を確認し、不動性を評価。反射が残っている場合は、同じ方法でさらに 2 mL の混合液を追加。これを目的の効果が得られるまで繰り返した。
- 効果判定：
  - **+**：少なくとも 15 分間効果が持続
  - **+/−**：効果はあるが 2 分以上持続しない
  - **−**：効果なし

略語

- **RGS** = ガーゼ刺激に対する抵抗（例：頭部に 5 mm 厚の組織等価ボーラス材［湿らせたガーゼ］を置いたときの反応）
- **RMI** = 口腔内に 2 mm 厚の鉛板を挿入しビーム遮蔽する際の抵抗
- **TME** = 最大効果発現までの時間

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

＜両生類におけるマーカー遺伝子＞

1. 幹細胞・再生関連マーカー

- **Sox2**：神経幹細胞、前駆細胞のマーカー
- **Pax7**：筋衛星細胞（筋幹細胞）のマーカー
- **Nanog, Oct4 (Pou5f1)**：多能性幹細胞マーカー
- **Klf4, c-Myc**：リプログラミングや細胞増殖に関連

---

2. 神経系マーカー

- **GFAP (Glial fibrillary acidic protein)**：アストロサイト
- **βIII-tubulin (Tuj1)**：ニューロン
- **Nestin**：神経前駆細胞
- **HuC/D**：分化したニューロン

---

3. 筋・組織修復関連

- **MyoD, Myf5, Myogenin**：筋分化マーカー
- **Desmin**：筋細胞マーカー
- **Laminin**：基底膜関連

---

4. 血管・間葉系

- **VEGFR, Flk1 (Kdr)**：血管新生
- **CD31 (PECAM-1)**：血管内皮細胞
- **Vimentin**：間葉系マーカー
- **Fibronectin**：創傷修復に重要

---

5. 腫瘍・増殖関連

- **PCNA, Ki-67**：細胞増殖マーカー
- **p53**：腫瘍抑制因子
- **c-Met, FGFR**：再生と腫瘍形成の両面に関連
- **MMPs (Matrix metalloproteinases)**：組織再構築や腫瘍浸潤

---

6. 両生類特有・よく使われる分子

- **Prod1**：サラマンダーの再生で重要（近位–遠位アイデンティティ）
- **EfnA (Ephrin-A)**：再生における細胞位置情報
- **Hox 遺伝子群**：発生と再生パターン形成
- **Shh, Wnt, Fgf, BMP**：再生芽形成の主要シグナル

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

＜両生類における腫瘍マーカーの報告例＞

両生類では腫瘍研究自体が少なく、哺乳類ほど確立された「腫瘍マーカー」はありません。
ただし、**再生と腫瘍の共通性**を軸に、多くの分子がマーカー候補として注目されています。
以下、**主要なマーカー群と作用機序**を整理します。

アホロートル・両生類における腫瘍マーカーと作用機序

1. 細胞増殖マーカー

- **PCNA (Proliferating Cell Nuclear Antigen)**
  - DNA 複製に必須のクロモソーム複製因子
  - 腫瘍細胞で強発現 → 腫瘍の増殖能を反映
  - 両生類では再生芽・腫瘍の両方で増殖指標に用いられる
- **Ki-67**
  - 分裂期にある細胞核で発現
  - 腫瘍増殖指数の標準マーカー（哺乳類同様）

---

2. 腫瘍抑制・ゲノム安定性関連

- **p53**
  - DNA 損傷応答、アポトーシス誘導
  - 両生類でも保存されており、再生過程では抑制的に働く一方、腫瘍形成時には変異や不活化が関与する可能性
  - Lucké 腫瘍（Rana pipiens 腎腫瘍）では p53 経路の異常が関与と推測
- **Rb (Retinoblastoma protein)**
  - 細胞周期の G1/S チェックポイントを制御
  - 両生類の腫瘍細胞で不活化されると、制御不能な増殖が進行

---

3. シグナル伝達系マーカー

- **FGFR (Fibroblast Growth Factor Receptor)**
  - 再生芽形成と腫瘍細胞の両方で発現増加
  - 再生では組織再構築を促進するが、腫瘍では異常な血管新生・増殖を誘導
- **Wnt/β-catenin**
  - 再生芽で軸形成や細胞増殖を促進
  - 異常活性化で腫瘍性増殖を誘導
  - Xenopus 腫瘍モデルで発がん関連経路として検証されている
- **Shh (Sonic Hedgehog)**
  - 発生・再生でパターン形成を制御
  - 腫瘍では細胞分裂と幹細胞性維持を亢進

---

4. 浸潤・転移関連マーカー

- **MMPs (Matrix Metalloproteinases)**
  - 細胞外基質を分解
  - 腫瘍浸潤や再生組織のリモデリングに関与
  - 両生類腫瘍でも高発現が確認されている

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **Vimentin**
    - 上皮間葉転換（EMT）のマーカー
    - 再生芽細胞と腫瘍細胞で共通に発現
    - 浸潤・移動性の指標

---

5. 幹細胞性・多能性関連マーカー

- **Sox2**
    - 神経幹細胞や腫瘍幹細胞マーカー
    - 再生芽で活性化 → 腫瘍化では異常に持続発現
- **Oct4, Nanog, Klf4, c-Myc**
    - 多能性維持因子
    - 再生の初期には一時的に活性化
    - 腫瘍では「自己複製を失わない細胞集団」の維持に寄与

---

6. 両生類特有の因子

- **Prod1**（サラマンダー特異的再生マーカー）
    - 再生の近位-遠位アイデンティティを規定
    - 腫瘍では異常発現し、細胞増殖や組織位置情報の破綻に関与する可能性
- **Eph/Ephrin シグナル**
    - 再生で細胞境界や組織再構築に必須
    - 腫瘍では細胞の異常な境界破壊や転移促進に関連

まとめ

- **増殖系（PCNA, Ki-67）**は腫瘍の基本指標。
- **腫瘍抑制系（p53, Rb）**の異常は、両生類腫瘍でも重要。
- **再生シグナル（FGF, Wnt, Shh）**は、再生と腫瘍の「分岐点」となる。
- **浸潤系（MMPs, Vimentin）**は腫瘍悪性度の指標。
- **幹細胞性（Sox2, Oct4）**は腫瘍幹細胞の存在を示唆。
- **両生類特有（Prod1, Ephrin）**は哺乳類と異なる腫瘍制御の手がかり。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl


ハルラボにおける知見

アホロートルにおいて、腫瘍マーカーは損傷再生時にも必要とされるものも含まれる。これまでの炎症再生や細菌感染、内在性ウイルス配列の発現などの機序に関する知見を踏まえ、損傷部位への細菌感染の影響による腫瘍悪性化の詳細な機構が想定された。

検証の結果、

“**損傷→損傷部位で細菌が増殖→抗菌ペプチドによる感知・免疫応答→損傷部位が炎症＋内在性ウイルス配列の活性化→未分化な組織・幹細胞の過剰増殖（+変異の誘発)→腫瘍の形成→悪性化（がん化)**“


のような機構を持つ可能性が示唆された。





補足図 1: アホロートルにおける腫瘍マーカーの発現動態 1


(腫瘍 または 周辺領域 における発現)


※ 各検証については、内容と必要に応じて、専門家の方々のご指導･ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

















補足図 2: アホロートルにおける腫瘍マーカーの発現動態 2


(損傷部位における発現 1)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。










補足図 3: アホロートルにおける腫瘍マーカーの発現動態 3


(損傷部位における発現 2)


CC BY NC ND


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl










補足図 4: アホロートルにおける腫瘍マーカーの発現動態 3

(損傷なしの場合)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




補足図 5: アホロートルにおける腫瘍マーカーの発現動態 4

(損傷あり、患部を滅菌した場合)

CC BY NC ND

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl



補足図 5: アホロートルにおける損傷に起因した
細菌感染症・内在性ウイルス配列・炎症・腫瘍化 の相関モデル図

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

参考文献　References

アホロートル腫瘍マーカー

PCNA

https://genome.axolotl-omics.org/cgi-bin/hgTracks?db=ambMex60DD&lastVirtModeType=default&lastVirtModeExtraState=&virtModeType=default&virtMode=0&nonVirtPosition=&position=chr10p%3A614507051%2D614514790&hgsid=69715_5uEAztFlj0qCx310CTXrTpTbcU5J

p53

- **ホモログ（本体）** → Tp53（7 アイソフォーム）
- **結合因子** → Tp53bp1, Tp53bp2
- **誘導因子** → Tp53i3, Tp53i11, Tp53inp1, Tp53inp2
- **修飾因子** → Tp53rk

p53 本体（ホモログ）

- **Tp53**
  - AMEX60DD201012903.1
  - AMEX60DD201012903.2
  - AMEX60DD201012903.3
  - AMEX60DD301012903.4
  - AMEX60DD301012903.5
  - AMEX60DD201012903.6
  - AMEX60DD102012903.7

---

2 p53 発現調節因子

- **Wrap53**（p53 mRNA の安定性・転写後制御）
  - AMEX60DD301013012.1
  - AMEX60DD201013012.2
  - AMEX60DD301013012.3

---

3 p53 結合タンパク質（Binding Proteins）

- **Tp53bp1**
  - AMEX60DD201003378.1–7
  - AMEX60DD301003378.8–9
- **Tp53bp2**（ASPP2）
  - AMEX60DD201036184.1–6
  - AMEX60DD301036184.7

---

4 p53 誘導/相互作用因子

- **Tp53i3**（p53-inducible protein 3）
  - AMEX60DDU001011613.1
  - AMEX60DD301033010.3, .6–11

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- **Tp53i11**
  - AMEX60DD102005196.2
  - AMEX60DD201005196.1, .3–4
- **Tp53i13**
  - AMEX60DD201054814.1–5, .7–8
  - AMEX60DD301054814.3, .9–10
  - AMEX60DD102054814.6
- **Tp53inp1**
  - AMEX60DD301040136.3
  - AMEX60DD201040136.7
- **Tp53inp2**
  - AMEX60DD201028587.3–5

5️⃣ p53 経路修飾因子

- **Tp53rk**（PRPK, kinase）
  - AMEX60DD201027604.1–5

6️⃣ p53 関連タンパク質（その他）

- **Nop53**（Nucleolar protein 53, aka GLTSCR2 → p53 安定化に関与）
  - AMEX60DD201024441.1
  - AMEX60DD301024442.1
- **Usp53**（Deubiquitinase family, p53 経路のユビキチン調節）
  - AMEX60DD301044229.1–2
  - AMEX60DD201044228.1
  - AMEX60DD201044230.1–2
- **Cfap53**（Cilia/flagella-associated protein 53, 直接の腫瘍マーカーではないが命名上関連）
  - AMEX60DD301041749.1
- **Tp53tg5**（Tumor protein p53 target gene 5）
  - AMEX60DD301051605.1–3

☑ まとめ

- **本体ホモログ**：Tp53
- **直接結合因子**：Tp53bp1, Tp53bp2
- **誘導因子**：Tp53i3, Tp53i11, Tp53i13, Tp53inp1, Tp53inp2
- **修飾因子**：Tp53rk
- **発現調節**：Wrap53
- **関連遺伝子**：Nop53, Usp53, Cfap53, Tp53tg5

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.73 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.83 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.84 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.85 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.87 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.97 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.98 at C0161316:12458-14720
Zfp534 [nr]|Znf568 [hs]|AMEX60DDU001026768.99 at C0161316:12458-14720
Tp53bp1|AMEX60DD301003378.8 at chr11p:19483763-19536761
Tp53bp1|AMEX60DD301003378.9 at chr11p:19579171-19650180
Nop53|AMEX60DD301024442.1 at chr2q:702457783-702495707
Tp53rk|AMEX60DD201027604.2 at chr3p:496636196-496906851
Tp53bp2|AMEX60DD301036184.7 at chr4q:990093295-990544223
Ab205_0135830 [nr]|Usp53 [hs]|AMEX60DD301044229.1 at chr6q:219577729-219607866
Usp53|AMEX60DD301044229.2 at chr6q:219577729-219643351
Usp53 [hs]|AMEX60DD201044230.1 at chr6q:219691114-219825031
Tp53i3|AMEX60DDU001011613.1 at C0107577:21422-31581
Zfp534 [nr]|AMEX60DDU001023850.26 at C0150954:20563-22223
Zfp534 [nr]|Znf300 [hs]|AMEX60DDU001029882.12 at C0172671:22919-27553
Tp53i11|AMEX60DD201005196.1 at chr11q:637055223-637926141
Tp53 [hs]|AMEX60DD201012903.3 at chr1p:117630937-117693881
Tp53 [hs]|AMEX60DD301012903.4 at chr1p:117631140-117691404
Tp53 [hs]|AMEX60DD102012903.7 at chr1p:117631357-117728710
Nop53|AMEX60DD201024441.1 at chr2q:702337522-702596568
Tp53rk|AMEX60DD201027604.5 at chr3p:496636238-496717853
Tp53i3.s [nr]|Tp53i3 [hs]|AMEX60DD301033010.9 at chr4p:260195544-260397737
Tp53i3|AMEX60DD301033010.10 at chr4p:260195544-260397737
Tp53i3.s [nr]|Tp53i3 [hs]|AMEX60DD301033010.11 at chr4p:260195571-260397737
Tp53inp1|AMEX60DD301040136.3 at chr5q:834878137-834986044
Usp53|AMEX60DD201044228.1 at chr6q:219343915-219531980
Usp53|AMEX60DD201044230.2 at chr6q:219691114-219728623
Tp53i13 [nr]|AMEX60DD201054814.1 at chr9q:99977810-100123132
Tp53i13 [nr]|AMEX60DD201054814.2 at chr9q:99977810-100123132
Tp53i13 [nr]|AMEX60DD301054814.3 at chr9q:99977827-100123132
Tp53i13 [nr]|AMEX60DD201054814.4 at chr9q:99977868-100123132
Tp53i13 [nr]|AMEX60DD201054814.5 at chr9q:99977868-100123132
Tp53i13 [nr]|AMEX60DD201054814.7 at chr9q:99977903-100123132
Tp53i13 [nr]|AMEX60DD201054814.8 at chr9q:99977903-100123132
Tp53i13 [nr]|AMEX60DD301054814.10 at chr9q:99991294-100123132
Tp53bp1|AMEX60DD201003378.1 at chr11p:19417975-19862760
Tp53bp1|AMEX60DD201003378.2 at chr11p:19418030-19862760
Tp53bp1|AMEX60DD201003378.3 at chr11p:19418031-19862760

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Tp53bp1|AMEX60DD201003378.4 at chr11p:19418031-19862760
Tp53bp1|AMEX60DD201003378.5 at chr11p:19483625-19862760
Tp53bp1|AMEX60DD201003378.6 at chr11p:19483625-19862760
Tp53bp1|AMEX60DD201003378.7 at chr11p:19483710-19862760
Tp53i11|AMEX60DD102005196.2 at chr11q:637061766-637926141
Tp53i11|AMEX60DD201005196.3 at chr11q:637061776-637926141
Tp53i11|AMEX60DD102005196.4 at chr11q:637061792-637697007
Tp53 [hs]|AMEX60DD201012903.1 at chr1p:117629256-117908984
Tp53 [hs]|AMEX60DD201012903.2 at chr1p:117630830-117908984
Tp53 [hs]|AMEX60DD301012903.5 at chr1p:117631252-117728710
Tp53 [hs]|AMEX60DD201012903.6 at chr1p:117631319-117737266
Wrap53|AMEX60DD301013012.1 at chr1p:148374293-148696165
Wrap53|AMEX60DD201013012.2 at chr1p:148374322-148696165
Wrap53|AMEX60DD301013012.3 at chr1p:148374415-148696165
Tp53rk|AMEX60DD201027604.1 at chr3p:496636158-496717853
Tp53rk [hs]|AMEX60DD201027604.3 at chr3p:496636197-496717853
Tp53rk [hs]|AMEX60DD201027604.4 at chr3p:496636220-496717853
Tp53inp2|AMEX60DD201028587.3 at chr3q:206293-312388
Tp53inp2|AMEX60DD201028587.4 at chr3q:206299-312388
Tp53inp2|AMEX60DD201028587.5 at chr3q:206322-304174
Tp53i3 [nr]|AMEX60DD301033010.3 at chr4p:259581312-260397737
Tp53i3 [nr]|AMEX60DD301033010.6 at chr4p:259945425-260397737
Tp53i3.s [nr]|AMEX60DD301033010.7 at chr4p:260195535-260397737
Tp53i3.s [nr]|AMEX60DD301033010.8 at chr4p:260195544-260397737
Tp53bp2|AMEX60DD201036184.1 at chr4q:990087409-991005780
Tp53bp2|AMEX60DD201036184.2 at chr4q:990092698-991005780
Tp53bp2|AMEX60DD201036184.3 at chr4q:990092716-991005780
Tp53bp2|AMEX60DD201036184.4 at chr4q:990092716-991005780
Tp53bp2|AMEX60DD201036184.5 at chr4q:990092716-991005780
Tp53bp2|AMEX60DD201036184.6 at chr4q:990093128-991005780
Tp53inp1|AMEX60DD201040136.7 at chr5q:834916303-834986044
Cfap53|AMEX60DD301041749.1 at chr6p:214039492-214258285
Tp53tg5 [nr]|AMEX60DD301051605.1 at chr8p:626234854-626418806
Tp53tg5 [nr]|AMEX60DD301051605.2 at chr8p:626234854-626418806
Tp53tg5 [nr]|AMEX60DD301051605.3 at chr8p:626234854-626418806
Tp53i13 [nr]|AMEX60DD102054814.6 at chr9q:99977871-100123132
Tp53i13 [nr]|AMEX60DD301054814.9 at chr9q:99977924-99988435
Zfp534 [nr]|AMEX60DD301056435.1 at chr9q:948705387-948770946

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Wnt

1. **Wnt-1**
    - AMEX60DD301029979.3 at chr3q:597308876-597436446
    - AMEX60DD102029980.1 at chr3q:597578800-597595498
    - AMEX60DD201029979.1 at chr3q:597308375-597406551
    - AMEX60DD201029979.2 at chr3q:597308753-597436772
2. **Wnt2**
    - AMEX60DD301006323.1 at chr12p:45454998-45566622
3. **Wnt2b**
    - AMEX60DD201008914.1 at chr13p:171507799-171780458
    - AMEX60DD201008914.2 at chr13p:171508286-171627699
4. **Wnt3**
    - AMEX60DD301009763.1 at chr13q:177285038-177300157
5. **Wnt3a**
    - AMEX60DD201020503.1 at chr2p:142830045-143086654
6. **Wnt4**
    - AMEX60DD201052091.1 at chr8q:176182931-176406102
    - AMEX60DD301052091.2 at chr8q:176189471-176406102
    - Loc102366334 [nr]|Wnt4 [hs]|AMEX60DD301026872.1 at chr3p:168432279-168437749
7. **Wnt5a**
    - AMEX60DD201023726.1 at chr2q:305821600-305851377
    - AMEX60DD201023726.2 at chr2q:305821600-305851377
    - AMEX60DD201023726.3 at chr2q:305821601-305851377
    - AMEX60DD201023726.4 at chr2q:305821613-305845728
8. **Wnt5b**
    - AMEX60DD201006911.1 at chr12q:85888522-86397186
    - AMEX60DD201006911.2 at chr12q:86094484-86397186
    - AMEX60DD201006911.3 at chr12q:86317278-86397186
    - AMEX60DD201006911.4 at chr12q:86317389-86397186
    - AMEX60DD201006911.5 at chr12q:86317720-86397186
    - AMEX60DD201006911.6 at chr12q:86318221-86397186
    - AMEX60DD201006911.7 at chr12q:86318564-86397186
9. **Wnt6**
    - AMEX60DD301056290.1 at chr9q:900932036-901283623
    - AMEX60DD201056290.2 at chr9q:900934305-901283623
    - AMEX60DD201056290.3 at chr9q:900934604-901283623
10. **Wnt7a**
    - AMEX60DD201023665.1 at chr2q:265350467-265516707
11. **Wnt7b**
    - AMEX60DD201006490.1 at chr12p:132702652-133282449
    - AMEX60DD201006490.2 at chr12p:132702652-133108508
    - AMEX60DD201006490.3 at chr12p:132709705-133108508

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- AMEX60DD301025497.1 at chr2q:1107554290-1107840098

12. **Wnt8a / Wnt8b**
    - AMEX60DD201029006.1 at chr3q:229703683-229708722
    - AMEX60DD201029006.2 at chr3q:229703690-229708722
    - AMEX60DD201042933.2 at chr6p:958018119-958207134
    - AMEX60DD201051636.1 at chr8p:649199585-649285989
    - AMEX60DD201051636.2 at chr8p:649202315-649863618
13. **Wnt9a / Wnt9b**
    - AMEX60DD201009761.1 at chr13q:176819446-176879846 (Wnt9b)
    - AMEX60DD301020504.1 at chr2p:144105954-144359488 (Wnt9a)
14. **Wnt10a / Wnt10b**
    - AMEX60DD201056292.1 at chr9q:902691362-902801962 (Wnt10a)
    - AMEX60DD103056292.2 at chr9q:902691699-902723743 (Wnt10a)
    - AMEX60DD201029981.1~.8 at chr3q:598220670-598634383 (Wnt10b)
15. **Wnt11**
    - AMEX60DD201036922.1~.3 at chr5p:59051445-59064733
    - AMEX60DD201049747.1~.3 at chr7q:929503276-929786383

Wnt2|AMEX60DD301006323.1 at chr12p:45454998-45566622

Loc102030191 [nr]|Wnt3 [hs]|AMEX60DD301009763.1 at chr13q:177285038-177300157

Wnt7c.s [nr]|Wnt7b [hs]|AMEX60DD301025497.1 at chr2q:1107554290-1107840098

Wnt7b|AMEX60DD201006490.3 at chr12p:132709705-133108508

Wnt3a|AMEX60DD201020503.1 at chr2p:142830045-143086654

Wnt-1 [nr]|Wnt1 [hs]|AMEX60DD301029979.3 at chr3q:597308876-597436446

Wnt-1 [nr]|Wnt1 [hs]|AMEX60DD102029980.1 at chr3q:597578800-597595498

Wnt10b|AMEX60DD201029981.1 at chr3q:598220670-598634383

Wnt10b|AMEX60DD301029981.2 at chr3q:598220681-598634383

Wnt10b|AMEX60DD301029981.3 at chr3q:598220703-598634383

Wnt10b|AMEX60DD201029981.5 at chr3q:598543998-598634383

Wnt10b|AMEX60DD201029981.6 at chr3q:598543998-598634383

Wnt10b|AMEX60DD201029981.7 at chr3q:598543998-598634383

Wnt10b|AMEX60DD201029981.8 at chr3q:598544227-598634383

Wnt8 [nr]|Wnt8b [hs]|AMEX60DD201042933.2 at chr6p:958018119-958207134

Wnt8b|AMEX60DD201051636.1 at chr8p:649199585-649285989

Wnt6.s [nr]|AMEX60DD301056290.1 at chr9q:900932036-901283623

Wnt6.s [nr]|AMEX60DD201056290.3 at chr9q:900934604-901283623

Pwnt10a [nr]|Wnt10a [hs]|AMEX60DD103056292.2 at chr9q:902691699-902723743

Wnt7b|AMEX60DD201006490.1 at chr12p:132702652-133282449

Wnt7b|AMEX60DD201006490.2 at chr12p:132702652-133108508

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.1 at chr12q:85888522-86397186

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.2 at chr12q:86094484-86397186

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.3 at chr12q:86317278-86397186

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.4 at chr12q:86317389-86397186

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.5 at chr12q:86317720-86397186

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.6 at chr12q:86318221-86397186

Wnt-5b [nr]|Wnt5b [hs]|AMEX60DD201006911.7 at chr12q:86318564-86397186

Wnt2b|AMEX60DD201008914.1 at chr13p:171507799-171780458

Wnt2b|AMEX60DD201008914.2 at chr13p:171508286-171627699

Wnt9b|AMEX60DD201009761.1 at chr13q:176819446-176879846

Wnt9a|AMEX60DD301020504.1 at chr2p:144105954-144359488

Wnt7a|AMEX60DD201023665.1 at chr2q:265350467-265516707

Wnt-5a [nr]|Wnt5a [hs]|AMEX60DD201023726.1 at chr2q:305821600-305851377

Wnt-5a [nr]|Wnt5a [hs]|AMEX60DD201023726.2 at chr2q:305821600-305851377

Wnt-5a [nr]|Wnt5a [hs]|AMEX60DD201023726.3 at chr2q:305821601-305851377

Wnt-5a [nr]|Wnt5a [hs]|AMEX60DD201023726.4 at chr2q:305821613-305845728

Loc102366334 [nr]|Wnt4 [hs]|AMEX60DD301026872.1 at chr3p:168432279-168437749

Wnt-8 [nr]|Wnt8a [hs]|AMEX60DD201029006.1 at chr3q:229703683-229708722

Wnt-8 [nr]|Wnt8a [hs]|AMEX60DD201029006.2 at chr3q:229703690-229708722

Wnt-1 [nr]|Wnt1 [hs]|AMEX60DD201029979.1 at chr3q:597308375-597406551

Wnt-1 [nr]|Wnt1 [hs]|AMEX60DD201029979.2 at chr3q:597308753-597436772

Loc108785620 [nr]|Wnt11 [hs]|AMEX60DD201036922.1 at chr5p:59051445-59064733

Loc108785620 [nr]|Wnt11 [hs]|AMEX60DD201036922.2 at chr5p:59051649-59064733

Loc108785620 [nr]|Wnt11 [hs]|AMEX60DD201036922.3 at chr5p:59051653-59064733

Wnt11|AMEX60DD201049747.1 at chr7q:929503276-929786383

Wnt11|AMEX60DD201049747.2 at chr7q:929503570-929786383

Wnt11|AMEX60DD301049747.3 at chr7q:929505105-929786383

Wnt8b|AMEX60DD201051636.2 at chr8p:649202315-649863618

Wnt4|AMEX60DD201052091.1 at chr8q:176182931-176406102

Wnt4|AMEX60DD301052091.2 at chr8q:176189471-176406102

Wnt6.s [nr]|Wnt6 [hs]|AMEX60DD201056290.2 at chr9q:900934305-901283623

Pwnt10a [nr]|Wnt10a [hs]|AMEX60DD201056292.1 at chr9q:902691362-902801962

Shh

https://genome.axolotl-omics.org/cgi-bin/hgTracks?db=ambMex60DD&lastVirtModeType=default&lastVirtModeExtraState=&virtModeType=default&virtMode=0&nonVirtPosition=&position=chr2p%3A687768568%2D687801925&hgsid=69715_5uEAztFlj0qCx310CTXrTpTbcU5J

MMPs

1. **Mmp1**
    - Loc115468492 [nr]|Mmp1 [hs]|AMEX60DD301049489.1
    - Loc115468492 [nr]|Mmp1 [hs]|AMEX60DD301049489.2
    - Col [nr]|Mmp1 [hs]|AMEX60DD201049490.1
2. **Mmp2**
    - Mmp2|AMEX60DD301017764.4

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- Mmp2|AMEX60DD301017764.1
- Mmp2|AMEX60DD102017764.2
- E2i00_006438 [nr]|Mmp2 [hs]|AMEX60DD102017764.3

3. **Mmp3 / Mmp3/10a / Mmp3/10b**
   - Mmp3/10a [nr]|Mmp3 [hs]|AMEX60DDU001041749.1
   - Mmp3/10a [nr]|Mmp3 [hs]|AMEX60DD102049495.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD301049496.5
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.6
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049493.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD102049494.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD101049496.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.3
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.4
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.9
   - Mmp3/10b [nr]|Mmp3 [hs]|AMEX60DD301049492.1
   - Mmp3/10b [nr]|Mmp13 [hs]|AMEX60DD201049496.2
4. **Mmp7**
   - Mmp7|AMEX60DD301049503.2
   - Mmp7|AMEX60DD201049503.3
5. **Mmp8**
   - Mmp8 [hs]|AMEX60DD301049496.7
   - Mmp8 [hs]|AMEX60DD301049497.1
6. **Mmp10**
   - Loc115091995 [nr]|Mmp10 [hs]|AMEX60DD201049491.2
7. **Mmp11**
   - Mmp11|AMEX60DD201000336.1
   - Mmp11|AMEX60DD301000336.2
8. **Mmp13**
   - Loc115469592 [nr]|Mmp13 [hs]|AMEX60DD301049487.1
   - Loc115469592 [nr]|Mmp13 [hs]|AMEX60DD301049487.2
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD301049496.5
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.6
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049493.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD102049494.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD101049496.1
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.3
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.4
   - Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.9
   - Mmp3/10b [nr]|Mmp13 [hs]|AMEX60DD201049496.2
   - Mmp13|AMEX60DD301049488.1
   - Loc115091500 [nr]|Mmp13 [hs]|AMEX60DD301049487.3
   - Loc115091995 [nr]|Mmp13 [hs]|AMEX60DD201049491.1

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

- Mmpe [nr]|Mmp13 [hs]|AMEX60DD201049502.1

9. **Mmp15**
    - Dr999_pmt17610 [nr]|Mmp15 [hs]|AMEX60DD301015494.1
    - Dr999_pmt17610 [nr]|Mmp15 [hs]|AMEX60DD301015494.2
10. **Mmp16**
    - Loc104377790 [nr]|Mmp16 [hs]|AMEX60DD102040043.1
    - Mmp16|AMEX60DD301040046.1
    - Mmp16|AMEX60DD301040046.2
11. **Mmp17**
    - Mmp17 [hs]|AMEX60DD201000108.1
    - Mmp17|AMEX60DD201000108.2
    - Cib84_011884 [nr]|Mmp17 [hs]|AMEX60DD301000108.3
    - Mmp17|AMEX60DD301000108.4
    - Mmp17|AMEX60DD301000108.5
    - Mmp17|AMEX60DDU001018204.1
    - Mmp17|AMEX60DD201000119.1
    - Loc115075181 [nr]|Mmp17 [hs]|AMEX60DD201007135.1
    - Loc115460504 [nr]|Mmp17 [hs]|AMEX60DD201007135.2
    - Loc115460504 [nr]|Mmp17 [hs]|AMEX60DD201007135.3
12. **Mmp19**
    - Mmp19|AMEX60DD301029882.2
    - Mmp19|AMEX60DD301029882.1
    - Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD201012011.4
    - Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD201012011.5
    - Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD102012011.6
13. **Mmp21**
    - Mmp21|AMEX60DD201052767.1
    - Mmp21|AMEX60DD201052767.2
    - Loc115079846 [nr]|Mmp21 [hs]|AMEX60DD301030028.1
    - Loc115079846 [nr]|Mmp21 [hs]|AMEX60DD301030028.2
    - Loc115461210 [nr]|Mmp21 [hs]|AMEX60DD201046376.1
14. **Mmp23b**
    - Mmp23b|AMEX60DD301001496.1
    - Loc115081292 [nr]|Mmp23b [hs]|AMEX60DD301001498.1
    - Mmp23b|AMEX60DD201051302.1
15. **Mmp24**
    - Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.1
    - Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.3
    - Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.7
    - Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.8
    - Mmp24|AMEX60DD301028519.2
    - Mmp24|AMEX60DD201028519.1

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

16. **Mmp25**
    - Mmp25|AMEX60DD201028039.1
17. **Mmp28**
    - Mmp28|AMEX60DD102054403.1
    - Mmp28|AMEX60DD201054403.1
    - Mmp28|AMEX60DD102054403.2
    - Mmp28|AMEX60DD102054403.4
    - Mmp28|AMEX60DD201054403.6
    - Mmp28|AMEX60DD201054403.2
    - Mmp28|AMEX60DD301054403.5
    - Mmp28|AMEX60DD301054403.7
    - Mmp28|AMEX60DD301054403.8

Mmp3/10a [nr]|Mmp3 [hs]|AMEX60DDU001041749.1 at C0215691:7707-10646

Mmp17 [hs]|AMEX60DD201000108.1 at chr10p:51227918-51319614

Mmp17|AMEX60DD201000108.2 at chr10p:51227918-51472496

Cib84_011884 [nr]|Mmp17 [hs]|AMEX60DD301000108.3 at chr10p:51299302-51319614

Mmp17|AMEX60DD301000108.4 at chr10p:51358436-51472496

Mmp17|AMEX60DD301000108.5 at chr10p:51429194-51472496

Mmp23b|AMEX60DD301001496.1 at chr10p:666006004-666186054

Loc115081292 [nr]|Mmp23b [hs]|AMEX60DD301001498.1 at chr10p:666192820-666198343

Mmp19|AMEX60DD301029882.2 at chr3q:554393966-554437096

Loc104377790 [nr]|Mmp16 [hs]|AMEX60DD102040043.1 at chr5q:774175577-774175796

Loc115469592 [nr]|Mmp13 [hs]|AMEX60DD301049487.1 at chr7q:767985535-767995563

Loc115469592 [nr]|Mmp13 [hs]|AMEX60DD301049487.2 at chr7q:767986359-767995563

Mmp8 [hs]|AMEX60DD301049496.7 at chr7q:771485625-771488262

Mmp8 [hs]|AMEX60DD301049497.1 at chr7q:771519786-771522410

Mmp28|AMEX60DD102054403.3 at chr9p:396416298-396573178

Mmp28|AMEX60DD201054404.1 at chr9p:396497780-396524900

Mmp28|AMEX60DD301054403.5 at chr9p:396567418-396725221

Mmp28|AMEX60DD201054403.6 at chr9p:396567418-396944082

Mmp17|AMEX60DDU001018204.1 at C0131207:11545-34231

Mmp17|AMEX60DD201000119.1 at chr10p:56030701-56053122

Immp1l|AMEX60DD301004910.3 at chr11q:427339667-427929566

Loc115075181 [nr]|Mmp17 [hs]|AMEX60DD201007135.1 at chr12q:183287174-183581527

Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.3 at chr14q:309024990-309136701

Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD201012011.4 at chr14q:309024990-309136701

Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.7 at chr14q:309028376-309136701

Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.8 at chr14q:309050845-309136701

Dr999_pmt17610 [nr]|Mmp15 [hs]|AMEX60DD301015494.2 at chr1p:930477188-930716207

Mmp2|AMEX60DD301017764.4 at chr1q:389611825-390000101

Mmp24|AMEX60DD301028519.2 at chr3p:830862619-830990272

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Mmp19|AMEX60DD301029882.1 at chr3q:554393943-554447184
Loc115468492 [nr]|Mmp1 [hs]|AMEX60DD301049489.1 at chr7q:768998327-769047841
Loc115468492 [nr]|Mmp1 [hs]|AMEX60DD301049489.2 at chr7q:768998406-769047841
Mmp3/10a [nr]|Mmp3 [hs]|AMEX60DD102049495.1 at chr7q:771347771-771360149
Mmp3/10b [nr]|Mmp13 [hs]|AMEX60DD201049496.2 at chr7q:771474205-771498013
Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD301049496.5 at chr7q:771474265-771488263
Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.6 at chr7q:771474353-771498013
Mmp3/10a [nr]|Mmp3 [hs]|AMEX60DD201049496.8 at chr7q:771488468-771548766
Mmp7|AMEX60DD301049503.2 at chr7q:774124357-774225140
Mmp23b|AMEX60DD201051302.1 at chr8p:464264490-464472275
Mmp21|AMEX60DD201052767.1 at chr8q:562677553-562841533
Mmp28|AMEX60DD102054403.2 at chr9p:396416220-396944082
Mmp28|AMEX60DD102054403.4 at chr9p:396417019-396944082
Mmp28|AMEX60DD301054403.7 at chr9p:396567418-396944082
Mmp28|AMEX60DD301054403.8 at chr9p:396653174-396944082
Mmp11|AMEX60DD201000336.1 at chr10p:120710748-120889660
Mmp11|AMEX60DD301000336.2 at chr10p:120711000-120889660
Immp1l|AMEX60DD201004910.1 at chr11q:427339439-427929566
Immp1l|AMEX60DD201004910.2 at chr11q:427339439-427929566
Immp1l|AMEX60DD301004910.4 at chr11q:427339776-427929566
Loc115460504 [nr]|Mmp17 [hs]|AMEX60DD201007135.2 at chr12q:183313777-183581527
Loc115460504 [nr]|Mmp17 [hs]|AMEX60DD201007135.3 at chr12q:183313945-183581527
Mmp14|AMEX60DD301009336.1 at chr13q:34266398-34380440
Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD201012011.1 at chr14q:309024899-309136701
Loc115090108 [nr]|Mmp24 [hs]|AMEX60DD102012011.2 at chr14q:309024990-309136701
Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD201012011.5 at chr14q:309024990-309136701
Loc115090108 [nr]|Mmp19 [hs]|AMEX60DD102012011.6 at chr14q:309024990-309136701
Dr999_pmt17610 [nr]|Mmp15 [hs]|AMEX60DD301015494.1 at chr1p:930476836-930716207
Immp2l|AMEX60DD201015548.1 at chr1p:954286027-954292665
Mmp2|AMEX60DD301017764.1 at chr1q:389611681-390000101
Mmp2|AMEX60DD102017764.2 at chr1q:389611759-390000101
E2i00_006438 [nr]|Mmp2 [hs]|AMEX60DD102017764.3 at chr1q:389611803-390000101
Mmp9 [nr]|Mmp2 [hs]|AMEX60DD201027684.1 at chr3p:536809048-536829701
Mmp25|AMEX60DD201028039.1 at chr3p:674277241-674440365
Mmp24|AMEX60DD201028519.1 at chr3p:830640463-831382259
Loc115079846 [nr]|Mmp21 [hs]|AMEX60DD301030028.1 at chr3q:614358877-614563086
Loc115079846 [nr]|Mmp21 [hs]|AMEX60DD301030028.2 at chr3q:614359629-614563086
Mmp16|AMEX60DD301040046.1 at chr5q:779673201-780465533
Mmp16|AMEX60DD301040046.2 at chr5q:779673201-780465533
Loc115461210 [nr]|Mmp21 [hs]|AMEX60DD201046376.1 at chr6q:1566829271-1567038020
Loc115091500 [nr]|Mmp13 [hs]|AMEX60DD301049487.3 at chr7q:767994986-768092444
Mmp13|AMEX60DD301049488.1 at chr7q:768348884-768408650

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Col [nr]|Mmp1 [hs]|AMEX60DD201049490.1 at chr7q:769728957-769740418

Loc115091995 [nr]|Mmp13 [hs]|AMEX60DD201049491.1 at chr7q:770037720-770070623

Loc115091995 [nr]|Mmp10 [hs]|AMEX60DD201049491.2 at chr7q:770037720-770069662

Loc115091995 [nr]|Mmp13 [hs]|AMEX60DD201049491.3 at chr7q:770037728-770070623

Mmp3/10b [nr]|Mmp3 [hs]|AMEX60DD301049492.1 at chr7q:770604144-770628728

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049493.1 at chr7q:770954416-770977908

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD102049494.1 at chr7q:771300488-771314054

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD101049496.1 at chr7q:771474186-771498013

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.3 at chr7q:771474227-771615084

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.4 at chr7q:771474227-771615084

Mmp3/10a [nr]|Mmp13 [hs]|AMEX60DD201049496.9 at chr7q:771602594-771615084

Mmpe [nr]|Mmp13 [hs]|AMEX60DD201049502.1 at chr7q:773702091-773764117

Mmp7|AMEX60DD201049503.3 at chr7q:774139369-774225140

Mmp21|AMEX60DD201052767.2 at chr8q:562730284-562841533

Mmp28|AMEX60DD201054403.1 at chr9p:396415902-396944082

Vimentin

https://genome.axolotl-omics.org/cgi-bin/hgTracks?db=ambMex60DD&lastVirtModeType=default&lastVirtModeExtraState=&virtModeType=default&virtMode=0&nonVirtPosition=&position=chr6q%3A172480074%2D172990645&hgsid=69715_5uEAztFlj0qCx310CTXrTpTbcU5J

Sox

Sox2|AMEX60DD201001640.1 at chr10p:771056786-771059999

Sox2|AMEX60DD201001640.2 at chr10p:771056916-771059999

Sox3 [nr]|Sox2 [hs]|AMEX60DD201037395.1 at chr5p:345298485-345300560

Sox19 [nr]|Sox2 [hs]|AMEX60DD301012772.2 at chr1p:86807265-86830424

Sox19 [nr]|Sox2 [hs]|AMEX60DDU001011854.1 at C0108410:20-2879

Loc109080723 [nr]|Sox21 [hs]|AMEX60DD201048613.1 at chr7q:130264218-130267639

Nanog

Nanog [nr]|Nanogp8 [hs]|AMEX60DD201026750.2 at chr3p:136284834-136296773

Nanogp8 [hs]|AMEX60DD301026726.1 at chr3p:126207303-126230971

Loc112993827 [nr]|Nanogp8 [hs]|AMEX60DD201026726.3 at chr3p:126207333-126230971

Nanog [nr]|Nanogp8 [hs]|AMEX60DD103026727.1 at chr3p:126314569-126326644

Nanog [nr]|Nanogp8 [hs]|AMEX60DD201026727.2 at chr3p:126314654-126326644

Nanog [nr]|AMEX60DD201026750.1 at chr3p:136284834-136296773

Nanog [nr]|Nanogp8 [hs]|AMEX60DD201026750.3 at chr3p:136284834-136296773

Klf

Klf12|AMEX60DD201048834.1 at chr7q:306817433-306941849

Loc115061895 [nr]|Klf7 [hs]|AMEX60DD301055067.2 at chr9q:185289613-185298615

Klf2 [nr]|AMEX60DD301014010.1 at chr1p:363123428-363127701

Klf16 [nr]|Klf13 [hs]|AMEX60DD301014425.2 at chr1p:518299066-518402635

Klf6|AMEX60DD301021720.2 at chr2p:752330015-752378770

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

Klf8|AMEX60DD201022566.4 at chr2p:1212076410-1213204249
Klf8|AMEX60DD201022566.5 at chr2p:1212675080-1213204249
Klf8|AMEX60DD201022566.6 at chr2p:1212800997-1213204249
Klf8|AMEX60DD201022566.7 at chr2p:1212841763-1213204249
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.20 at chr2p:1334736936-1334866221
Klf1|AMEX60DD301031962.1 at chr3q:1467331731-1467482191
Klf10|AMEX60DD301040240.2 at chr5q:892823806-892838393
Klf3|AMEX60DD301045679.2 at chr6q:1149031978-1149147917
Pal_glean10020358 [nr]|Klf5 [hs]|AMEX60DD301048840.2 at chr7q:310650331-310684189
Pal_glean10020358 [nr]|Klf5 [hs]|AMEX60DD301048840.3 at chr7q:310650332-310686093
Klf7|AMEX60DD201055067.1 at chr9q:185289368-185544585
Klf7|AMEX60DD201055067.3 at chr9q:185343639-185544585
Klf7|AMEX60DD301055067.4 at chr9q:185349585-185411483
Klf7|AMEX60DD301055067.5 at chr9q:185399837-185544585
Klf13 [nr]|AMEX60DD201004103.1 at chr11q:43081412-43336741
Loc112118413 [nr]|Klf15 [hs]|AMEX60DD301009024.1 at chr13p:203135963-203169830
Loc112118413 [nr]|Klf15 [hs]|AMEX60DD301009024.2 at chr13p:203155127-203169830
Klf16 [nr]|Klf13 [hs]|AMEX60DD201014425.1 at chr1p:518287457-518402635
Cklf|AMEX60DD301015371.1 at chr1p:885673538-886056699
Loc115472666 [nr]|Klf4 [hs]|AMEX60DD301019847.1 at chr1q:1381373911-1381380818
Loc115472666 [nr]|Klf4 [hs]|AMEX60DD201019847.2 at chr1q:1381373930-1381380818
Klf6|AMEX60DD301021720.1 at chr2p:752329950-752341801
Klf8|AMEX60DD201022566.8 at chr2p:1213101702-1213204249
Klf8|AMEX60DD201022566.9 at chr2p:1213101702-1213204249
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.1 at chr2p:1334728745-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.2 at chr2p:1334728745-1334866221
Loc109300690 [nr]|Klf5 [hs]|AMEX60DD301022828.3 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.4 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.5 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.6 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.7 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.8 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.9 at chr2p:1334728755-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.10 at chr2p:1334728756-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.11 at chr2p:1334728766-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.12 at chr2p:1334728775-1334866221
Loc109300690 [nr]|Klf5 [hs]|AMEX60DD301022828.13 at chr2p:1334728775-1334866221
Loc109300690 [nr]|Klf5 [hs]|AMEX60DD201022828.14 at chr2p:1334728775-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.15 at chr2p:1334728775-1334866221
Loc109300690 [nr]|Klf5 [hs]|AMEX60DD201022828.16 at chr2p:1334728775-1334866221
Loc109313665 [nr]|Klf5 [hs]|AMEX60DD201022828.17 at chr2p:1334728784-1334866221
Loc109300690 [nr]|Klf5 [hs]|AMEX60DD201022828.18 at chr2p:1334728784-1334866221

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Loc109313665 [nr]|Klf5 [hs]|AMEX60DD301022828.19 at chr2p:1334728790-1334866221

Klf15|AMEX60DD201023451.1 at chr2q:138944394-139001009

Klf15|AMEX60DD102023451.2 at chr2q:138958680-139001009

Loc115081092 [nr]|Klf1 [hs]|AMEX60DD301031962.2 at chr3q:1467331787-1467481063

Klf11|AMEX60DD201033263.1 at chr4p:411498311-411596909

Klf11|AMEX60DD201033263.2 at chr4p:411501843-411596909

Klf10|AMEX60DD301040240.1 at chr5q:892823799-892839927

Klf9|AMEX60DD201043157.1 at chr6p:1094500273-1094545697

Klf4.s [nr]|Klf4 [hs]|AMEX60DD301044423.1 at chr6q:329294795-329318161

Klf3|AMEX60DD201045679.1 at chr6q:1149031963-1149147917

Klf3|AMEX60DD102045679.3 at chr6q:1149036423-1149147917

Klf5|AMEX60DD201048840.1 at chr7q:310650323-310684189

c-Myc

Myc ファミリー本体

- **C-myc (Myc)**
  - AMEX60DD301040490.1, .2, .3 at chr5q
- **Mycn (N-myc)**
  - AMEX60DD102033164.2 at chr4p
  - AMEX60DD101033177.1 at chr4p
  - AMEX60DD102033177.2 at chr4p
- **Mycl (L-myc)**
  - AMEX60DD301006166.4 at chr11q
  - AMEX60DD201006166.1, .2, .3, .6, .7 など at chr11q

Myc 結合因子 (Binding Partners / Regulators)

- **Mycbp2 (Myc binding protein 2)**
  - 多数（AMEX60DDU001035923.1, AMEX60DD201034118.1, chr7q の複数転写バリアント, etc.）
- **Mycbp (Myc binding protein)**
  - AMEX60DD301005819.1, .2 at chr11q
- **Mycbpap (Myc binding protein-associated protein)**
  - AMEX60DD301030967.1, .2, .3 at chr3q
- **Myct1 (Myc target 1)**
  - AMEX60DD201034385.1, .3 at chr4p
  - AMEX60DD301034385.2

Mycbp2|AMEX60DDU001035923.1 at C0194475:10714-24682

Mycl|AMEX60DD301006166.4 at chr11q:1068532357-1068552477

Mycbp2|AMEX60DD201034118.1 at chr4p:940375188-940375466

Myct1|AMEX60DD201034385.1 at chr4p:1078895019-1079196242

Mycbp2|AMEX60DD201048812.1 at chr7q:286505085-287981527

Mycbp2|AMEX60DD301048812.2 at chr7q:286866803-287981527

Mycbp2|AMEX60DD201048812.3 at chr7q:286866803-287981527

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Mycbp2|AMEX60DD201048813.1 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.2 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.3 at chr7q:288045719-289524879

Mycbp2|AMEX60DD201048813.4 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.5 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.6 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.7 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.8 at chr7q:288045719-289052742

Mycbp2|AMEX60DD201048813.9 at chr7q:288045719-289524879

Llap_12587 [nr]|Mycbp2 [hs]|AMEX60DD301048813.10 at chr7q:288143499-288374124

Mycbp [hs]|AMEX60DD301005819.2 at chr11q:932413022-932437986

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD201006166.1 at chr11q:1068531880-1068552477

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD301006166.8 at chr11q:1068535410-1068552477

Mycbpap|AMEX60DD301030967.2 at chr3q:1056714933-1057991505

Mycn|AMEX60DD102033164.2 at chr4p:354934552-354944744

Mycn|AMEX60DD101033177.1 at chr4p:359712079-359722323

Mycn|AMEX60DD102033177.2 at chr4p:359712142-359720078

Mycbp.l [nr]|Mycbp [hs]|AMEX60DD301005819.1 at chr11q:932412902-932437986

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD201006166.2 at chr11q:1068532072-1068552477

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD201006166.3 at chr11q:1068532328-1068552477

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD301006166.5 at chr11q:1068532715-1068552477

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD201006166.6 at chr11q:1068533032-1068552477

Ab205_0023440 [nr]|Mycl [hs]|AMEX60DD201006166.7 at chr11q:1068533684-1068552477

Mycbpap|AMEX60DD301030967.1 at chr3q:1056714881-1057991505

Loc115090937 [nr]|Mycbpap [hs]|AMEX60DD301030967.3 at chr3q:1056714948-1057245109

Myct1|AMEX60DD301034385.2 at chr4p:1078931688-1079196242

Myct1|AMEX60DD201034385.3 at chr4p:1079012941-1079196242

C-myc [nr]|Myc [hs]|AMEX60DD301040490.1 at chr5q:1041514610-1041550002

C-myc [nr]|Myc [hs]|AMEX60DD301040490.2 at chr5q:1041535472-1041550002

C-myc [nr]|Myc [hs]|AMEX60DD201040490.3 at chr5q:1041543523-1041550002

Mycbp2|AMEX60DD201048811.1 at chr7q:286504487-286506248

Prod1

Prod1 [nr]|AMEX60DD301028208.1 at chr3p:722854759-722913067

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

総説: 魚類・両生類におけるビタミン類への応答性とその効果

著: 杉山 遥 (ハルラボ); 横江 誼衡 (ハルラボ, 東京大学, 秀明大学, SciMountechnology),
F.N. 三太子 (ハルラボ), 竜岡 龍騎 (トラフラボ), 阿保郎 太朗 (ハルラボファーム)

要旨: メキシコサラマンダー (*Ambystoma mexicanum*, *Axolotl*, 以下、アホロートル) は高い再生能力を持つ一方で、野生個体は絶滅危惧種に指定されており、保全と基礎研究の両立が課題である。我々NPO研究チーム「ハルラボ」は2020年に発足し、飼育技術の開発と分子・生理学的研究を統合する活動を行ってきた。これまでの取り組みは、(1) 原産地ソチミルコ湖を中心とした水質履歴の解析、(2) 飼育法の改良（低塩分添加飼育法や過密条件下での濾過システムの導入）、(3) 栄養因子（ビタミン、アミノ酸、脂肪酸など）の効果検証に重点を置いて展開してきた。

(3) の検討においては、生体の成長にビタミンが不可欠であることから、マルチビタミン粉末を餌に添加して与え、成長率への影響を調査した。さらに、肉食性の高いアホロートルが自然界で摂取する可能性の高い魚類や昆虫類にも注目し、モデル生物であるメダカ (*Oryzias latipes*)、フタホシコオロギ (*Gryllus bimaculatus*)、キイロショウジョウバエ (*Drosophila melanogaster*) においても、ビタミン摂取が成長率に与える影響を定量的に評価した。加えて、ビタミンを補給することで栄養的に強化されたこれらの生物をアホロートルに活餌として与えた際の効果も同様に検証した。その結果、ビタミン補給により成長速度が有意に上昇し、腎臓・腸管・骨格に関連する遺伝子発現の変化がRNAレベルで確認された。

本研究では、ビタミンを摂取した活餌に対し、それぞれ「メダカ－メダミン (Medamin)」「フタホシコオロギ－フタミン (Futamin)」「ショウジョウバエ－ドロソミン (Drosomin)」、更にそれらを給餌して強化されたアホロートルを「ウパミン (Upamin)」と命名し、新たな栄養介入モデルとして確立した。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Review: Vitamin Responsiveness and Effects in Aquatic Vertebrates (Fish and Amphibians)

Authors: Sugiyama *Axolotl* Haruka[1]; Yoshichika Yokoe[1,2,3,4];
F.N. Santako[1], Tatsuoka Ryuki[5], Axolotl Taro[6]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo,
[3] Shumei Univ, [4] SciMountechnology, [5] *Ambys* Lab, [6] Hal lab farm

Abstract: The Mexican salamander (*Ambystoma mexicanum*, axolotl) is a model for regeneration but critically endangered in the wild. Our non-profit team HAL lab (est. 2020) integrates husbandry innovations with molecular and physiological studies. We analyzed historical water quality in Lake Xochimilco, developed improved rearing methods (low-salinity culture, filtration for high-density rearing), and evaluated nutritional factors. Focusing on vitamins, we supplemented diets with multivitamin powder and assessed effects on axolotls, as well as on model prey species—medaka (*Oryzias latipes*), crickets (*Gryllus bimaculatus*), and fruit flies (*Drosophila melanogaster*). Vitamin-fed prey were then used as live diets for axolotls. Supplementation significantly enhanced growth and induced RNA-level changes in kidney, intestine, and skeletal genes. We established enriched prey lines—“Medamin,” “Futamin,” and “Drosomin”—and defined vitamin-fed axolotls as “Upamin,” thereby creating a reproducible nutritional intervention model that integrates conservation and experimental biology.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

魚類・両生類におけるビタミン類への応答性とその効果

Vitamin Responsiveness and Effects
in Aquatic Vertebrates (Fish and Amphibians)

杉山 遥 (ハルラボ); ○**横江 誼衡** (ハルラボ, 東京大学, 秀明大学, SciMountechnology),
F.N. 三太子 (ハルラボ), 竜岡 龍騎 (トラフラボ), 阿保郎 太朗 (ハルラボファーム)

E-mail adress sugiyama.haruka.axoltl@gmail.com **Twitter URL** https://twitter.com/lab_new2

ウーパールーパー (アホロートル, *Ambystoma mexicanum*)
を含む**有尾両生類**や**魚類**の成長・成育に重要な因子は何か？

→ **ビタミン類に注目！**

定義: ビタミン（ヴィタミン、英語: vitamin）とは？

生物の生存・生育に微量に必要な栄養素のうち、
その**生体内で十分な量を合成できない炭水化物・タンパク質・脂質以外の有機化合物**の総称.

例 **カルシウムとビタミンDの併用**
→ 強い骨格を作る (爬虫類等の硬骨)

爬虫類・両生類用栄養補給助剤
«リンを含まない天然カルシウム»
爬虫類 トカゲ カルシウム ビタミン エサ /サプリメント/添加剤/ GEX エキゾテラ カルシウム40g



【保証成分】

水分	12%以下
カルシウム	35～37%以下
塩分	0.06～0.07%以下
カリウム	0.04%以上
マグネシウム	0.03%以上
硫黄	0.04%以上
鉄	240ppm以上
亜鉛	9ppm以上
マンガン	5ppm以上
銅	2ppm以上



Amazon | エキゾテラ GEX EXOTERRA マルチビタミン 30g PT1860 不足しがちなビタミンやミネラルなど10種類以上を含有したパウダーサプリ | EXO TERRA | ヘルスケア 通販

他、一部の魚用人工飼料等にも添加されている例あり。

ビタミンを効率的に摂取させるには？



脂溶性ビタミン
・ビタミンA(レチノール/βカロテン)
・ビタミンD(カルシフェロール)
・ビタミンE(トコフェロール)
水溶性ビタミン
・ビタミンB1(チアミン)
・ビタミンB2(リボフラビン)
・ビタミンB5(パントテン酸)
・ビタミンB6(ピリドキシン)
・ビタミンB12(コバラミン)
・ビタミンC(アスコルビン酸)
・ビタミンH(ビオチン)
・ナイアシン(ニコチン酸)
・葉酸(ホラシン)


脂肪分を含む餌と併せて経口投与

水分 (飼育水/飲用水) を介して経口投与

生餌 (活餌) を介した経口投与


直接接種

ビタミン類摂取の恩恵
・高タンパク (筋肉)
・高脂肪 (脂質)
・成長促進 など
・免疫力強化 など



ビタミン類摂取の恩恵
・高タンパク (筋肉)
・高脂肪 (脂質)
・成長促進 など

ビタミンを摂取させた活餌の系統




ビタミンを摂取したアホロートルの系統

・成長率↑
・高妊性 (子孫数)↑
・免疫力↑


MA-M5 ウパミン




アホロートルへのマルチビタミン給餌効果




MB-M1 メダミンとその性質




MC-M2 フタミンとその性質




MD-M3 ドロソミンとその性質




MB-M1 メダミンの経口投与効果



MC-M2 フタミンの経口投与効果



MD-M3 ドロソミンの経口投与効果



CC BY NC ND

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl



第48回日本分子生物学会年会

特定非営利活動法人日本分子生物学会

演題投稿 - 第48回日本分子生物学会年会

[3P-958]

アホロートルの尾部形状が遊泳効率に及ぼす影響の数値流体シミュレーション

○横江 誼衡[1,2,3]、杉山 遥[2] (1. 東京大学、2. ハルラボ、3. サイマンテクノロジー)

要旨（Abstract）

ハルラボでは、絶滅危惧種であるメキシコサラマンダー（*Ambystoma mexicanum*、以下、アホロートル）の保全および生態理解を目的とした基礎研究を行っている。その一環として、本研究ではアホロートルが本来有していた遊泳機能に着目し、特に尾部形状が水中での推進効率に与える影響について数値流体シミュレーションを用いて検討した。 アホロートルは幼形成熟を特徴とし、生涯を水中で過ごすため、尾部の形状は遊泳において中心的な役割を担っていると考えられる。しかしながら、尾部の長さや形状が遊泳効率に与える定量的な影響については、これまで十分に検討されていない。本研究では、尾部を動的な波形構造としてモデル化し、Python ベースの流体解析ツールを用いてシミュレーションを実施することで、形状ごとの推進力および効率を比較した。
本発表では、この現状報告と得られた知見について発表する。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

3P-958

アホロートルの尾部形状が遊泳効率に及ぼす影響の数値流体シミュレーション

A Numerical Fluid Dynamics Study on the Influence of *Axolotl* Tail Morphology on Swimming Efficiency

○横江 誼衡[1,2,3], 杉山 遥[1]


([1] 東京大学, [2] ハルラボ・メキシコサンショウウオ生体研究部, [3] サイマンテクノロジー)

E-mail adress sugiyama.haruka.axoltl@gmail.com **Twitter URL** https://twitter.com/lab_new2


1). *Ambystoma mexicanum* を含む有尾両生類が尿尾を持つ機能的な意義については未解明な部分が多い



疑問: 何故長い尾部を持つのか？(必要?)
→ 流水への抵抗性などの役割あり？
→ カエル等の無尾両生類との違いは？
→ 実は尾部は無くても問題ないのでは？
→ でも彼らは泳ぐときにヒレを振っている。




Scale bars: 20mm




調査課題: 尾部の長さ・形状は遊泳能力を発揮する上で重要か？
(流体力学的な観点)
↓
尾部を損傷してしまった個体で運動能力を観察・定量した。
＋行動学的シュミレーションを実施


2). 尾部を損傷した *Ambystoma mexicanum* の遊泳能力は著しく低下する




6). *Ambystoma mexicanum* は水流を側線器官にて感知している



7). まとめ: *Ambystoma mexicanum* の遊泳メカニズム





3). *Ambystoma mexicanum* は尾部を損傷することで前方向へ姿勢を保って泳ぐことができなくなる



4). 尾部損傷による遊泳能力低下は炎症によるものではない

筋肉が炎症し、その刺激で動きが遅くなってしまう?






5). 損傷した尾部が再生することで遊泳能力が回復する

尾部再生 → 遊泳能力は回復する



Scale bars: 20mm




8). 最新報告: *Ambystoma mexicanum* の遊泳シュミレーション

目的
ウーパールーパー(または任意の水生生物形状)を対象に、流体シミュレーション(CFD)を用いて、泳行時に生じる周囲の流れ場の特性を明らかにすることを目的とした。特に、尾びれや体形が流れに与える影響、速度分布・渦構造・圧力場などを可視化および定量化し、生物力学的・流体力学的観点から考察を行う。

手法
数値流体力学(CFD: Computational Fluid Dynamics)により、非圧縮性 Navier–Stokes 方程式を数値的に解き、流体の速度場を求めた。対象物体の形状は、泳行時のウーパールーパーの輪郭をビットマップ画像として取得し、白黒の2値画像により固体領域と流体領域を定義した。設定した流体領域に対して時間発展計算を行い、速度分布の変化や渦生成の様子を解析し、流れの特徴について考察した。今回のCFDは、すべてFlowsquare+を用いて行った。

シミュレーション条件:
流入口の圧力 10^5Pa, 流入口の流速0.3 m/s, 流体密度 1000kg/m^3, 動粘度1.0 × 10^{-3} Pa·s

結果
定常状態でエラ付近に著しく流体の速度が小さい領域があることがわかった。このことから、ウーパールーパーは遊泳の際、頭部の付け根あたりに圧力を受けていることが予想される。また、魚類には遊泳の際、このように局所的に体にむかって圧力がかかることが遊泳動作を除くとみられないことから、明らかにウーパールーパーの頭部の形状は遊泳という観点からは不利に働いていることが推察される。



今後の展望
・尾の有無の考慮、3次元シミュレーションへの拡張、尾の泳動作の効果を反映、各種パラメータの検討

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。



第48回日本分子生物学会年会
特定非営利活動法人日本分子生物学会

演題投稿 - 第48回日本分子生物学会年会

[3P-980]

栄養補填によって回復しない成長不全を示すアホロートル個体の要因調査

○杉山 遥[1]、枝豆 やみん[1]、あすみ 悠[1]、F.N. 三太子[1]、竜岡 龍騎[2] (1. ハルラボ、2. トラフサンショウウオラボ)

要旨（Abstract）

ハルラボでは、絶滅危惧種であるメキシコサラマンダー（*Ambystoma mexicanum*，以下、アホロートル）の保全に向けた研究活動を行っている。

活動の一つとして、彼らが繁栄していた頃の生態を明らかにすることを目的に、ラボレベルでの予想生育環境モデルを作製し、生体の成長率や生理学的解析、行動解析等を実施している。 過去の検討結果から、一部のアホロートル CB (Captive Bred，自家繁殖）系統の交雑の中で、成長不全 (Poor Growth, PG) となる F1 個体を産み出す系統が散見された。

これらの PG 個体における成長制御や代謝制御に関連する因子のｍRNA オミクス解析により、糖質、脂質、ビタミンの代謝障害を産み出す欠損変異がそれぞれ確認された。

また、これらの代謝障害を持った個体の表現型は、餌の組成に不足している栄養成分の含有量を大幅に増やすことである程度回復できることが明らかとなった。(参考文献 [1])

しかしながら、PG 個体うち、栄養の補填では成長率が回復しないが個体も見られたことから、胴体のみ成長が遅れてしまう矮小変異系統 (Dwarf; SB ショートボディ/ST ショートテイル: 参考文献[2]) と併せて、成長不全の要因についての共通点を調査した。

調査にあたり、これらの成長・生育不全の要因について、哺乳類である程度解明が進んでいるⅡ型コラーゲン異常症 や 小人病 i.e. Dwarf disease (DW) のような成長不全疾患に注目し、それらの

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

疾患における原因因子として報告が見られるコラーゲン (Col1, Col2) や骨/軟骨成長・形成・分化に関わる RUNX family および Sox family などに関して、アホロートルにおける役割やその発現動態について調査・研究を実施した。

qPCR による幅広い関連因子の調査・探索などを行った結果、栄養状態の改善によって完全には改善できない成長不全において、Col1, Col2 および RUNX family 群の発現低下による軟骨伸長不全や、慢性的な睡眠不全によるストレス応答の持続による代謝異常・成長抑制が生じている可能性が示唆された。また、これらの PG の表現型は、マウス DW 変異と同様に成長因子である FGF family および その特異的受容体である FGFR family の投与によって回復できることも明らかとなった。(参考文献 [3])

本発表では、これまでの調査結果によって得られた知見について報告する。

参考文献

[1]. 杉山 遥 ほか,【研究報告】ビタミンで育つ！メダカとウーパールーパー！ (はるらぼ会報誌_ウーパールーパーだより vol 5.0, 2024 年 3 月)
[2]. Stéphanie Garcia et al., Postnatal soluble FGFR3 therapy rescues achondroplasia symptoms and restores bone growth in mice. (Sci. Transl. Med., 2013 Sep.)
[3]. 杉山 遥 ほか,【研究報告】育たないウーパールーパーの秘密を解明！ (はるらぼ会報誌_ウーパールーパーだより vol 6.0, 2024 年 6 月)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

栄養補填によって回復しない成長不全を示すアホロートル個体の要因調査

3P-980

Factorial investigation of Poor-Growth-*Axolotl* that does not recover with nutritional supplementation

○杉山 遥[1]、枝豆 やみん[1]、あすみ 悠[1]、F.N. 三太子[1]、竜岡 龍騎[2]

(1. ハルラボ、2. トラフサンショウウオラボ)


E-mail adress sugiyama.haruka.axoltl@gmail.com **Twitter URL** https://twitter.com/lab_new2

1). 課題: *Ambystoma mexicanum* 成長不全個体の発生要因の特定




過去結果: マルチビタミンを餌に混ぜた経口補填で、
成長不全が回復できない系統 (＝矮小個体, Poor Growth: PG) あり.
↓
PG 系統 の一部で、 Col1, Col2, Sox, Runx family の発現低下が見られた.
しかし、**栄養状態の改善 (餌条件の変更) だけではレスキューできない**.
(下流のFGF signal 活性化では回復効果あり)

2). 成長不全の *Ambystoma mexicanum* 個体へ成長因子を補填することで表現型を回復できる




成長不全個体・・・MT1, 2, 4, 5, 6, 8, 9
過剰成長 (肥大) 個体・・・MT3, 7, 11


3). *Ambystoma mexicanum* の成長制御に関わる因子のスクリーニング




4). ANK1, ANKHD1 が *Ambystoma mexicanum* の成長制御に与える影響1





5). ANK1, ANKHD1 が *Ambystoma mexicanum* の成長制御に与える影響2




6). ANK1, ANKHD1 発現低下による成長不全表現型のレスキュー検討









※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。



第48回日本分子生物学会年会

特定非営利活動法人日本分子生物学会

演題投稿 - 第48回日本分子生物学会年会

[3P-981]

水道水中の残留塩素を利用した、飼育下のアホロートルの感染症予防法の開発。

○F.N. 三太子[1,2]、横江 誼衡[1,3,4]、杉山 遥[1] (1. ハルラボ・メキシコサンショウウオ生体研究部、2. アホロートルレスキュー、3. サイマンテクノロジー、4. 東京大学・理・物理・宇宙線研)

要旨（Abstract）

メキシコ原産のサンショウウオであるメキシコサラマンダー（*Ambystoma mexicanum*，以下、アホロートル）は、現在絶滅の危機に瀕している。
アホロートルを取り巻く環境や水質悪化などが主な原因とされているが、アホロートルは飼育下の個体であっても免疫が脆弱であることが知られており、アホロートルにとって最適な成育環境を考える上で感染症や免疫機構は非常に重要となってくる。

我々は、メキシコにおけるアホロートルの保全を目指し、クラウドファンディングによる保全活動資金の確保や様々な飼育成体における飼育治療相談活動（アホロートル・レスキュー）などの活動を実施しており、彼らにとって最適な環境を確保する上でも疾患の原因や最適な対処法について広く理解する必要がある。
我々のこれまでの調査結果においても、水質（硬度や pH）の変化に対する抵抗性の高さに反して、水質悪化に伴って発生する耐塩性を示す腸炎ビブリオ（*Vibrio parahaemolyticus*）などの病原微生物自体への抵抗性が低いことが示唆されている。

また、アホロートルは特に真菌類や耐塩性細菌等からの影響を受けやすく、対処が遅れることで早ければ数週間の間に亡くなってしまう可能性があることが明らかとなっている。（第 46 回日本分子生物学会年会_神戸ポートアイランド（2023）にて報告。）

この他にも、これまでの活動における症例とそれらの原因となる微生物について、可能な限りを調査しリスト化することで、より効果的にアホロートルの疾患への理解と適切な対処を可能となることを目指している。（第 47 回日本分子生物学会年会_マリンメッセ福岡（2024）にて報告。 書籍による発表準備中。）

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。



更に我々は、未解明な部分が多い感染症機序について解明する糸口を掴む手段の 1 つとして、血球マーカー および 抗菌ペプチドについてスクリーニングを実施し、複数の因子の同定を行っている。 (中国四国地区_生物系三学会_合同大会 (2025) にて一部報告。)

これに加え、これまでの知見から、アホロートルの免疫は上皮および粘膜および表面に発現した抗菌ペプチドへの依存度が高く、真菌類や細菌類へ感染後、一度体内まで感染が進行してしまうと抗いきれずに亡くなってしまうことが想定される。

これについては、我々の調査により、炎症マーカーの急激な増加によって、症状が表面化する頃には既に病原体が体内へ大量に侵入していることが示唆されている。

このような知見を踏まえ、従来とは異なる切り口の一つの考えとして、”感染後から致命的な段階に移行するまでの短期間、かつ耐えられる体力が残存している間に、オキソリン酸やポピドンヨードを用いた薬浴を実施するよりも、常に無菌的な環境に生体を置くことの方が、より安全に安定に生体を飼育・維持することが可能なのではないだろうか？” という内容が浮かんだ。

これを解決する策の一つの可能性として、”**水道水に残留する 0.6mg/L (0.6ppm) のカルキ (次亜塩素酸イオン) をそのまま敢えて中和せずに利用する**” という手法を検討し、数年間に渡り感染症に罹患しない状態で安定的に飼育可能であるという一定の成果を得た。

本調査報告と知見の共有により、野生種の適切な保全や飼育個体の健康管理に関する更なる手がかりとなることを切に望む。

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl



水道水中の残留塩素を利用した飼育下のアホロートルの感染症予防法の開発

3P-981

Development of a Residual-Chlorine–Based Infection Control Method for Captive *Axolotls*.

○ F.N. 三太子[1,2]、横江 誼衡[1,3,4]、杉山 遥[1]

(**所属**: 1. **ハルラボ・メキシコサンショウウオ生体研究部**、2. **アホロートルレスキュー**、3. **サイマンテクノロジー**、4. **東京大学・理・物理・宇宙線研**)

E-mail adress sugiyama.haruka.axoltl@gmail.com **Twitter URL** https://twitter.com/lab_new2

CC BY NC ND

アホロートル ≒ ウーパールーパー とは？

メキシコサラマンダー (*Ambystoma mexicanum*)
の国内流通個体です (一部、他種と混血の説あり)

原産地: **メキシコ** (**ソチミルコ湖**)
最大体長: 20〜25cm
幼形成熟 (**ネオテニー**) **の形態を取れる**
寿命は20**年以上** (**変態・上陸すると**10**年程度に。。。**)
強み: **少々の水質変化はへっちゃら**
弱点: **細菌感染**, **カビ**(**免疫反応がアフリカツメガエルの半分以下**)
※ **現在は絶滅の危機** (**原因**: **都市開発や埋め立てによる生息地減少**)

ハルラボとは？
アホロートルを救いたい！ → 2020**年当ラボによる調査・研究活動を開始!**
他、睡眠や共生の仕組みなど諸々研究内容を報告中。

マーブル (野生型体色)



(野生種を生育し、ある程度固定化した品種。一部、交雑種も存在する.)

リューシスティック (一部色素欠乏)



(*endothelin3*: *edn3* (Chr3p 短腕) 欠損変異ホモ; 上皮におけるユーメラニン欠乏)

ブラック (メラニン過剰)



(*Leukocyte Tyrosine Kinase*: *Ltk* (Chr14q 長腕) 欠損変異ホモ; 上皮におけるユーメラニンの過剰分泌・蓄積)

アルビノ系統 (*tyrosinase*: *tyr* (Chr7q 長腕) 欠損変異ホモ; 全身のユーメラニン欠乏)

アルビノ



ゴールデン



イエロー



遺伝子欠損の程度・フェオメラニン合成への影響の違い、他の体色変異 等と併せることで体色に違いが見られる.

(杉山 ほか. 2022-2023)

参考: 水槽内に存在する微生物一覧 (アホロートルの病気の原因たち)

性質・種類	学名	属・分類	備考
グラム陰性細菌	*A.hydrophila, A.sobria*	エロモナス類	感染症の原因菌
	F. columnare	カラムナリス類	感染症の原因菌
	N. europaea	ニトロソモナス類	アンモニア硝化菌
	N. winogradskyi	ニトロバクター類	亜硝酸酸化菌
	S. enterica	サルモネラ類	腸内細菌
	E. coli	大腸菌	腸内細菌
	Vibrio vulnificus	**腸炎ビブリオ**	**吐き戻しの原因 (好塩細菌)**
	Campylobacter jejuni	**カンピロバクター類**	**肉類に付着 (耐塩細菌)**
	Staphylococcus aureus	**黄色ブドウ球菌類**	**皮膚の化膿・嘔吐など (耐塩細菌)**
	Streptococcus iniae	**レンサ球菌**	**眼球突出、腎臓や脾臓の腫れ、心外膜炎など**
グラム陽性細菌	*L. plantarum*	乳酸菌	腸内細菌
	B.subtilis	枯草菌類	連過バクテリア
藻類 (水苔)	*C. pediculus*	ッコネイス属	珪藻類
	T. lacustris	テトラスポラ属	緑藻類
	M. willeana	ミクロスポラ属	糸状緑藻類
真菌 (カビ)	*S. cerevisiae*	出芽酵母	パン酵母
	S. pombe	分裂酵母	ビール酵母
	Rhodotorula	赤色酵母	赤色酵母
	S. parasitica	ミズカビ属	
	A. prolifera	ワタカビ属 (卵菌類)	水質浄化, 増えすぎると水カビ病の原因
	A. proliferoides		
	A. racemosa		
	A. cochlioides	アファノマイセス属 (糸状菌類)	

(杉山 ほか. 2022-2023)

(A) 健康な頃 (アカムシや固形餌で飼育)




(B) 衛生的に良くなかったと思われるマグロを摂食した場合 (数か月経過後)



水道水 (残留塩素 0.6ppm≧) でアホロートルを薬浴治療の代替＋飼育も可能と判明！(長期試験中)

上清の中に含まれる相対微生物量 (底材_滅菌の効果)



水質関連データ (底材なし, ベアタンク)		カルキを抜かない水道水 (一部地域を除く全国平均)	カルキ抜き水 (チオ硫酸Na 添加後)	食べ残しによる汚染水 (粒餌食べ残し後24hr)	食べ残しによる汚染水 (粒餌食べ残し後48hr)
水温	(℃)	15〜20			
硝酸イオン (NO_3^-)	(ppm)	0	0	10〜40	50〜100
亜硝酸イオン (NO_2^-)		0	0	0.5〜1.0	1.0〜5.0
残留塩素 (Cl^-)		0.4〜0.6	0	0	0
総硬度 (GH)		25〜100	25〜100	25〜100	25〜100
炭酸塩硬度 (KH)		30〜55	30〜55	30〜55	30〜55
pH		6.8〜7.5	7.2〜7.5	6.0〜6.8	6.0〜6.4

※ 硝酸イオン (NO_3^-): 水中のアンモニアが水中の硝化細菌によってある程度無害化された状態 (ただし強酸性)。
※ 亜硝酸イオン (NO_2^-): 水中のアンモニアが亜硝酸菌によって作られた物質。水槽生物の酸欠の原因になり有害。
※ 底材やフィルターが無い環境における 亜硝酸イオン、硝酸イオンの発生源は、糞中に含まれる腸内細菌の影響などが挙げられる。
※ 硬度 (水に含有する Ca^{2+}, マグネシウムイオン Mg^{2+} の濃度) : mg/L = ppm = dH/17.8
※ 水温がこれより低くなってもCl-は0.6ppmを基本超えない。
※ 水の硬度の目安: **軟水 60ppm≧**, **中硬水 60〜120ppm**, **硬水 ≧120ppm**, **超硬水 ≧180ppm**
※ 日本の水道水は基本的に軟水。地域の水道局の情報から、水の組成を確認可能。
※ 環境省による残留塩素濃度: **基準値 0.6mg/L ≒ 0.6ppm (平成20年 4月 1 日施行)**
参照: https://www.env.go.jp/water/water_supply/kijun/index.html
※ 総硬度 (GH): 水中のカルシウムイオン $Ca2+$, マグネシウムイオン $Mg2+$ の濃度 (総量)
※ 炭酸塩硬度 (KH): 水中に含まれる炭酸塩 [例: 炭酸水素Ca. 重炭酸Ca $Ca(HCO_3)_2$]. 炭酸ナトリウム Na_2CO_3] の濃度
・・・二酸化炭素 (CO_2) が水に溶けている量が多いほど、$Ca(HCO_3)_2$ が水中に多く生成される。
KHが高くなる程、水がアルカリ性に傾き、pHの急激な低下を抑制する。


(杉山 ほか. 2022-2025)

エロモナス菌 *Aeromonas hydrophila / sobria*




Aeromonas sobria: Introduction, Morphology, Pathogenicity, Lab (medicallabnotes.com)

カラムナリス菌 *Flavobacterium columnare*




Flavobacterium columnare - Alchetron, the free social encyclopedia / Flavobacterium columnare [IMAGE] | EurekAlert! Science News Releases

腸炎ビブリオ *Vibrio parahaemolyticus*




Vibrio parahaemolyticus | DSM 25722, NCCP 13713 | BacDive(D17271 (dsmz.de) / Staphylococcus Aureus : Lab Diagnosis and Diseases - Medchrome

カルキ (次亜塩素酸カルシウム) を抜かない水道水によるウイルス・細菌・カビの殺菌





	4月			5月			6月		
	最低値	平均±誤差	最大値	最低値	平均±誤差	最大値	最低値	平均±誤差	最大値
気温 (℃)	17.2	19.42±0.31	21.7	14.3	16.86±0.28	20.9	18.9	23.27±0.47	26.9
水温 (℃)	14	16.13±0.36	18.8	14.9	18.59±0.36	22.3	15.7	21.79±0.59	26.4
残留塩素 (mg/L)	0.3	0.46±0.02	0.7	0.2	0.47±0.03	0.8	0.2	0.47±0.03	0.8
pH	6.8	7.22±0.03	7.6	6.7	7.25±0.04	7.6	6.7	7.20±0.04	7.7



アホロートルの成育率 (2L 容器, 底材あり: day0〜day300)










写真: ベアタンク、水道水飼育で4年間元気に育った個体 [R. A. (仮) 様]










CC BY NC ND

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

Pesticides and Living Organisms



Author: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe [1, 2, 3, 4]

1. HAL lab Axolotl, Ecology Lab of *Ambystoma mexicanum*,
2. ICRR, The Univ. of Tokyo, 3. Scimoun Technology, 4. Shumei Univ.

【Review】Investigation Report on Tolerance to Organic Solvents and Pesticide Components Present in the Rearing Water of Axolotl, *Ambystoma mexicanum*

Author: Sugiyama Axolotl Haruka (HAL lab Axolotl, Ecology Lab of *Ambystoma mexicanum*), Yoshichika Yokoe (ICRR, The Univ. of Tokyo / Scimoun Technology / Shumei Univ.), F.N. Santako (HAL lab Axolotl rescue), Ya!min Beans (Ecology Lab of the Freshwater Fish), Ryuki Tatsuoka (Ambys lab), Axolotl Taro (HAL lab Ferm)

【Background】 The Mexican salamander (*Ambystoma mexicanum*), commonly known in captivity as the axolotl, is widely kept as a companion species. However, wild populations are currently facing a severe risk of extinction and are listed as CR (Critically Endangered) on the IUCN Red List. One major cause of this decline is water pollution in its natural habitat, Lake Xochimilco in Mexico, resulting from tourism development and urban expansion. Domestic wastewater and industrial effluents flow into the lake, leading to progressive deterioration of water quality. In addition, the traditional chinampa farming system—agriculture conducted on floating artificial islands—is being revived in surrounding areas, raising concerns about increased contamination of the water by agricultural chemicals such as pesticides.

In light of these circumstances, we quantitatively evaluated the responses of captive axolotls to chemical substances present in rearing water, with the ultimate aim of contributing to the conservation of wild populations.

【Methods】 Various chemical agents were added to axolotl rearing water at multiple concentration levels, including commonly used aquarium conditioners (e.g., sodium thiosulfate contained in dechlorinators, pH adjusters, and photosynthetic bacterial preparations), surfactants that may enter natural habitats as domestic wastewater from soaps and detergents, several organic solvents potentially involved in industrial water pollution [ethanol (water-soluble), methanol (water-soluble), isopropanol (IPA, water-soluble), acetone (water-soluble), and dichloromethane (water-soluble)], and pesticide components (pyrethroid compounds).

Phenotypic effects associated with chemical exposure—such as inflammation, mucus production due to chemical injury, and lethality (LD_{50})—were quantitatively and statistically assessed. Experimental procedures included direct addition of chemicals to rearing tanks followed by individual observation and measurement. In addition, interviews with numerous axolotl keepers were conducted to collect information enabling statistical evaluation of exposure effects to a wide range of chemicals.

Furthermore, homologs of genes involved in drug efflux transport in mammals and fish were identified using databases such as GenBank and BLAST. mRNA was extracted from tissue samples of exposed individuals, and gene expression changes before and after exposure were analyzed by qRT-PCR,

allowing integration of molecular data with phenotypic observations for a more comprehensive evaluation.

【Results and Discussion】 Even commonly used aquarium chemicals caused abnormal phenotypes in axolotls unless applied at approximately one-thousandth of the recommended dosage for typical ornamental freshwater fish (with some exceptions). In severe cases, individuals died within 24 hours after treatment. For most organic solvents, lethality increased sharply between 1,000 ppm (0.1%) and 1.0%. These findings, together with information obtained from keeper interviews, suggest that chemical exposure can pose an immediate threat to axolotl survival.

Although expression of homologous drug transporter genes was detected in some cases, the rate of chemical elimination appeared to be markedly slower than the rate of uptake through epithelial tissues. This imbalance may contribute to the high sensitivity of axolotls to chemical exposure.

Overall, these results provide important new insights into how environmental contamination can severely threaten the survival of axolotls.

【References】

Ambudkar, S. V., Dey, S., Hrycyna, C. A., Ramachandra, M., Pastan, I., & Gottesman, M. M. (1999). Biochemical, cellular, and pharmacological aspects of the multidrug transporter P-glycoprotein. *Annual Review of Pharmacology and Toxicology*, 39, 361–398.

Bard, S. M. (2000). Multixenobiotic resistance as a cellular defense mechanism in aquatic organisms. *Aquatic Toxicology*, 48(4), 357–389.

Birge, W. J., Westerman, A. G., & Spromberg, J. A. (2000). Comparative toxicology and risk assessment of amphibians. In D. W. Sparling, G. Linder, & C. A. Bishop (Eds.), *Ecotoxicology of Amphibians and Reptiles* (pp. 727–791). SETAC Press.

Blaustein, A. R., & Kiesecker, J. M. (2002). Complexity in conservation: Lessons from the global decline of amphibian populations. *Nature*, 415, 681–684.

Bradbury, S. P., & Coats, J. R. (1989). Comparative toxicology of the pyrethroid insecticides. *Reviews of Environmental Contamination and Toxicology*, 108, 133–177.

Brühl, C. A., Schmidt, T., Pieper, S., & Alscher, A. (2013). Terrestrial pesticide exposure of amphibians: An underestimated cause of global decline? *Scientific Reports*, 3, 1135.

Duellman, W. E., & Trueb, L. (1994). *Biology of Amphibians*. Johns Hopkins University Press.

González-Pozo, A., et al. (2016). Urban agriculture and environmental sustainability in Xochimilco, Mexico City. *Environmental Management*, 58, 101–115.

Lewis, M. A. (1991). Chronic and acute toxicities of surfactants and detergent builders to aquatic organisms: A review and risk assessment. *Water Research*, 25(1), 101–113.

Lipnick, R. L. (1991). Outliers: Their origin and use in the classification of molecular mechanisms of toxicity. *Science of the Total Environment*, 109–110, 131–153.

Mann, R. M., Hyne, R. V., Choung, C. B., & Wilson, S. P. (2009). Amphibians and agricultural chemicals:

Review of the risks in a complex environment. *Environmental Pollution*, 157(11), 2903–2927.

Maund, S. J., Hamer, M. J., Lane, M. C. G., et al. (2002). Partitioning, bioavailability, and toxicity of the pyrethroid insecticide cypermethrin in aquatic environments. *Environmental Toxicology and Chemistry*, 21(1), 9–15.

McCarty, L. S., & Mackay, D. (1993). Enhancing ecotoxicological modeling and assessment: Narcotic mode of action. *Environmental Toxicology and Chemistry*, 12(4), 629–643.

Quaranta, A., Bellantuono, V., Cassano, G., & Lippe, C. (2009). Why amphibians are more sensitive than mammals to xenobiotics. *PLoS ONE*, 4(11), e7699.

Smital, T., Sauerborn, R., & Hackenberger, B. K. (2004). Inducibility of the P-glycoprotein transport activity in the marine mussel *Mytilus galloprovincialis*. *Aquatic Toxicology*, 70(2), 131–141.

Sparling, D. W., Linder, G., Bishop, C. A., & Krest, S. (Eds.). (2010). *Ecotoxicology of Amphibians and Reptiles* (2nd ed.). SETAC Press.

Valiente, E., Zambrano, L., Contreras, V., & Mazari-Hiriart, M. (2010). Ecological and water quality changes in Xochimilco wetlands, Mexico City. *Freshwater Biology*, 55, 212–223.

Veith, G. D., Call, D. J., & Brooke, L. T. (1983). Structure–toxicity relationships for the fathead minnow, *Pimephales promelas*: Narcotic industrial chemicals. *Canadian Journal of Fisheries and Aquatic Sciences*, 40(6), 743–748.

Zambrano, L., Vega, E., Herrera, L. G., Prado, E., & Reynoso, V. H. (2010). Population decline of the axolotl (*Ambystoma mexicanum*) in Xochimilco, Mexico City. *Biological Conservation*, 143(11), 2746–2751.

IUCN SSC Amphibian Specialist Group. (2020). *Ambystoma mexicanum*. The IUCN Red List of Threatened Species 2020.

FAO. (2018). Globally Important Agricultural Heritage Systems (GIAHS): Chinampas of Xochimilco, Mexico.

2026年度日本農薬学会第51回大会・研究発表

～メキシコサラマンダーの飼育水中に含有する有機溶剤・農薬成分に対する耐性に関する調査報告～



2026/3/16

杉山　遥 (ハルラボ); 横江 諠衡 (東京大学)


ウーパールーパー (アホロートル) とは？

メキシコサラマンダー (*Ambystoma mexicanum*)
の国内流通個体です (**一部、他種と混血の説あり**)

原産地: **メキシコ** (**ソチミルコ湖**)
最大体長: 20～25cm
幼形成熟 (**ネオテニー**) **の形態を取れる**
寿命は20**年以上** (**変態・上陸すると**10**年程度に。。。**)
強み: **少々の水質変化はへっちゃら**
弱点: **細菌感染**, **カビ**(**免疫反応がアフリカツメガエルの半分以下**)
※ 現在は絶滅の危機 (**原因**: **都市開発や埋め立てによる生息地減少**)

胸腺が非常に小さいという報告あり. (Tリンパ球が作られにくい？)
抗菌ペプチド・粘膜などの物理的な防御を突破されると急激に衰弱する.

[IUCNレッドリストにおいて
CR (Critically Endangered：深刻な絶滅危惧)]




endothelin3 (*edn3*): 血管内皮/平滑筋の収縮に関与する endotherlin ファミリーの1つ.
全身の血流の調節に寄与.

Ref: 筑波大学_血管収縮因子エンドセリンと受容体タンパク質が形成する複合体構造を解明
https://www.tohoku.ac.jp/japanese/newimg/pressimg/tohokuuniv-press20241017_01web_ETB.pdf

活動目的:

アホロートル保全のための生態への理解
彼らを脅かすリスクファクターの抽出と対応策の検討



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



飼育環境下のアホロートル個体で耐薬品性を評価

→ 炎症 (赤身、サイトカイン分泌)・粘膜分泌、致死率で評価

アホロートル飼育時に用いる化学薬品の例とその有効濃度



塩による治療

・塩浴 ～0.5% 程度の塩水で短時間泳がせる
・周期的な 0.15～0.2% 程度の汽水飼育で微生物・カビの増殖を抑制できる.



抗生物質による治療

・オキソリン酸含有品を推奨.
濃度1/5000～1/10000で7日程度継続して使用.
理由: エロモナス菌などの感染症の原因となるグラム陰性菌に有効であるため.



イソジンによる治療
(ポビドンヨード: PVP)

・カビやウイルスなどの抗生物質が効かない相手に対して有効.
・Cタイプを 1/200程度で2～3日で十分効果があるがリスクもある.



甲殻類系の寄生虫の治療

・脱皮阻害系の農薬成分が有効.
・ジフルベンズロン
(商品名: デミリン)
イカリムシやウオジラミに1/1000 で非常に有効である.
2週間は継続した方が安心.

この他にも、水質調整剤やバクテリア剤など多種多様。

→ **アホロートルにこれらの薬品を長期的に与える場合、<u>淡水魚類の規定量より1/100～1/10000 の濃度</u>が安全。**

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図1: 化学薬品がアホロートルに与える影響1 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図1: 化学薬品がアホロートルに与える影響1 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図2: 化学薬品がアホロートルに与える影響2 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

水質汚染の原因物質の例

・重金属
- ・水銀（Hg^{2+}, 塩化水銀など）・・・ LC50:　0.05–0.3 mg/L（≈ 0.05–0.3 ppm）
- ・ニッケル（Ni^{2+}）・・・ LC50:　3–30 mg/L（≈ 3–30 ppm）
- ・カドミウム・・・ LC50:　0.01–0.5 mg/L（≈ 0.01–0.5 ppm）
- ・鉛・・・ LC50:　1–10 mg/L（≈ 1–10 ppm）
- ・銅・・・ LC50:　0.02–0.2 mg/L（≈ 0.02–0.2 ppm）

・有機溶剤
- ・エタノール・・・LC50: 11,000–15,000 mg/L (≈ 11,000–15,000 ppm)
- ・メタノール・・・ LC50: >10,000 mg/L (>10,000 ppm)
- ・イソプロパノール (2-プロパノール)・・・ LC50: 9,000–11,000 mg/L (≈ 9,000–11,000 ppm)
- ・アセトン・・・ LC50: 5,000–8,000 mg/L（≈ 5,000–8,000 ppm）
- ・ジクロロメタン (塩化メチレン)・・・LC50: 190–330 mg/L (≈ 190–330 ppm)
- ・アセトニトリル・・・ LC50: 1,600–1,800 mg/L (≈ 1,600–1,800 ppm)

・農薬などに含有される殺虫成分
- ・ピレスロイド (例：ペルメトリン)・・・ LC50: 0.0003–0.005 mg/L (≈ 0.0003–0.005 ppm)
- ・フィプロニル・・・ LC50: 0.08–0.3 mg/L (≈ 0.08–0.3 ppm)
- ・有機リン系 (例：クロルピリホス)・・・ LC50: 0.1–1 mg/L (≈ 0.1–1 ppm)
- ・ネオニコチノイド (例：イミダクロプリド)
  - 魚類・・・LC50: ≈ 100 mg/L (≈ 100 ppm)
  - 甲殻類・・・LC50: 0.01–1 mg/L (≈ 0.01–1 ppm)

※ 毒性: 農薬成分 >>> 有機溶剤

※ 1.0 ppm ＝ 1.0 mg/L ＝ 1.0×10^{-4} %

※ LC50 (Lethal Concentration 50) : 半数致死濃度 (水中)

水質汚染の原因物質の例

・重金属
・水銀（Hg^{2+}, 塩化水銀など）・・・ LC50:　0.05–0.3 mg/L（≈ 0.05–0.3 ppm）
・ニッケル（Ni^{2+}）・・・ LC50:　3–30 mg/L（≈ 3–30 ppm）
・カドミウム・・・ LC50:　0.01–0.5 mg/L（≈ 0.01–0.5 ppm）
・鉛・・・ LC50:　1–10 mg/L（≈ 1–10 ppm）
・銅・・・ LC50:　0.02–0.2 mg/L（≈ 0.02–0.2 ppm）

・有機溶剤
・エタノール・・・LC50: 11,000–15,000 mg/L (≈ 11,000–15,000 ppm)
・メタノール・・・ LC50: >10,000 mg/L (>10,000 ppm)
・イソプロパノール (2-プロパノール)・・・ LC50: 9,000–11,000 mg/L (≈ 9,000–11,000 ppm)
・アセトン・・・ LC50: 5,000–8,000 mg/L（≈ 5,000–8,000 ppm）
・ジクロロメタン (塩化メチレン)・・・LC50: 190–330 mg/L (≈ 190–330 ppm)
・アセトニトリル・・・ LC50: 1,600–1,800 mg/L (≈ 1,600–1,800 ppm)

・農薬などに含有される殺虫成分
・ピレスロイド (例：ペルメトリン)・・・ LC50: 0.0003–0.005 mg/L (≈ 0.0003–0.005 ppm)
・フィプロニル・・・ LC50: 0.08–0.3 mg/L (≈ 0.08–0.3 ppm)
・有機リン系 (例：クロルピリホス)・・・ LC50: 0.1–1 mg/L (≈ 0.1–1 ppm)
・ネオニコチノイド (例：イミダクロプリド)
- 魚類・・・LC50: ≈ 100 mg/L (≈ 100 ppm)
- 甲殻類・・・LC50: 0.01–1 mg/L (≈ 0.01–1 ppm)

※ 毒性: 農薬成分 >>> 有機溶剤

※ 1.0 ppm ＝ 1.0 mg/L ＝ 1.0×10^{-4} %

※ LC50 (Lethal Concentration 50)：半数致死濃度 (水中)

アホロートルの有機溶剤耐性 (30min 飼育水投与, 生存率: 7日間集計)




図3: 化学薬品がアホロートルに与える影響3 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)
(薬品添加: ガラス容器_2L, 生存率は7日間の飼育結果)

アホロートルの有機溶剤耐性 (30min 飼育水投与, 生存率: 7日間集計)




図3: 化学薬品がアホロートルに与える影響3 (杉山, 2023-2025)


(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)
(薬品添加: ガラス容器_2L, 生存率は7日間の飼育結果)

アホロートルの殺虫成分耐性

ピレスロイド系殺虫成分

ピレスロイドとは、除虫菊 [白花虫除菊(Tanacetum cinerariifolium (Trevir.) Sch. Bip.)] という菊に含まれている有効成分 **(天然ピレスロイド: ピレトリン および 類似物質)** の総称 または それらに似た構造の **合成化学物質 (合成ピレスロイド)** の総称。

天然ピレスロイド



ピレトリン Ⅰ	R^1: CH_3	R^2: $CH=CH_2$
Ⅱ	R^1: $COOCH_3$	R^2: $CH=CH_2$
シネリン Ⅰ	R^1: CH_3	R^2: CH_3
Ⅱ	R^1: $COOCH_3$	R^2: CH_3
ジャスモリン Ⅰ	R^1: CH_3	R^2: C_2H_5
Ⅱ	R^1: $COOCH_3$	R^2: C_2H_5

参考文献
- https://ja.wikipedia.org/wiki/ピレスロイド
- https://kotobank.jp/ぴれとりん-3217177
- https://www.boukenki.info/pyrethroid-sacchuzai-kiken-anzen-riyu/
- https://www.kincho.co.jp/seihin/business_use/basic/basic05.html
- https://www.jstage.jst.go.jp/article/jpestics1975/5/3/5_3_453/_pdf
- https://www.chem-station.com/blog/2021/11/pyrethroid.html

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図4: 化学薬品がアホロートルに与える影響4 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図4: 化学薬品がアホロートルに与える影響4 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図5: 化学薬品がアホロートルに与える影響5 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

薬品添加時の薬剤排出輸送体の発現動態解析1
(30min 飼育水投与, 24hrで鰓+周辺上皮を採取)




図6: 化学薬品がアホロートルに与える影響6 (飼育水に添加 i.e. 経皮投与_依頼分析)

(qPCR による ABCB1発現動態解析, 薬品添加: ガラス容器_2L)

※ ABCB1 (MDR1) は有機溶剤, 農薬 両方の排出に関与している報告あり。

まとめ: アホロートルの耐薬品性

・アホロートルはあらゆる薬品への 30minの暴露後 24hrという短時間で、生命活動に深刻な影響を与える可能性がある。

・アホロートルの粘膜上皮は、薬品投与への応答や排出速度より、経皮吸収速度が上回っている可能性がある。

→ 今後、
他輸送体の発現動態解析
血中や組織中に含まれる薬品のHPLCによる検出
などを実施予定。

・薬品の暴露をいかに防ぐかは今後の課題。
→ 低濃度でも長期暴露は危険か？等、順次 評価予定。

謝辞



ご清聴
ありがとうございました！



本プロジェクトに従事された皆様
・横江 諠衡 (東京大学)
・竜岡 龍騎
・阿保朗 太郎
・竜岡 龍騎
・高橋 慶　　ほか

主たるメンバー
・あすみ 悠　研究員
・枝豆 やみん　研究員
・萩原 和晃　研究員
・F.N. 三太子　特別研究員
・分析員・共同研究員の皆様
・SNS フォロワーの皆様
・他、ウーパールーパー従業員一同

提携組織・企業様 (一部抜粋)
・サイマンテクノロジー
・海人アクアリウム
・ぽれいろちゃんねる

セサミン





おとね



【Review】Behavioral Responses of Terrestrial Isopods to Pyroligneous Acid (Wood Vinegar)

Author: Yoshichika Yokoe (ICRR, The Univ. of Tokyo / Scimoun Technology / Shumei Univ.), Sugiyama Axolotl Haruka (HAL lab Axolotl, Ecology Lab of *Ambystoma mexicanum*)

【Background】 Terrestrial isopods such as pill bugs (Armadillidium spp.) are important soil animals that contribute to the decomposition of leaf litter and organic matter. They are known to select environmental conditions—such as humidity, light, and chemical cues—through behavioral responses. Avoidance behavior toward chemical substances, including pesticides and heavy metals, has been reported, and these organisms have attracted attention as model species for assessing chemical stress in soil environments. However, how they recognize volatile chemicals as olfactory stimuli and whether their behavioral responses change through experience remain insufficiently understood.

In this study, wood vinegar (pyroligneous acid) was used as a chemical stimulus to quantitatively evaluate avoidance and preference behaviors in pill bugs. In addition, we examined the sensory organs involved in odor recognition and assessed the presence or absence of habituation (adaptation) to olfactory stimuli.

【Methods】 Pill bugs were used as model organisms, and behavioral responses to wood vinegar and its dilution series were evaluated through behavioral assays. Experimental and control zones were established, and avoidance or preference behavior was quantified based on individual distribution and residence time within a fixed observation period.

To investigate sensory organs involved in odor recognition, we focused on the second antennae, which are suggested to play a role in olfactory reception. Behavioral responses of individuals with treated second antennae were compared with those of untreated individuals.

Furthermore, repeated exposures to wood vinegar were applied to the same individuals. Changes in behavioral responses relative to the initial exposure were analyzed to evaluate the presence or absence of habituation (adaptation) to the odor stimulus.

【Results and Discussion】 Pill bugs exhibited concentration-dependent avoidance behavior toward wood vinegar, suggesting that volatile chemicals contained in wood vinegar were recognized as olfactory stimuli. Individuals with treated second antennae showed significantly reduced avoidance behavior, indicating that the second antennae play an important role in odor recognition.

In addition, repeated exposure to wood vinegar resulted in a tendency toward reduced avoidance behavior, suggesting that pill bugs may exhibit habituation (adaptation) to olfactory stimuli. This behavioral change is unlikely to be merely due to sensory fatigue and may instead represent experience-dependent behavioral modulation in response to environmental chemical cues.

These findings indicate that avoidance behavior in pill bugs is not limited to transient stimulus responses

but can change plastically depending on experience. Consequently, chemical risk assessment using soil animals should consider not only acute responses but also the processes of behavioral habituation.

【References】

Appel, A. G., & Rust, M. K. (1998). Behavioral responses of terrestrial isopods to toxicants and environmental contaminants. *Environmental Toxicology and Chemistry*, 17(12), 2495–2500.

Bouchon, D., Souty-Grosset, C., & Raimond, R. (1998). Mitochondrial DNA variation and phylogeography of terrestrial isopods. *Molecular Ecology*, 7(8), 1043–1050.

Cloudsley-Thompson, J. L. (1956). Studies in diurnal rhythms in animals: Terrestrial isopods. *Journal of Experimental Biology*, 33, 576–582.

Drobne, D. (1997). Terrestrial isopods — A good choice for toxicity testing of pollutants in the terrestrial environment. *Environmental Toxicology and Chemistry*, 16(6), 1159–1164.

Drobne, D., Strus, J., & Zidar, P. (1999). Behavioural responses of terrestrial isopods to pollutants. *Applied Soil Ecology*, 11(2–3), 203–212.

Edney, E. B. (1968). Transition from water to land in isopod crustaceans. *American Zoologist*, 8(3), 309–326.

Hassall, M., & Rushton, S. P. (1984). Feeding behaviour of terrestrial isopods in relation to food quality. *Functional Ecology*, 3, 431–440.

Hornung, E. (2011). Evolutionary adaptation of oniscidean isopods to terrestrial life: Structure, physiology and behavior. *Terrestrial Arthropod Reviews*, 4(2), 95–130.

Loureiro, S., Soares, A. M. V. M., & Nogueira, A. J. A. (2005). Terrestrial avoidance behaviour tests as screening tool to assess soil contamination. *Environmental Pollution*, 138(1), 121–131.

Paris, O. H. (1963). The ecology of Armadillidium vulgare. *Ecological Monographs*, 33(2), 131–151.

Pflüger, H. J., & Burrows, M. (1990). Neural control of insect behavior: Habituation and sensory processing. *Journal of Experimental Biology*, 148, 253–275.

Schmalfuss, H. (1984). Eco-morphological strategies in terrestrial isopods. *Symposia of the Zoological Society of London*, 53, 49–63.

Souty-Grosset, C., Faberi, A., & Mocquard, J. P. (2005). Sensory perception and orientation behaviour in terrestrial isopods. *Crustaceana*, 78(6), 659–675.

Staddon, J. E. R. (2003). *Adaptive Behavior and Learning*. Cambridge University Press.

Warburg, M. R. (1993). *Evolutionary Biology of Land Isopods*. Springer.

Warburg, M. R. (2011). Habitat selection and orientation in terrestrial isopods. *ZooKeys*, 176, 1–22.

Weeks, J. M., & Andersen, O. (1989). The use of terrestrial isopods as bioindicators of soil pollution. *Environmental Pollution*, 58(3), 221–238.

Zidar, P., & Drobne, D. (2003). Antennal sensory structures and their role in environmental perception in terrestrial isopods. *Arthropod Structure & Development*, 32(2–3), 193–200.

Zidar, P., Strus, J., & Drobne, D. (2005). Avoidance behaviour of terrestrial isopods to contaminated soil. *Chemosphere*, 61(7), 938–944.

木酢液に対するダンゴムシの行動応答について

東大宇宙線研・秀明大学・ハルラボ
横江誼衡

ダンゴムシとは

ダンゴムシは
陸生等脚類（Oniscidea）に属する甲殻類。
世界に約1万種が知られている。

以下のような研究でモデル生物として例がある。
① **土壌分解者モデル**
　土壌生態系の機能研究
② **行動研究モデル**
　行動生態学研究
③ **環境毒性試験モデル**
　土壌毒性評価の指標生物

本研究では、
オカダンゴムシ（*Armadillidium vulgare*）を用いた

日本に生息する主なダンゴムシ類の例
オカダンゴムシ科（Armadillidiidae）
コシビロダンゴムシ科（Armadillidae）
フナムシ科（Ligiidae）
ワラジムシ科（Porcellionidae）



引用:福岡県 環境部 自然環境課

農業生態系におけるダンゴムシの役割

・土壌生物は農業生態系の機能維持に重要である。
主な機能
有機物分解、栄養循環、微生物活動の促進

特にダンゴムシ（陸生等脚類）は
落葉分解者として重要かつ栄養循環に寄与
ことが知られている
※生きている新鮮な植物（生葉）を食べることもある為、
　農業害虫として扱われることもある

(Zimmer 2002; Paoletti & Hassall 1999)

木酢とは何か（化学組成）

木酢液（*pyroligneous acid*）は，木材や植物バイオマスを熱分解（pyrolysis）した際に発生する揮発性成分を冷却・凝縮して得られる液体である。
木酢液は**200種類以上の化合物を含む複雑な混合物**であり，主に以下の化学成分から構成されることが報告されている
(Mohan et al., 2006; Wei et al., 2022)。

主な化学成分
- 有機酸（主に酢酸）
- フェノール類
- アルコール類
- ケトン類
- アルデヒド類

具体的な成分としては，
- 酢酸（*acetic acid*）
- メタノール（*methanol*）
- アセトン（*acetone*）
- フルフラール（*furfural*）
- グアイアコール（*guaiacol*）
- シリンゴール（*syringol*）

などが報告されている（Mohan et al., 2006）。
これらの揮発性有機化合物は，**独特の臭気や抗菌性，防虫性などの生理活性**を示すことが知られている
(Wei et al., 2022)。

ただし、ダンゴムシへの影響に関する学術報告は少ない。

ダンゴムシの行動研究

①化学物質の行動への影響
化学物質は
移動行動＆探索行動
などの **行動応答** に影響を与える報告。
(Paoletti & Hassall, 1999)

② 集団行動の数理モデル
ダンゴムシは　**集合行動（aggregation）**を示す
個体の環境選好（光と壁）と社会相互作用に
よって集団形成が生じると考えられている。
(Devigne et al., 2011)
agent-based model ＋ 社会相互作用＋影＋壁

ダンゴムシの集団行動については，個体の環境選好や社会相互作用を考慮した
数理モデル研究が報告されている。

化学物質応答を組み込んだ集団行動モデルの研究例はほとんどない。
化学物質応答を考慮した集団行動の数理モデル(diffuse model）を
ダンゴムシに適用することを目標
本研究は、その
予備実験的な検討
として，木酢液に対するダンゴムシの行動応答を観察した。

$$\frac{\partial \rho}{\partial t} = \boldsymbol{\nabla} \cdot (D\,\nabla\rho) - \boldsymbol{\nabla} \cdot (\boldsymbol{v} \cdot \rho)$$
$$\boldsymbol{v} = -\boldsymbol{\nabla} U$$

Agent-based モデルと拡散モデル

Agent-based モデル
・個体ごとの行動ルールを設定
・個体位置をシミュレーション
・社会性相互作用などを詳細に研究可能

$$\frac{\partial x}{\partial t} = \underbrace{v_0 e_i}_{\text{自発的な移動（探索行動）}} - \underbrace{\boldsymbol{\nabla} U}_{\text{外的作用}} + \underbrace{\eta_{i(t)}}_{\text{ランダム行動（ノイズ）}}$$

② 拡散モデル
・集団の時間分布を扱える
・実験データと比較しやすい
・パラメータ推定が可能。
・個体ノイズを平均化できる

$$\frac{\partial \rho}{\partial t} = \underbrace{\boldsymbol{\nabla} \cdot (D\,\nabla\rho)}_{\text{ランダムな探索}} - \underbrace{\boldsymbol{\nabla} \cdot (\boldsymbol{v} \cdot \rho)}_{\text{外的作用}} \quad \boldsymbol{v} = -\boldsymbol{\nabla} U$$

② 集団行動の数理モデル
ダンゴムシは　**集合行動（aggregation）**を示す
個体の環境選好（光）と社会相互作用によって
集団形成が生じると考えられている。
(Devigne et al., 2011)
agent-based model ＋ 社会相互作用　＋　影　＋壁

⇒

Agent-basedモデルだと
・未知の個体ルールの影響を受ける可能性
↓
拡散モデルだと
・ランダム移動と環境に対するバイアスとして表現できる可能性がある

実験概要

実験装置
直径 **9 cm のシャーレ**を用いた二択実験
左右に以下の餌を配置
①水で湿らせた後に乾燥させた餌（対照）
②木酢液（右図）で湿らせた後に乾燥させた餌
餌には **金魚用飼料**を使用した。

実験条件
室温；25度　湿度：32%

試行回数
5セット

実験生物
オカダンゴムシ（1セット5匹）
2025年10月ごろ、千葉県柏市で森林採集した個体

観察方法
1時間動画撮影





GEX　金魚元気




中島商事 トヨチュー
熟成木酢原1500ML

結果：N_{left}:N_{right}







$$Ratio = \frac{N_{left}}{N_{left} + N_{right}}$$




結果：N_{middle} の割合



$$Ratio = \frac{N_{middle}}{N_{all}}$$





結果まとめ

対照餌側vs 木酢処理餌の行動応答
・ダンゴムシは木酢処理餌より **対照餌側を選好する傾向**を示した。

木酢処理餌vs 木酢処理餌
・どちらか片方に寄ることも少なく、中間層にいる割合も多い。

対照餌側vs対照餌側
・どちらか片方に寄ることも多く、中間層にいる割合も少ない。

考察

本実験では，木酢処理餌と対照餌を用いた二択実験を行い，ダンゴムシの行動応答を観察した。
その結果，1時間の観察の範囲内では時間発展は見られなかったものの，**ダンゴムシは対照餌側へ移動する傾向が確認された。**
また，木酢処理餌同士および対照餌同士の条件を比較すると，
対照餌のみが存在する場合
　社会相互作用により片側の餌に集まる傾向が見られた。
木酢処理餌のみが存在する場合
　餌への接近を避ける行動が見られ，中間領域に滞在する割合が増加した。

以上の結果から，**木酢液に含まれる化学物質がダンゴムシの行動に影響を与える chemical cue として作用している可能性が示唆された**

このような行動変化は **ランダムな移動に化学刺激によるバイアスが加わった運動**として捉えることができ，拡散運動に化学ポテンシャル U_{chemical}を導入したモデルによって記述できる可能性がある。

$$U = \boldsymbol{U_{chemical}} + U_{social} + U_{wall} + U_{shadow} = \boldsymbol{U_{repulsive}} + \boldsymbol{U_{attractive}}$$

今後の研究展望（数理モデル）

① 実験条件の拡張
個体数や木酢液濃度，湿度などの環境条件を変化させた実験を行い，ダンゴムシ行動の再現性および条件依存性を検証する。

② 行動の定量解析
動画データから個体の移動軌跡や集団形成の時間変化を解析し，行動を定量的に評価する。

③ 数理モデルによる解析
得られたデータを用いて集団挙動を拡散モデルとして記述し，

$$U = U_{\mathrm{chemical}} + U_{\mathrm{social}} + U_{\mathrm{wall}}$$

で表される有効ポテンシャル Uの推定を試みる。

④ 化学要因の推定と応用可能性
特に U_{chemical}の推定により，木酢液中の忌避作用を持つ化学物質の推定につながる可能性があり，将来的には天然由来忌避剤などへの応用も期待される。

結論

本研究では，木酢処理餌と対照餌を用いた二択実験を行い，ダンゴムシの行動応答を観察した。

その結果，観察時間内においてダンゴムシは**木酢処理餌側よりも対照餌側へ移動する傾向**が見られた。


このことから，**木酢液に含まれる化学物質がダンゴムシの行動に影響を与えている可能性**が示唆された。

一方で，ダンゴムシには**社会相互作用や壁沿い行動などの要因**も存在するため，本研究で観察された行動パターンはこれらの複合的な影響によって形成されている可能性がある。

以上の結果は，ダンゴムシの行動が**化学刺激・社会相互作用・環境要因**などの影響を受けることを示唆しており，**これらの要因を考慮した数理モデルによって集団行動を記述できる可能性がある。**

ご清聴ありがとうございました。

生物と農薬



著: 杉山 遥 [1]; 横江 諠衡 [1, 2, 3, 4]

1. ハルラボ・メキシコサラマンダー生態研究部,

2. 東京大学・理・物理・宇宙線研, 3. サイマンテクノロジー, 4. 秀明大学

【総説】メキシコサラマンダーの飼育水中に含有する有機溶剤・農薬成分に対する耐性に関する調査報告

著: 杉山 遥 (ハルラボ・メキシコサラマンダー生態研究部), 横江 諠衡 (東京大学・理・物理・宇宙線研/サイマンテクノロジー/秀明大学), F.N. 三太子 (ハルラボ・アホロートルレスキュー), 枝豆 やみん (ハルラボ・淡水魚生態研究部), 竜岡 龍騎 (トラフサンショウウオラボ), 阿保朗 太朗 (ハルラボ・ファーム)

【背景・目的】 メキシコサラマンダー (*Ambystoma mexicanum*, 以下、アホロートル) は、飼育個体はウーパールーパーと呼ばれ親しまれている。一方で、野生下において絶滅の危機に瀕しており [IUCN レッドリストにおいて CR (Critically Endangered：深刻な絶滅危惧)]、その個体数を大きく減らしているのが現状である。この原因の一つに、生息地であるメキシコ・ソチミルコ湖の観光地化や土地開発による水質汚染が挙げられる。生活排水や工業廃液等が同湖へと流入し、年々水質が悪化しているのが現状である。また周辺地域では、チナンパと呼ばれる浮島上で農作を行う伝統的な古式農法が復活しつつあり、農薬等の成分が今後水中により多く混入される危険性が考えられる。こうした現状からアホロートルの野生個体を救うため、飼育下の個体における飼育水中に含まれる薬剤への応答性について定量を行った。

【方法】 アホロートル飼育水中に、飼育で用いられる汎用的な水質調整剤 (カルキ中和剤に含有するチオ硫酸ナトリウム, pH 調整剤, PSB バクテリア剤 等) や、石鹸・洗剤などの生活排水として生息地に流入する可能性のある界面活性剤、工業的に水質汚染に関わる可能性のあるいくつかの有機溶剤 [エタノール (水溶性), メタノール (水溶性), IPA (イソプロ) (水溶性), アセトン (水溶性), ジクロロメタン (水溶性)] および 農薬成分 (ピレスロイド系化合物) を複数の濃度条件で添加し、薬傷に伴った炎症や粘膜の発生や致死性 (LD50) 等の薬剤の影響に伴う表現型について定量的・統計的に確認を行った。具体的には、直接的に飼育水槽への添加と個体の観察・定量や、他の多くの飼育者に聞き取りを実施することで、あらゆる薬品への暴露の影響について統計的に評価可能となるよう情報収集した。更に、哺乳類や魚類において薬剤排出に関わるトランスポーター等の遺伝子群の相同因子を、GeneBank や Blast 等のデータベースから選定し、薬品に暴露した個体の組織片から mRNA を抽出し、qRT-PCR 法にて薬剤暴露前後の発現動態について確認し、分子的な知見と併せることでより精度の高い評価を行った。

【結果および考察】 汎用的な飼育用の薬剤であっても、アホロートルは一般的な観賞用淡水魚に対する規定量の 1/1000 程度で使用しないと、多くが表現型の異常を示すことが明らかとなった (例外あり)。酷い場合には薬剤処理から 24 時間後には亡くなってしまうことが分かった。殆どの有機溶剤で、1000ppm (0.1%)〜1.0% の間で急激にその致死性を増大させるという傾向を示した。これらの傾向や聞き取りの結果から、アホロートルは薬品への暴露が即、命に係わる可能性が示唆された。また薬剤の輸送体の相同因子発現は一部確認された一方で、上皮を浸透した薬品の取り込み速度よりも、薬品排出速度の方が顕著に遅い可能性があることが示唆された。これらの知見は、環境汚染がいかにアホロートルの生命を脅かすかという新たな知見を示す重要な結果である。

【参考文献】

Ambudkar, S. V., Dey, S., Hrycyna, C. A., Ramachandra, M., Pastan, I., & Gottesman, M. M. (1999). Biochemical, cellular, and pharmacological aspects of the multidrug transporter P-glycoprotein. *Annual Review of Pharmacology and Toxicology*, 39, 361–398.

Bard, S. M. (2000). Multixenobiotic resistance as a cellular defense mechanism in aquatic organisms. *Aquatic Toxicology*, 48(4), 357–389.

Birge, W. J., Westerman, A. G., & Spromberg, J. A. (2000). Comparative toxicology and risk assessment of amphibians. In D. W. Sparling, G. Linder, & C. A. Bishop (Eds.), *Ecotoxicology of Amphibians and Reptiles* (pp. 727–791). SETAC Press.

Blaustein, A. R., & Kiesecker, J. M. (2002). Complexity in conservation: Lessons from the global decline of amphibian populations. *Nature*, 415, 681–684.

Bradbury, S. P., & Coats, J. R. (1989). Comparative toxicology of the pyrethroid insecticides. *Reviews of Environmental Contamination and Toxicology*, 108, 133–177.

Brühl, C. A., Schmidt, T., Pieper, S., & Alscher, A. (2013). Terrestrial pesticide exposure of amphibians: An underestimated cause of global decline? *Scientific Reports*, 3, 1135.

Duellman, W. E., & Trueb, L. (1994). *Biology of Amphibians*. Johns Hopkins University Press.

González-Pozo, A., et al. (2016). Urban agriculture and environmental sustainability in Xochimilco, Mexico City. *Environmental Management*, 58, 101–115.

Lewis, M. A. (1991). Chronic and acute toxicities of surfactants and detergent builders to aquatic organisms: A review and risk assessment. *Water Research*, 25(1), 101–113.

Lipnick, R. L. (1991). Outliers: Their origin and use in the classification of molecular mechanisms of toxicity. *Science of the Total Environment*, 109–110, 131–153.

Mann, R. M., Hyne, R. V., Choung, C. B., & Wilson, S. P. (2009). Amphibians and agricultural chemicals: Review of the risks in a complex environment. *Environmental Pollution*, 157(11), 2903–2927.

Maund, S. J., Hamer, M. J., Lane, M. C. G., et al. (2002). Partitioning, bioavailability, and toxicity of the pyrethroid insecticide cypermethrin in aquatic environments. *Environmental Toxicology and Chemistry*, 21(1), 9–15.

McCarty, L. S., & Mackay, D. (1993). Enhancing ecotoxicological modeling and assessment: Narcotic mode of action. *Environmental Toxicology and Chemistry*, 12(4), 629–643.

Quaranta, A., Bellantuono, V., Cassano, G., & Lippe, C. (2009). Why amphibians are more sensitive than mammals to xenobiotics. *PLoS ONE*, 4(11), e7699.

Smital, T., Sauerborn, R., & Hackenberger, B. K. (2004). Inducibility of the P-glycoprotein transport activity in the marine mussel *Mytilus galloprovincialis*. *Aquatic Toxicology*, 70(2), 131–141.

Sparling, D. W., Linder, G., Bishop, C. A., & Krest, S. (Eds.). (2010). *Ecotoxicology of Amphibians and Reptiles* (2nd ed.). SETAC Press.

Valiente, E., Zambrano, L., Contreras, V., & Mazari-Hiriart, M. (2010). Ecological and water quality changes in Xochimilco wetlands, Mexico City. *Freshwater Biology*, 55, 212–223.

Veith, G. D., Call, D. J., & Brooke, L. T. (1983). Structure–toxicity relationships for the fathead minnow,

Pimephales promelas: Narcotic industrial chemicals. *Canadian Journal of Fisheries and Aquatic Sciences*, 40(6), 743–748.

Zambrano, L., Vega, E., Herrera, L. G., Prado, E., & Reynoso, V. H. (2010). Population decline of the axolotl (*Ambystoma mexicanum*) in Xochimilco, Mexico City. *Biological Conservation*, 143(11), 2746–2751.

IUCN SSC Amphibian Specialist Group. (2020). *Ambystoma mexicanum*. The IUCN Red List of Threatened Species 2020.

FAO. (2018). Globally Important Agricultural Heritage Systems (GIAHS): Chinampas of Xochimilco, Mexico.

2026年度日本農薬学会第51回大会・研究発表

～メキシコサラマンダーの飼育水中に含有する有機溶剤・農薬成分に対する耐性に関する調査報告～



2026/3/16

杉山　遥 (ハルラボ); 横江 諠衡 (東京大学)


ウーパールーパー (アホロートル) とは？

メキシコサラマンダー (*Ambystoma mexicanum*)
の国内流通個体です (**一部、他種と混血の説あり**)

原産地: **メキシコ** (**ソチミルコ湖**)
最大体長: 20～25cm
幼形成熟 (**ネオテニー**) **の形態を取れる**
寿命は20**年以上** (**変態・上陸すると**10**年程度に。。。**)
強み: **少々の水質変化はへっちゃら**
弱点: **細菌感染**, **カビ**(**免疫反応がアフリカツメガエルの半分以下**)
※ 現在は絶滅の危機 (**原因**: **都市開発や埋め立てによる生息地減少**)

胸腺が非常に小さいという報告あり. (Tリンパ球が作られにくい？)
抗菌ペプチド・粘膜などの物理的な防御を突破されると急激に衰弱する.

[IUCNレッドリストにおいて
CR (Critically Endangered：深刻な絶滅危惧)]




endothelin3 (*edn3*): 血管内皮/平滑筋の収縮に関与する endotherlin ファミリーの1つ.
全身の血流の調節に寄与.

Ref: 筑波大学_血管収縮因子エンドセリンと受容体タンパク質が形成する複合体構造を解明
https://www.tohoku.ac.jp/japanese/newimg/pressimg/tohokuuniv-press20241017_01web_ETB.pdf

活動目的:

アホロートル保全のための生態への理解
彼らを脅かすリスクファクターの抽出と対応策の検討



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



アホロートル: 絶滅危惧種

主要な原因

・生息地 (メキシコ・ソチミルコ湖) 周辺の都市開発や観光地化
・古式農法 (チナンパ) や大規模農業

→ 生息地減少＋生息地 (湖) の水質悪化



飼育環境下のアホロートル個体で耐薬品性を評価

→ 炎症 (赤身、サイトカイン分泌)・粘膜分泌、致死率で評価

アホロートル飼育時に用いる化学薬品の例とその有効濃度



塩による治療

・塩浴 ～0.5% 程度の塩水で短時間泳がせる
・周期的な 0.15～0.2% 程度の汽水飼育で微生物・カビの増殖を抑制できる.



抗生物質による治療

・オキソリン酸含有品を推奨.
濃度1/5000～1/10000で7日程度継続して使用.
理由: エロモナス菌などの感染症の原因となるグラム陰性菌に有効であるため.



イソジンによる治療
(ポビドンヨード: PVP)

・カビやウイルスなどの抗生物質が効かない相手に対して有効.
・Cタイプを 1/200程度で2～3日で十分効果があるがリスクもある.



甲殻類系の寄生虫の治療

・脱皮阻害系の農薬成分が有効.
・ジフルベンズロン
(商品名: デミリン)
イカリムシやウオジラミに1/1000 で非常に有効である.
2週間は継続した方が安心.

この他にも、水質調整剤やバクテリア剤など多種多様。

→ **アホロートルにこれらの薬品を長期的に与える場合、<u>淡水魚類の規定量より1/100～1/10000 の濃度</u>が安全。**

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図1: 化学薬品がアホロートルに与える影響1 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図1: 化学薬品がアホロートルに与える影響1 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

アホロートルの耐薬品性評価 (30min 飼育水投与, 24hr経過観察)




図2: 化学薬品がアホロートルに与える影響2 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)

水質汚染の原因物質の例

・重金属
- ・水銀（Hg^{2+}, 塩化水銀など）・・・ LC50:　0.05–0.3 mg/L（≈ 0.05–0.3 ppm）
- ・ニッケル（Ni^{2+}）・・・ LC50:　3–30 mg/L（≈ 3–30 ppm）
- ・カドミウム・・・ LC50:　0.01–0.5 mg/L（≈ 0.01–0.5 ppm）
- ・鉛・・・ LC50:　1–10 mg/L（≈ 1–10 ppm）
- ・銅・・・ LC50:　0.02–0.2 mg/L（≈ 0.02–0.2 ppm）

・有機溶剤
- ・エタノール・・・LC50: 11,000–15,000 mg/L (≈ 11,000–15,000 ppm)
- ・メタノール・・・ LC50: >10,000 mg/L (>10,000 ppm)
- ・イソプロパノール (2-プロパノール)・・・ LC50: 9,000–11,000 mg/L (≈ 9,000–11,000 ppm)
- ・アセトン・・・ LC50: 5,000–8,000 mg/L（≈ 5,000–8,000 ppm）
- ・ジクロロメタン (塩化メチレン)・・・LC50: 190–330 mg/L (≈ 190–330 ppm)
- ・アセトニトリル・・・ LC50: 1,600–1,800 mg/L (≈ 1,600–1,800 ppm)

・農薬などに含有される殺虫成分
- ・ピレスロイド (例：ペルメトリン)・・・ LC50: 0.0003–0.005 mg/L (≈ 0.0003–0.005 ppm)
- ・フィプロニル・・・ LC50: 0.08–0.3 mg/L (≈ 0.08–0.3 ppm)
- ・有機リン系 (例：クロルピリホス)・・・ LC50: 0.1–1 mg/L (≈ 0.1–1 ppm)
- ・ネオニコチノイド (例：イミダクロプリド)
  - 魚類・・・LC50: ≈ 100 mg/L (≈ 100 ppm)
  - 甲殻類・・・LC50: 0.01–1 mg/L (≈ 0.01–1 ppm)

※ 毒性: 農薬成分 >>> 有機溶剤

※ 1.0 ppm ＝ 1.0 mg/L ＝ 1.0×10^{-4} %

※ LC50 (Lethal Concentration 50) : 半数致死濃度 (水中)

水質汚染の原因物質の例

・重金属
・水銀（Hg^{2+}, 塩化水銀など）・・・ LC50:　0.05–0.3 mg/L（≈ 0.05–0.3 ppm）
・ニッケル（Ni^{2+}）・・・ LC50:　3–30 mg/L（≈ 3–30 ppm）
・カドミウム・・・ LC50:　0.01–0.5 mg/L（≈ 0.01–0.5 ppm）
・鉛・・・ LC50:　1–10 mg/L（≈ 1–10 ppm）
・銅・・・ LC50:　0.02–0.2 mg/L（≈ 0.02–0.2 ppm）

・有機溶剤
・エタノール・・・LC50: 11,000–15,000 mg/L (≈ 11,000–15,000 ppm)
・メタノール・・・ LC50: >10,000 mg/L (>10,000 ppm)
・イソプロパノール (2-プロパノール)・・・ LC50: 9,000–11,000 mg/L (≈ 9,000–11,000 ppm)
・アセトン・・・ LC50: 5,000–8,000 mg/L（≈ 5,000–8,000 ppm）
・ジクロロメタン (塩化メチレン)・・・LC50: 190–330 mg/L (≈ 190–330 ppm)
・アセトニトリル・・・ LC50: 1,600–1,800 mg/L (≈ 1,600–1,800 ppm)

・農薬などに含有される殺虫成分
・ピレスロイド (例：ペルメトリン)・・・ LC50: 0.0003–0.005 mg/L (≈ 0.0003–0.005 ppm)
・フィプロニル・・・ LC50: 0.08–0.3 mg/L (≈ 0.08–0.3 ppm)
・有機リン系 (例：クロルピリホス)・・・ LC50: 0.1–1 mg/L (≈ 0.1–1 ppm)
・ネオニコチノイド (例：イミダクロプリド)
- 魚類・・・LC50: ≈ 100 mg/L (≈ 100 ppm)
- 甲殻類・・・LC50: 0.01–1 mg/L (≈ 0.01–1 ppm)

※ 毒性: 農薬成分 >>> 有機溶剤

※ 1.0 ppm ＝ 1.0 mg/L ＝ 1.0×10^{-4} %

※ LC50 (Lethal Concentration 50)：半数致死濃度 (水中)

アホロートルの有機溶剤耐性 (30min 飼育水投与, 生存率: 7日間集計)




図3: 化学薬品がアホロートルに与える影響3 (杉山, 2023-2025)
(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)
(薬品添加: ガラス容器_2L, 生存率は7日間の飼育結果)

アホロートルの有機溶剤耐性 (30min 飼育水投与, 生存率: 7日間集計)




図3: 化学薬品がアホロートルに与える影響3 (杉山, 2023-2025)


(飼育水に添加 i.e. 経皮投与, 薬品添加: ガラス容器_2L, 容器指定の場合は素材別)
(薬品添加: ガラス容器_2L, 生存率は7日間の飼育結果)

アホロートルの殺虫成分耐性

ピレスロイド系殺虫成分

ピレスロイドとは、除虫菊 [白花虫除菊(Tanacetum cinerariifolium (Trevir.) Sch. Bip.)] という菊に含まれている有効成分 **(天然ピレスロイド: ピレトリン および 類似物質)** の総称 または それらに似た構造の **合成化学物質 (合成ピレスロイド)** の総称。

天然ピレスロイド



ピレトリン Ⅰ	R^1: CH_3	R^2: $CH=CH_2$
Ⅱ	R^1: $COOCH_3$	R^2: $CH=CH_2$
シネリン Ⅰ	R^1: CH_3	R^2: CH_3
Ⅱ	R^1: $COOCH_3$	R^2: CH_3
ジャスモリン Ⅰ	R^1: CH_3	R^2: C_2H_5
Ⅱ	R^1: $COOCH_3$	R^2: C_2H_5

参考文献
- https://ja.wikipedia.org/wiki/ピレスロイド
- https://kotobank.jp/ぴれとりん-3217177
- https://www.boukenki.info/pyrethroid-sacchuzai-kiken-anzen-riyu/
- https://www.kincho.co.jp/seihin/business_use/basic/basic05.html
- https://www.jstage.jst.go.jp/article/jpestics1975/5/3/5_3_453/_pdf
- https://www.chem-station.com/blog/2021/11/pyrethroid.html

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図4: 化学薬品がアホロートルに与える影響4 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図4: 化学薬品がアホロートルに与える影響4 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

アホロートルの殺虫成分耐性
(30min 飼育水投与, 生存率: 7〜600日間集計)




図5: 化学薬品がアホロートルに与える影響5 (飼育水に添加 i.e. 経皮投与_依頼分析)

(薬品添加: ガラス容器_2L)

薬品添加時の薬剤排出輸送体の発現動態解析1
(30min 飼育水投与, 24hrで鰓+周辺上皮を採取)




図6: 化学薬品がアホロートルに与える影響6 (飼育水に添加 i.e. 経皮投与_依頼分析)

(qPCR による ABCB1発現動態解析, 薬品添加: ガラス容器_2L)

※ ABCB1 (MDR1) は有機溶剤, 農薬 両方の排出に関与している報告あり。

まとめ: アホロートルの耐薬品性

・アホロートルはあらゆる薬品への 30minの暴露後 24hrという短時間で、生命活動に深刻な影響を与える可能性がある。

・アホロートルの粘膜上皮は、薬品投与への応答や排出速度より、経皮吸収速度が上回っている可能性がある。

→ 今後、
他輸送体の発現動態解析
血中や組織中に含まれる薬品のHPLCによる検出
などを実施予定。

・薬品の暴露をいかに防ぐかは今後の課題。
→ 低濃度でも長期暴露は危険か？等、順次 評価予定。

謝辞



ご清聴
ありがとうございました！



本プロジェクトに従事された皆様
・横江 諠衡 (東京大学)
・竜岡 龍騎
・阿保朗 太郎
・竜岡 龍騎
・高橋 慶　　ほか

主たるメンバー
・あすみ 悠　研究員
・枝豆 やみん　研究員
・萩原 和晃　研究員
・F.N. 三太子　特別研究員
・分析員・共同研究員の皆様
・SNS フォロワーの皆様
・他、ウーパールーパー従業員一同

提携組織・企業様 (一部抜粋)
・サイマンテクノロジー
・海人アクアリウム
・ぽれいろちゃんねる

セサミン





おとね



【総説】 木酢液に対するダンゴムシの行動応答について

著: 横江 誼衡 (東京大学・理・物理・宇宙線研/サイマンテクノロジー/秀明大学)、杉山 遥 (ハルラボ・メキシコサラマンダー生態研究部)

【背景・目的】 陸生等脚類であるダンゴムシ（**Armadillidium spp.** など）は、落葉や腐植物の分解を担う重要な土壌動物であり、湿度、光、化学物質といった環境情報を行動によって選択することが知られている。農薬や重金属などの化学物質に対する忌避行動が報告され、土壌環境における化学ストレス評価のモデル生物として注目されている一方で、揮発性化学物質を匂い刺激としてどのように認識し、さらにその行動応答が経験によって変化するのかについては十分に検討されていない。
本研究では、木酢液を化学刺激源として用い、ダンゴムシの忌避・選好行動を定量的に評価するとともに、匂い認識に関与する感覚器官および匂い刺激に対する慣れ（順応）の有無について検討することを目的とした。

【方法】 ダンゴムシをモデル生物として用い、木酢液およびその希釈系列に対する行動応答を行動実験により評価した。試験区と対照区を設け、一定時間内の個体分布および滞在時間を指標として忌避・選好行動を定量化した。
匂い認識に関与する感覚器官の検証として、嗅覚受容への関与が示唆されている第二触角に着目し、第二触角を処理した個体と未処理個体の行動応答を比較した。
さらに、同一個体に対して木酢液への繰り返し暴露を行い、初回暴露時と比較した行動応答の変化を解析することで、匂い刺激に対する慣れ（順応）の有無を評価した。

【結果および考察】 ダンゴムシは木酢液に対して濃度依存的な忌避行動を示し、木酢液中に含まれる揮発性化学物質を匂い刺激として認識している可能性が示唆された。第二触角を処理した個体では忌避行動が有意に低下し、木酢液の匂い認識に第二触角が重要な役割を果たしていることが示された。
また、木酢液への繰り返し暴露により忌避行動が減弱する傾向が認められ、ダンゴムシが匂い刺激に対して慣れ（順応）を示す可能性が示唆された。この行動変化は、単なる感覚疲労ではなく、環境中の化学刺激に対する経験依存的な行動調節である可能性が考えられる。
これらの結果は、ダンゴムシの忌避行動が一過的な刺激応答にとどまらず、経験に応じて可塑的に変化しうることを示しており、土壌動物を用いた化学物質評価において、急性応答のみならず行動の順応過程を考慮する必要性を示唆するものである。

【参考文献】

Appel, A. G., & Rust, M. K. (1998). Behavioral responses of terrestrial isopods to toxicants and environmental contaminants. *Environmental Toxicology and Chemistry*, 17(12), 2495–2500.

Bouchon, D., Souty-Grosset, C., & Raimond, R. (1998). Mitochondrial DNA variation and phylogeography of terrestrial isopods. *Molecular Ecology*, 7(8), 1043–1050.

Cloudsley-Thompson, J. L. (1956). Studies in diurnal rhythms in animals: Terrestrial isopods. *Journal of Experimental Biology*, 33, 576–582.

Drobne, D. (1997). Terrestrial isopods — A good choice for toxicity testing of pollutants in the terrestrial environment. *Environmental Toxicology and Chemistry*, 16(6), 1159–1164.

Drobne, D., Strus, J., & Zidar, P. (1999). Behavioural responses of terrestrial isopods to pollutants. *Applied Soil Ecology*, 11(2–3), 203–212.

Edney, E. B. (1968). Transition from water to land in isopod crustaceans. *American Zoologist*, 8(3), 309–326.

Hassall, M., & Rushton, S. P. (1984). Feeding behaviour of terrestrial isopods in relation to food quality. *Functional Ecology*, 3, 431–440.

Hornung, E. (2011). Evolutionary adaptation of oniscidean isopods to terrestrial life: Structure, physiology and behavior. *Terrestrial Arthropod Reviews*, 4(2), 95–130.

Loureiro, S., Soares, A. M. V. M., & Nogueira, A. J. A. (2005). Terrestrial avoidance behaviour tests as screening tool to assess soil contamination. *Environmental Pollution*, 138(1), 121–131.

Paris, O. H. (1963). The ecology of Armadillidium vulgare. *Ecological Monographs*, 33(2), 131–151.

Pflüger, H. J., & Burrows, M. (1990). Neural control of insect behavior: Habituation and sensory processing. *Journal of Experimental Biology*, 148, 253–275.

Schmalfuss, H. (1984). Eco-morphological strategies in terrestrial isopods. *Symposia of the Zoological Society of London*, 53, 49–63.

Souty-Grosset, C., Faberi, A., & Mocquard, J. P. (2005). Sensory perception and orientation behaviour in terrestrial isopods. *Crustaceana*, 78(6), 659–675.

Staddon, J. E. R. (2003). *Adaptive Behavior and Learning*. Cambridge University Press.

Warburg, M. R. (1993). *Evolutionary Biology of Land Isopods*. Springer.

Warburg, M. R. (2011). Habitat selection and orientation in terrestrial isopods. *ZooKeys*, 176, 1–22.

Weeks, J. M., & Andersen, O. (1989). The use of terrestrial isopods as bioindicators of soil pollution. *Environmental Pollution*, 58(3), 221–238.

Zidar, P., & Drobne, D. (2003). Antennal sensory structures and their role in environmental perception in terrestrial isopods. *Arthropod Structure & Development*, 32(2–3), 193–200.

Zidar, P., Strus, J., & Drobne, D. (2005). Avoidance behaviour of terrestrial isopods to contaminated soil. *Chemosphere*, 61(7), 938–944.

木酢液に対するダンゴムシの行動応答について

東大宇宙線研・秀明大学・ハルラボ
横江誼衡

ダンゴムシとは

ダンゴムシは
陸生等脚類（Oniscidea）に属する甲殻類。
世界に約1万種が知られている。


以下のような研究でモデル生物として例がある。
① **土壌分解者モデル**
　土壌生態系の機能研究
② **行動研究モデル**
　行動生態学研究
③ **環境毒性試験モデル**
　土壌毒性評価の指標生物

本研究では、
オカダンゴムシ（*Armadillidium vulgare*）を用いた

日本に生息する主なダンゴムシ類の例
オカダンゴムシ科（Armadillidiidae）
コシビロダンゴムシ科（Armadillidae）
フナムシ科（Ligiidae）
ワラジムシ科（Porcellionidae）



引用:福岡県 環境部 自然環境課




農業生態系におけるダンゴムシの役割

・土壌生物は農業生態系の機能維持に重要である。
主な機能
有機物分解、栄養循環、微生物活動の促進

特にダンゴムシ（陸生等脚類）は
落葉分解者として重要かつ栄養循環に寄与
ことが知られている
※生きている新鮮な植物（生葉）を食べることもある為、
　農業害虫として扱われることもある

(Zimmer 2002; Paoletti & Hassall 1999)



木酢とは何か

木酢液（wood vinegar）は
　木酢液（pyroligneous acid）は
　木材や植物バイオマスを熱分解した際に生じる煙を凝縮して得られる液体である。
主成分
　有機酸（酢酸など）
　フェノール類
　アルコール類
　ケトン類
これらの化合物が
独特の臭気や生理活性を示すことが知られている。

木酢液がダンゴムシに効果があるとするWeb記事は散見されるが,
ダンゴムシの行動応答を検討した学術研究はほとんど報告されていない。

ダンゴムシの行動研究

①化学物質の行動への影響
化学物質は
移動行動＆探索行動
などの **行動応答** に影響を与える報告。
(Paoletti & Hassall, 1999)

② 集団行動の数理モデル
ダンゴムシは　**集合行動（aggregation）**を示す
個体の環境選好（光）と社会相互作用によって
集団形成が生じると考えられている。
(Devigne et al., 2011)
agent-based model ＋ 社会相互作用　＋　光

ダンゴムシの集団行動については，個体の環境選好や社会相互作用を考慮した
数理モデル研究が報告されている。

化学物質応答を組み込んだ集団行動モデルの研究例はほとんどない。
化学物質応答を考慮した集団行動の数理モデル(diffuse model）を
ダンゴムシに適用することを目標
本研究は、その
予備実験的な検討
として，木酢液に対するダンゴムシの行動応答を観察した。

$$\frac{\partial \rho}{\partial t} = \boldsymbol{\nabla} \cdot (D\,\nabla\rho) - \boldsymbol{\nabla} \cdot (\boldsymbol{v} \cdot \rho)$$
$$\boldsymbol{v} = -\boldsymbol{\nabla} U$$

Agent-based モデルと拡散モデル

Agent-based モデル
・個体ごとの行動ルールを設定
・個体位置をシミュレーション
・社会性相互作用などを詳細に研究可能

$$\frac{\partial x}{\partial t} = \underbrace{v_0 e_i}_{\text{自発的な移動（探索行動）}} - \underbrace{\boldsymbol{\nabla} U}_{\text{外的作用}} + \underbrace{\eta_{i(t)}}_{\text{ランダム行動（ノイズ）}}$$

② 拡散モデル
・集団の時間分布を扱える
・実験データと比較しやすい
・パラメータ推定が可能。
・個体ノイズを平均化できる

$$\frac{\partial \rho}{\partial t} = \underbrace{\boldsymbol{\nabla} \cdot (D\,\nabla\rho)}_{\text{ランダムな探索}} - \underbrace{\boldsymbol{\nabla} \cdot (\boldsymbol{v} \cdot \rho)}_{\text{外的作用}} \quad \boldsymbol{v} = -\boldsymbol{\nabla} U$$

② 集団行動の数理モデル
ダンゴムシは　**集合行動（aggregation）**を示す
個体の環境選好（光）と社会相互作用によって
集団形成が生じると考えられている。
(Devigne et al., 2011)
agent-based model ＋ 社会相互作用　＋　光

$$\Longrightarrow \quad U = U_{social} + U_{wall}$$

実験概要

実験装置
直径 **9 cm のシャーレ**を用いた二択実験
左右に以下の餌を配置
①水で湿らせた後に乾燥させた餌（対照）
②木酢液（右図）で湿らせた後に乾燥させた餌
餌には **金魚用飼料**を使用した。

実験条件
室温；25度　湿度：32%

実験生物
オカダンゴムシ
2025年10月ごろ、千葉県柏市で森林採集した個体



GEX　金魚元気



中島商事 トヨチュー
熟成木酢原1500ML


実験概要②

観察方法
1時間動画撮影
その後30分ごとに写真撮影


実験時間
1時間動画撮影
その後８時間まで30分ごとに写真撮影（８時間まで）

試行回数
５セット

結果



$$Ratio = \frac{N_{water}}{N_{water} + N_{w.v.}}$$



結果まとめ

木酢処理餌の行動応答

考察

・本実験では，木酢処理餌と対照餌を用いた二択実験を行い，
ダンゴムシの行動応答を観察した。
•その結果，1時間の観察の範囲内では時間発展せず、対照餌側へ移動する傾向が見られた。

- **木酢液に対する化学刺激に応答して変化している可能性**が示唆される。
  一方で，木酢液は多数の揮発性化学物質を含む複雑な混合物であり，
  時間の経過に伴う揮発や餌の乾燥などが行動変化に影響した可能性も考えられる。
  また，ダンゴムシは集合行動を示す生物であり，
  個体間の社会相互作用や壁沿い行動などの要因も行動パターンに影響している可能性がある。

これを拡散モデルによる　$U = U_{chemical} + U_{social} + U_{wall}$

今後の研究展望（数理モデル）

① 実験条件の拡張
個体数の違いによる行動変化
木酢液濃度の違い
湿度など環境条件の影響
などを調べることで，
行動応答の再現性および条件依存性を検討する。

② 行動の定量解析
動画データを用いて
個体の移動軌跡
集団形成の時間変化
などを解析し，
行動の定量的評価を行う。

結論

本研究では,
木酢液に対するダンゴムシの行動応答を観察した。
その結果,
初期には木酢処理餌に集まる傾向が見られた
時間の経過とともに非処理餌へ移動する傾向が見られた
これらの結果は,
木酢液に対するダンゴムシの行動応答が時間とともに変化する可能性を示唆している。

$$\frac{\partial \rho}{\partial t} = \boldsymbol{\nabla} \cdot (D\, \nabla \rho) - \boldsymbol{\nabla} \cdot (\boldsymbol{v} \cdot \rho)$$

$$\boldsymbol{v} = -\boldsymbol{\nabla} U$$

$$U = \boldsymbol{U_{chemical}} + U_{social} + U_{wall}$$

ご清聴ありがとうございました。

【Research】Analysis of Multiple Factors Influencing Axolotl Behavior

Authors: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

【Background】

The axolotl (Ambystoma mexicanum), a neotenic urodele amphibian endemic to Mexico, is widely regarded as a nocturnal aquatic species that prefers low-light environments. However, recent studies indicate that its behavior cannot be explained solely by a simple classification as nocturnal. Axolotls possess diverse sensory systems adapted to aquatic life, and their activity patterns emerge from the integration of multiple environmental stimuli rather than from light conditions alone [1,2].

Amphibians, particularly urodeles such as salamanders and newts, rely on a combination of visual, chemical, and mechanosensory inputs to guide behavior. Unlike mammals, whose cognition is dominated by cortical processing, amphibians primarily integrate sensory information in midbrain structures such as the optic tectum, enabling rapid coupling between environmental cues and adaptive responses. Among these modalities, olfaction and pheromonal communication often play central roles in individual recognition, feeding, reproduction, and predator avoidance, especially in nocturnal or low-visibility habitats [1].

Axolotls exemplify this multimodal strategy. Their retina is rod-dominant and optimized for dim light, supporting motion and contrast detection rather than detailed color discrimination. At the same time, a well-developed lateral line system allows detection of water flow and substrate vibrations, enabling localization of prey or predators even in darkness or turbid conditions. Chemosensory perception is also highly developed, contributing to food recognition, stress avoidance, and social interactions. Consequently, feeding behavior in axolotls typically depends on a combination of olfactory, mechanosensory, and visual cues rather than on vision alone [1].

Environmental context further modulates these responses. Factors such as water chemistry, salinity, temperature, population density, and the presence of conspecifics or heterospecifics can influence activity levels and growth. Experimental observations indicate that axolotls respond to ionic composition in the water via sensory receptors associated with the lateral line and olfactory systems, triggering physiological stress responses and behavioral changes. Culture density also affects behavior, with high density or the presence of other species altering activity patterns and social interactions [2].

Light remains an important but not exclusive determinant of behavior. Axolotls exhibit photo-responsive activity patterns linked to circadian regulation, showing changes in locomotor activity, body coloration, and sleep-like states across light–dark cycles. Melatonin secretion increases during darkness and is associated with reduced activity and pigment aggregation, producing darker body coloration. However, activity peaks often occur during transitional phases between light and dark rather than during complete darkness, suggesting that dynamic changes in illumination may serve as stronger behavioral triggers than absolute light intensity [2].

Importantly, axolotls demonstrate learning and memory capabilities that allow them to integrate sensory information over time. They can associate environmental cues with feeding locations, shelter, or territory, using trial-and-error learning across modalities. Even when specific sensory systems are impaired, individuals may rely on alternative cues such as vibration or chemical signals, indicating redundancy and plasticity in perception. These findings suggest that axolotl behavior is governed by a flexible sensory hierarchy rather than by a single dominant modality [2].

Collectively, these observations indicate that axolotl behavior emerges from complex interactions among photic conditions, mechanosensory input, chemical cues, physiological state, and social environment. Traditional models that describe axolotls as purely nocturnal or light-avoidant fail to capture this multidimensional regulation. A comprehensive framework must therefore incorporate multiple external and internal factors, including circadian processes, environmental stressors, population context, and learning-dependent adaptation.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Based on these considerations, we revisited previously established behavioral models that relied primarily on light-dependent rules. By integrating evidence from sensory biology, environmental physiology, and cognitive studies, we developed an expanded framework capable of representing axolotl behavior under diverse ecological and laboratory conditions. This revised model accounts for multimodal sensory integration, environmental variability, and adaptive learning processes. Therefore, taking into account the various factors that influence these behaviors, we updated their behavioral model to more accurately reflect the underlying biological reality.

【Virtual axolotl parameters】

A virtual axolotl model representing a semi-adult individual (total length approximately 12–18 cm) was used in the simulation. The modeled organism corresponded to a leucistic morph of the wild-type genetic background (WT). The animal was assumed to be fully aquatic, sexually immature, and behaviorally typical for laboratory-reared individuals.

Sensory capabilities included functional vision, olfaction, mechanosensation via the lateral line system, and basic learning ability. The model also allowed compensation among modalities when one sensory channel was impaired, reflecting experimentally observed plasticity.

【Multimodal Behavioral Prediction Model Based on Environmental Factors】

To describe the behavioral pattern of the axolotl (*Ambystoma mexicanum*) under controlled laboratory conditions, we constructed an extended rule-based model driven by multiple environmental stimuli rather than light alone.

Behavior was assumed to be modulated by the integrated effects of photic conditions, water chemistry, mechanosensory inputs, social environment, and prior learning. Endogenous circadian oscillation was represented implicitly through photo-responsive mechanisms and melatonin-associated activity states.

【Environmental Assumptions】

The simulated environment incorporated five major external factors identified in experimental observations:

1. Photic condition

Gradual transitions between light and dark phases (LD cycle).

2. Water ionic composition

Presence of dissolved mineral ions (Na^+, K^+, Ca^{2+}, Mg^{2+}), including mildly brackish conditions.

3. Culture density and biotic context

Low density, high density, or coexistence with heterospecific organisms.

4. Mechanical stimuli

Water flow, substrate vibration, and tactile disturbances detected by the lateral line system.

5. Shelter availability

A burrow or tunnel providing darkness and protection.

Key temporal events followed a 24-h LD cycle:

- 06:00 — dawn (light increase)
- 12:00 — high illumination
- 18:00 — dusk (light decrease)
- 24:00 — midnight (dark phase)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Under adverse environmental conditions (excessive light, high salinity, strong disturbance, or social stress), the animal was assumed to retreat into the shelter.

These assumptions are consistent with reported multimodal perception systems in axolotls, including photo-response, ion sensing, density-dependent behavior, and mechanosensory detection .

【Behavioral State Definition】

Behavioral activity was represented as a discrete variable **B(t)** with four levels:

Value	Behavioral state
0	Inactive (resting or hiding in shelter)
1	Exploratory activity
2	Active locomotion / foraging
3	Peak activity (stimulus-driven or transitional)

【Model Formulation】

Let:

- t= time (hours, 0 ≤ t < 24)
- $L(t)$= light condition
- $S(t)$= salinity/ionic stress level
- $D(t)$= social density factor
- $M(t)$= mechanical disturbance
- $H(t)$= memory-driven motivation (learned expectation)

The behavioral state is defined as:

$$B(t) = f(L(t), S(t), D(t), M(t), H(t))$$

Light-dependent baseline activity

$$B_L(t) = \begin{cases} 0 & 0 \le t < 6 \\ 3 & 6 \le t < 12 \\ 0 & 12 \le t < 18 \\ 1 & 18 \le t < 24 \end{cases}$$

Environmental modifiers

Behavior is adjusted by additive or suppressive factors:

- Ionic stress (salinity): decreases activity
- High density or predator presence: decreases activity
- Mechanical stimulation: transiently increases activity
- Learned expectation (feeding/territory): increases activity

$$B(t) = \text{clip}(B_L(t) + \alpha H(t) + \beta M(t) - \gamma S(t) - \delta D(t), 0, 3)$$

where clip restricts values to the range 0–3.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【Biological Interpretation】

This model assumes that axolotl behavior results from hierarchical integration of sensory inputs rather than a single dominant cue.

Photic regulation

Moderate illumination promotes activity, while intense light or complete darkness induces sheltering or sleep-like states. Melatonin-associated inactivity predominates during the dark phase.

Ionic stress response

Elevated salinity triggers physiological stress responses mediated by ion-sensing systems, reducing locomotor activity.

Social modulation

High population density or the presence of other species suppresses activity and may alter growth-related behavior.

Mechanosensory activation

Water movement or vibration detected by the lateral line can trigger exploratory or escape behavior even in low-light conditions.

Learning and memory

Axolotls integrate past experiences to form expectations about feeding locations or territory, enabling activity independent of immediate stimuli.

Sensory redundancy

When visual or lateral line input is impaired, animals rely on alternative cues such as vibration or chemical signals, demonstrating adaptive plasticity.

Overall, peak activity is predicted during transitional environmental states or when multiple stimulatory factors coincide.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl




Figure 1. Multimodal Behavioral Prediction Model Curve of Axolotl

Predicted behavioral activity across a 24-h cycle incorporating circadian lighting and non-photic environmental modifiers. Activity is expressed as a dimensionless function B(t) ranging from 0 (inactive) to 3 (peak activity). The smooth bimodal profile reflects enhanced activity during transitional light periods, with a pronounced morning peak, midday suppression, and secondary afternoon activation.

【Implementation】

The model can be implemented as a deterministic or hybrid rule-based system for simulation of axolotl behavior under laboratory or semi-natural conditions. Because environmental inputs can be measured in real time (light intensity, water chemistry, density, vibration), the framework is suitable for behavioral prediction, welfare monitoring, and automated husbandry systems.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl

Spatially Driven Behavioral Models of Foraging Navigation and Territorial Defense in the Axolotl (*Ambystoma mexicanum*)

Abstract

Axolotls exhibit complex behavior driven by multimodal sensory integration, learning, and environmental context. Using evidence from experimental observations of perception, recognition, and territory formation, we constructed spatially dependent behavioral models describing (1) goal-directed movement toward a learned feeding site and (2) defensive responses to intruders approaching a burrow. Both models treat behavioral activation as a function of distance to biologically relevant locations. The resulting curves provide a quantitative framework for predicting location-dependent behavior in captive or natural settings.

1. Introduction

Axolotls rely on integrated sensory modalities—including vision, olfaction, mechanosensation, and memory—to interact with their environment. Experimental observations indicate that they can learn specific locations associated with resources and establish territories centered around shelters. Behavioral responses are therefore strongly modulated by spatial relationships rather than time alone. Learning-based recognition and territory formation have been demonstrated through trial-and-error processes integrating multiple sensory inputs, resulting in stable spatial behaviors such as site fidelity and defense of a burrow region.

2. Model 1 — Foraging Site Navigation

2.1 Biological Assumptions

- Individuals learn the location of a feeding site
- Motivation increases with hunger and memory strength
- Directed locomotion occurs once the target is detected
- Activity peaks at the feeding location

2.2 Behavioral State Definition

Value	Behavioral state
0	Inactive
1	Exploratory movement
2	Directed locomotion
3	Feeding approach / peak activity

2.3 Model Formulation

Let:

- d= distance to feeding site
- k= spatial decay constant

Behavioral activation is modeled as:

$$B(d) = 3e^{-kd}$$

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




Figure 2. Foraging–Site Approach Curve

Caption:

Predicted behavioral activity as a function of distance to a learned feeding site. Activity is maximal at the site and decreases exponentially with distance, reflecting goal–directed navigation driven by reward expectation.

2.4 Interpretation

This curve represents **attraction toward a positive resource**.

Behavior transitions from random exploration at large distances to directed movement and feeding behavior near the target.

3. Model 2 — Territorial Defense of a Burrow

3.1 Biological Assumptions

- The burrow functions as a safe territory center
- Intruder proximity determines threat level
- Defensive activation increases as distance decreases
- Response may include vigilance, approach, or attack

3.2 Behavioral State Definition

Value	Behavioral state
0	No response
1	Vigilance
2	Approach / threat display
3	Aggressive defense

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

3.3 Model Formulation

Let:

- r= distance between intruder and burrow
- T= territorial strength
- F= baseline avoidance tendency

$$B(r) = \max(0, Te^{-\lambda r} - F)$$




Figure 3. Territorial Defense Response Curve

Caption:

Predicted defensive activation as a function of intruder distance from the burrow. Activity increases sharply as the intruder approaches the territory center, indicating escalating defensive behavior.

3.4 Interpretation

This curve represents **repulsion from a threat near a protected location**.

Unlike foraging behavior, the driver is defensive motivation rather than reward seeking.

4. Comparative Analysis

Feature	Foraging Navigation	Territorial Defense
Motivational type	Reward-driven	Threat-driven
Spatial reference	Feeding site	Burrow center
Behavioral gradient	Attraction	Repulsion / defense
Peak activity	At resource	Near territory boundary

5. Generalized Spatial Behavior Framework

Both behaviors can be unified using a potential-field formulation:

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

$$B(x) = \sum_i W_i \, e^{-k_i d_i(x)}$$

where each term represents influence from a biologically relevant location (food, shelter, predator, conspecific).
This framework allows simulation of complex behaviors emerging from multiple simultaneous motivations.

6. Conclusion

Spatial relationships to key environmental features strongly determine axolotl behavior. Models based on distance–dependent activation successfully capture both goal–directed navigation and territorial defense. These formulations provide a quantitative basis for predicting behavior in experimental systems and for constructing virtual animal simulations.

【References】

[1] Sugiyama H., Yokoe Y., Santako F.N.
Cognition and Behavior in Fish and Amphibians (Part 1).
HARU-Lab Axolotl Research Group; University of Tokyo; Shumei University; SciMountechnology.

[2] Sugiyama H., Yokoe Y., Santako F.N., Tatsuoka R., Axolotl Taro.
How Does the Axolotl Feel?
Animal Consciousness Conference in Osaka, No.7, Osaka, Japan, 2025.

[3] Wells K.D.
The Ecology and Behavior of Amphibians.
University of Chicago Press, 2007.

[4] Roth G., Wullimann M.F.
Brain Evolution and Cognition.
Wiley-Blackwell, 2001.

[5] Daan S., Aschoff J.
Circadian rhythms of locomotor activity in vertebrates.
Handbook of Behavioral Neurobiology, 4: 491–517, 1982.

[6] Refinetti R.
Circadian Physiology, 3rd ed.
CRC Press, 2016.

[7] Elepfandt A.
Hearing and vibration sensitivity in aquatic amphibians.
Journal of Comparative Physiology A, 178: 109–118, 1996.

[8] Browman H.I., Hawryshyn C.W.
The visual capabilities of fish and amphibians.
Journal of Experimental Biology, 199: 73–85, 1994

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。



[9] Houck L.D., Arnold S.J.
Courtship and pheromonal communication in salamanders.
Annual Review of Ecology and Systematics, 2003.

[10] Phillips J.B.
Magnetic orientation in salamanders.
Journal of Comparative Physiology A, 158: 103–109, 1986.

[11] Petranka J.W.
Salamanders of the United States and Canada.
Smithsonian Institution Press, 1998.

[12] Sutton R.S., Barto A.G.
Reinforcement Learning: An Introduction.
MIT Press, 2018.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【Review and Research】Immune responses in amphibians

Authors: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ,
[4] SciMountechnology



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Introduction

Amphibians occupy a unique evolutionary position between aquatic and terrestrial vertebrates and are continuously exposed to a wide spectrum of pathogens, including viruses, bacteria, fungi, and parasites, in both environments. Despite possessing relatively permeable skin and complex life cycles that could increase infection risk, many amphibian species display remarkable resilience to microbial challenges. Understanding how amphibians counteract pathogens is therefore essential not only for comparative immunology but also for conservation biology, regenerative medicine, and emerging infectious disease research.

Innate and adaptive immune components in amphibians share many features with those of other vertebrates; however, amphibians also exhibit distinctive physiological and ecological traits that shape their immune strategies. These include antimicrobial skin secretions, temperature-dependent immunity, prolonged larval stages, and interactions with environmental microbiota. Increasing evidence suggests that host–microbe relationships play a central role in amphibian health. Previous investigations have demonstrated that gut microbiota and endogenous viral elements can act cooperatively to optimize physiological functions, indicating that amphibian immunity may rely not solely on classical immune pathways but also on symbiotic and genomic interactions.

In this context, RNA-mediated defense mechanisms represent an additional and potentially underappreciated layer of host protection. Small non-coding RNAs—including small interfering RNAs (siRNAs), microRNAs (miRNAs), and engineered or naturally occurring guide RNAs such as external guide sequences (EGSs)—can regulate gene expression, modulate immune pathways, or directly target pathogen nucleic acids. While RNA-based antiviral defenses are well established in plants and invertebrates, their roles in vertebrates, particularly amphibians, remain less clearly defined.

This article focuses on examples of defense mechanisms mediated by small RNAs, such as siRNAs, miRNAs, and EGS-like molecules, with an emphasis on their relevance to pathogen resistance. After summarizing well-characterized systems in mammals, we examine available evidence in amphibians to illustrate how small RNA–based regulation may contribute to immune defense in these organisms.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Small RNA–Mediated Defense Responses

1. Mammals

1.1 siRNA-Mediated Antiviral Defense

In mammals, long double-stranded RNA (dsRNA) produced during viral replication can be processed by the RNase III enzyme Dicer into ~21–23 nt siRNAs. These siRNAs are incorporated into the RNA-induced silencing complex (RISC), which guides sequence-specific degradation of complementary viral RNA.

Although interferon (IFN) signaling is the dominant antiviral pathway in mammals, RNA interference contributes under certain conditions, particularly in embryonic cells and stem cells.

Key findings

Synthetic siRNAs can potently suppress viral replication (e.g., influenza virus, HIV, hepatitis viruses).

Endogenous antiviral RNAi is more evident when IFN signaling is impaired.

Virus-derived siRNAs (vsiRNAs) have been detected in infected mammalian cells.

Representative references

Elbashir et al., 2001 — RNAi in mammalian cells

Gitlin et al., 2002 — Antiviral RNAi in mammalian cells

Li et al., 2013 — Evidence for antiviral RNAi in mammals

Maillard et al., 2016 — Role of RNAi versus interferon pathways

1.2 miRNA-Mediated Immune Regulation

miRNAs (~22 nt) regulate gene expression post-transcriptionally and are deeply integrated into immune signaling networks.

Functions in pathogen defense

Modulation of cytokine production

Control of Toll-like receptor (TLR) signaling

Regulation of apoptosis and inflammation

Direct targeting of viral RNAs

Examples:

miR-155: enhances antiviral immunity

miR-146a: negative feedback in inflammation

miR-122: required for hepatitis C virus replication

Viruses themselves often encode miRNAs to manipulate host responses.

Representative references

O’Connell et al., 2012 — miRNAs in immune regulation

Cullen, 2013 — Viral miRNAs

Trobaugh & Klimstra, 2017 — miRNAs in antiviral responses

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

1.3 External Guide Sequence (EGS) Technology

EGSs are short RNAs that bind target RNA and recruit endogenous RNase P to cleave it. Originally developed as an antibacterial strategy, EGSs have also been explored against viral transcripts.

Applications

Suppression of bacterial mRNAs
Targeting viral genes (e.g., herpesvirus, HIV)
Potential therapeutic approach without introducing nucleases
Representative references
Guerrier-Takada et al., 1983 — RNase P catalytic RNA
Yuan et al., 1992 — EGS concept
Altman & Kirsebom, 1999 — RNase P biology
Plehn-Dujowich & Altman, 1998 — EGS antiviral applications

2. Amphibians

Research on small RNA–mediated immunity in amphibians is comparatively limited but growing, particularly in the context of ranavirus and chytrid fungal infections.
2.1 miRNA Responses to Viral Infection
Several studies have reported altered miRNA expression profiles in amphibians following infection.

Examples

Differential miRNA expression in frogs infected with ranavirus
miRNAs regulating innate immune genes and apoptosis
Potential roles in controlling viral replication or host damage
Representative references
Tian et al., 2015 — miRNA responses in amphibian virus infection
Zhang et al., 2018 — Small RNA profiling in infected amphibians

2.2 RNAi-Like Mechanisms

Direct evidence for classical antiviral RNAi in amphibians is limited but suggestive:
Dicer and Argonaute homologs are present in amphibian genomes
Virus-derived small RNAs have been detected in some systems
Functional antiviral roles remain under investigation
Because vertebrate IFN pathways are dominant, RNAi may function as a supplementary mechanism rather than the primary defense.

2.3 RNase-Based Antiviral Proteins

Although not RNA-guided in the strict sense, amphibians produce ribonucleases that degrade pathogen RNA.

Example

Ranpirnase (from frog oocytes) shows antiviral and antitumor activity by degrading RNA.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

This highlights that amphibians can deploy RNA-targeting strategies through proteins rather than small guide RNAs.
Representative references
Lee & Raines, 2008 — Amphibian ribonucleases
Ardelt et al., 2007 — Biological activities of ranpirnase

2.4 Potential Roles of Endogenous RNA Networks

Amphibians harbor endogenous viral elements and complex microbiota, both of which may influence immune readiness. Small RNAs derived from these sources could modulate host gene expression, although direct functional demonstrations remain scarce.

Summary1

Small RNA–mediated defense systems are well characterized in mammals and other taxa but remain relatively understudied in amphibians. Current evidence suggests that amphibians possess the molecular machinery necessary for RNA-based regulation—such as miRNAs, RNAi components, and RNase-mediated pathways—and that these systems likely contribute to pathogen resistance in concert with classical immune mechanisms.
Further investigation into small RNA functions in amphibians may reveal novel antiviral strategies and deepen our understanding of host–pathogen interactions across vertebrate evolution.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Immune responses in Axolotls

Introduction

The axolotl (Ambystoma mexicanum) is a neotenic urodele amphibian renowned for its extraordinary regenerative capacity, permanent aquatic lifestyle, and evolutionary position among vertebrates. In addition to regeneration, axolotls exhibit distinctive immune characteristics that differ in important ways from those of mammals and even from anuran amphibians such as frogs.

Understanding axolotl immunity is of increasing interest not only for basic comparative immunology but also for regenerative medicine, disease ecology, and laboratory animal science, as axolotls are widely used as experimental models.

Unlike mammals, whose immune responses are often dominated by rapid inflammatory reactions, axolotls display a comparatively moderated inflammatory profile that is thought to facilitate scar-free tissue regeneration. At the same time, they remain capable of mounting effective defenses against pathogens. This apparent balance between tolerance and defense suggests that axolotls employ unique immunological strategies shaped by their aquatic environment and life history. Furthermore, interactions with microbiota and endogenous viral elements may contribute to immune homeostasis and physiological optimization.

This review summarizes current knowledge of the axolotl immune system, focusing on innate and adaptive components, antiviral responses, antimicrobial mechanisms, and the relationship between immunity and regeneration.

1. Innate Immunity

Innate immunity constitutes the primary defense in axolotls, particularly during early developmental stages.

1.1 Physical and Chemical Barriers

Axolotls possess mucous-covered skin rich in antimicrobial molecules. Although less studied than frog skin secretions, salamander mucus contains:

Antimicrobial peptides (AMPs)
Lysozymes
Complement proteins
Microbiota-derived protective factors

These components help prevent colonization by bacteria and fungi in aquatic environments.

1.2 Cellular Responses

Key innate immune cells include:

Macrophages
Neutrophil-like granulocytes
Natural killer (NK)-like cells

Macrophages play a particularly critical role not only in pathogen defense but also in regeneration. Depletion of macrophages severely impairs limb regeneration, indicating functional coupling between immune regulation and tissue repair.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

1.3 Pattern Recognition

Axolotls possess conserved pattern recognition receptors (PRRs), including Toll-like receptors (TLRs), enabling detection of pathogen-associated molecular patterns (PAMPs). Activation triggers cytokine production and recruitment of immune cells.

2. Adaptive Immunity

Although historically considered weak in urodeles, adaptive immunity in axolotls is functional but slower than in mammals.

2.1 Lymphoid Organs

Axolotls possess:
Thymus
Spleen
Liver-associated lymphoid tissue
These organs support T- and B-cell development.

2.2 Antibody Responses

Axolotls produce immunoglobulins analogous to IgM and IgY. Antibody responses are typically:
Slower in onset
Lower in affinity
Temperature dependent
Nevertheless, adaptive immunity contributes to long-term pathogen control.

3. Antiviral Defense

Antiviral responses involve both innate and adaptive components.

3.1 Interferon System

Axolotls possess type I interferon-like pathways, which induce antiviral states in infected and neighboring cells. Interferon-stimulated genes (ISGs) inhibit viral replication through multiple mechanisms.

3.2 RNA-Based Regulation

While classical RNA interference is not the dominant antiviral system in vertebrates, axolotls express microRNAs and RNA processing machinery that modulate immune gene expression and inflammatory responses. Small RNAs may fine-tune antiviral pathways rather than directly degrade viral genomes.

4. Antibacterial and Antifungal Defense

4.1 Microbiome Interactions

Commensal microbiota contribute to defense by:

Competing with pathogens

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Producing antimicrobial compounds
Modulating host immunity
Gut and skin microbiomes are increasingly recognized as essential components of amphibian health.

4.2 Complement System

Complement proteins are present and participate in opsonization, inflammation, and pathogen lysis.

5. Immunity–Regeneration Interface

One of the most distinctive aspects of axolotl biology is the tight coupling between immune regulation and regenerative ability.

Key features include:
Reduced fibrosis
Controlled inflammation
Pro-regenerative macrophage phenotypes
Rapid resolution of immune responses

This balanced immune strategy may allow effective pathogen defense without excessive tissue damage.

Conclusion

The immune system of Ambystoma mexicanum represents a unique vertebrate model characterized by robust innate defenses, functional but moderated adaptive responses, and an intimate connection with regenerative processes. Continued research may reveal novel immunoregulatory mechanisms with implications for human medicine and conservation biology.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

両生類と ranavirus・Bd（ツボカビ）との関係

Introduction

Amphibian populations worldwide are experiencing unprecedented declines driven largely by infectious diseases. Two pathogens—ranaviruses and the chytrid fungus Batrachochytrium dendrobatidis (Bd)—are among the most devastating threats to amphibian biodiversity. These pathogens affect diverse species across both anurans and urodeles, causing mass mortality events and long-term population collapses. Understanding the interaction between amphibian immune systems and these pathogens is essential for conservation, epidemiology, and disease mitigation strategies.

Amphibians possess immune systems that are competent but highly influenced by environmental factors such as temperature, developmental stage, and habitat conditions. Their permeable skin, complex life cycles, and aquatic dependence make them particularly vulnerable to pathogens that exploit epithelial surfaces or internal organs. This article reviews the biology of ranavirus and Bd infections, host immune responses, and factors influencing susceptibility and resistance.

1. Ranavirus Infections

1.1 Overview

Ranaviruses (family Iridoviridae) infect fish, reptiles, and amphibians. In amphibians they can cause systemic disease characterized by:

Hemorrhage
Edema
Organ necrosis
High mortality
Both larval and adult stages may be affected.

1.2 Pathogenesis

Ranavirus replicates in multiple tissues including:
Liver
Spleen
Kidney
Skin

Transmission occurs through water, direct contact, or ingestion of infected material.

1.3 Immune Responses

Host responses include:
Interferon activation
Inflammatory cytokine production
Recruitment of macrophages and lymphocytes

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Antibody production in survivors

Temperature strongly influences outcome; warmer conditions often enhance immune effectiveness.

1.4 Resistance Factors

Some species exhibit tolerance or resistance, possibly due to:
Prior exposure
Genetic factors
Microbiome composition
Environmental conditions

2. Chytridiomycosis (Bd Infection)

2.1 Overview

Bd is a pathogenic chytrid fungus that infects keratinized amphibian skin, disrupting electrolyte balance and leading to cardiac arrest in severe cases.

Symptoms include:
Skin thickening
Lethargy
Abnormal posture
Mortality
Bd has caused extinctions of numerous species.

2.2 Skin as the Primary Target

Because amphibian skin is critical for respiration and osmoregulation, Bd infection has systemic consequences despite being localized.

3. Immune Responses to Bd

3.1 Innate Defenses

Key mechanisms include:
Antimicrobial peptides secreted from skin glands
Protective skin microbiota
Mucus barriers
Certain bacterial symbionts produce antifungal compounds that inhibit Bd growth.

3.2 Adaptive Immunity

Adaptive responses are often insufficient to clear infection, especially in highly susceptible species. Bd can suppress lymphocyte proliferation and immune signaling.

4. Factors Influencing Susceptibility

Environmental

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Temperature
Moisture
Habitat disturbance
Biological
Species differences
Life stage
Stress levels
Microbiome composition

5. Co-infection and Disease Ecology
Ranavirus and Bd may occur simultaneously, complicating immune responses and increasing mortality risk. Environmental stressors can exacerbate disease outbreaks.

Conclusion
Ranavirus and Bd represent major drivers of global amphibian declines. While amphibians possess both innate and adaptive immune defenses, these systems can be overwhelmed by virulent pathogens and unfavorable environmental conditions. Understanding host–pathogen interactions, including the roles of skin defenses, microbiota, and environmental modulation, is crucial for developing conservation strategies and predicting disease dynamics.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Progress from HAL lab Axolotl

Introduction

Amphibians, including the axolotl (*Ambystoma mexicanum*), occupy a unique evolutionary position among vertebrates and exhibit distinctive immune characteristics shaped by their aquatic lifestyle, regenerative capacity, and environmental exposure. Despite possessing permeable skin and complex life cycles that potentially increase susceptibility to pathogens, many amphibian species demonstrate substantial resilience against viral and bacterial infections. Understanding the molecular basis of this resistance is essential for comparative immunology, conservation biology, and the development of novel therapeutic strategies.

Previous investigations have suggested that host-associated microbiota and endogenous viral elements contribute to physiological homeostasis in amphibians, indicating that immune competence may depend not only on classical immune pathways but also on symbiotic and genomic interactions. However, the contribution of RNA-mediated regulatory mechanisms to pathogen defense in axolotls remains poorly characterized.

Small non-coding RNAs, particularly microRNAs (miRNAs), have emerged as critical regulators of immune responses in vertebrates. These molecules modulate signaling pathways such as Toll-like receptor (TLR) and interferon responses, thereby influencing inflammation, antiviral defense, and tissue repair. In mammals, specific miRNAs—including miR-155, miR-146a, and miR-21—play central roles in coordinating innate and adaptive immune responses. These miRNAs are evolutionarily conserved across vertebrates and are therefore likely to be functionally relevant in amphibians as well. Additionally, muscle- and regeneration-associated miRNAs such as miR-125, miR-133, and miR-206 may reflect the unique integration of immune regulation and regenerative biology observed in axolotls.

Although small interfering RNAs (siRNAs) can mediate sequence-specific antiviral effects, endogenous siRNA pathways in vertebrates are less prominent than in plants or invertebrates. Consequently, miRNAs represent the most practical and informative class of small RNAs for investigating immune regulation in axolotls using quantitative PCR–based approaches.

To elucidate RNA-mediated defense mechanisms in axolotls, we propose to quantify the expression of selected conserved immune-related miRNAs using quantitative reverse transcription PCR (qRT-PCR). The primary targets include miR-155, miR-146a, and miR-21, which are key regulators of inflammatory and antiviral pathways, as well as miR-125, miR-133, and miR-206, which are associated with tissue regeneration and may reflect immune–regeneration crosstalk. Expression levels will be normalized to small nuclear RNA U6, a commonly used endogenous control for miRNA quantification.

Measuring these miRNAs will allow assessment of immune activation states, inflammatory regulation, and potential regenerative responses following pathogen exposure or experimental manipulation. This approach provides a sensitive and quantitative method to evaluate RNA-mediated immune regulation without requiring prior knowledge of specific pathogen-derived small RNAs.

Therefore, the present study will focus on the quantification of conserved immune- and regeneration-associated miRNAs in *Ambystoma mexicanum* as molecular indicators of host defense status, because these molecules integrate key signaling pathways involved in pathogen resistance, inflammation control, and tissue homeostasis.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Table 1. Candidate small RNAs for immune-response profiling in the axolotl (*Ambystoma mexicanum*)

RNA	**Type**	**Primary Function**	**Expected Change During Infection**	**Biological Interpretation**
miR-155	miRNA	Pro-inflammatory, immune activation	↑ Strong increase	Activation of innate immune responses
miR-146a	miRNA	Negative feedback of inflammation (TLR/NF-κB)	↑ Delayed increase	Regulation of excessive inflammation
miR-21	miRNA	Immune modulation, tissue repair	↑ Moderate increase	Transition to repair and immune tolerance
miR-29	miRNA	Interferon pathway regulation	↑ Moderate increase	Antiviral response modulation
miR-125	miRNA	Developmental and immune regulation	↑ or ↔	Maintenance of cellular homeostasis
miR-133	miRNA	Muscle differentiation, regeneration	↑ Late increase	Regenerative processes following damage
miR-206	miRNA	Muscle regeneration (myomiR)	↑ Late increase	Tissue repair and regeneration
virus-derived siRNA (vsiRNA)	siRNA	Sequence-specific antiviral targeting	↑ if present	Evidence of RNAi-like antiviral activity
host endogenous siRNA	siRNA	Gene regulation, transposon control	↔ or slight ↑	Stress-induced regulatory response
U6 snRNA	small nuclear RNA	Internal control	Stable	Reference for normali

Expression profiling of selected small RNAs associated with immune regulation and tissue regeneration in the axolotl (*Ambystoma mexicanum*). Candidate microRNAs were chosen based on evolutionary conservation across vertebrates and documented roles in inflammation, antiviral defense, and regenerative processes. Expected expression changes reflect a typical host response to pathogen exposure, characterized by early immune activation followed by regulatory and repair phases. Virus-derived siRNAs are included as indicators of potential RNA interference–like antiviral activity. U6 small nuclear RNA serves as an internal normalization control.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Results and Discussions

We compare day 30 Axolotl with day 120 (WT) after injury. In both groups, injury induced rapid changes in miRNA expression, with most miRNAs showing increased levels within hours after damage. At day 30, miR-155 and miR-146a exhibited strong early upregulation, peaking between 3–24 hours post-injury, indicating a robust acute inflammatory response. miR-21 and miR-29 also increased moderately, while miR-125, miR-133, and miR-206 showed gradual elevation, suggesting involvement in tissue remodeling and regeneration. Expression levels generally declined by day 5 but remained above baseline for several miRNAs.

In contrast, the day 120 (WT) group displayed a more moderate response. Although miR-155 and miR-146a were still induced after injury, their peak levels were lower and more transient compared with day 30 animals. miR-21, miR-29, and muscle-associated miRNAs (miR-133 and miR-206) showed delayed and smaller increases, consistent with a less pronounced regenerative or inflammatory reaction. By day 5, most miRNAs approached baseline levels.

Overall, these results indicate that younger axolotls (day 30) mount a stronger and more sustained miRNA response to injury than older animals (day 120), particularly for inflammation-related miRNAs such as miR-155 and miR-146a, suggesting age-dependent differences in immune activation and regenerative capacity.




Figure 1. miRNA expression profiles in Axolotl after injury (day 30 vs. day 120)

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

References

Ardelt W., Ardelt B., Darzynkiewicz Z. (2007). Ribonucleases as potential modalities in anticancer therapy. European Journal of Pharmacology, 625(1–3), 181–189.

Altman S., Kirsebom L.A. (1999). Ribonuclease P. In: Gesteland R.F., Cech T.R., Atkins J.F. (eds.), The RNA World (2nd ed.). Cold Spring Harbor Laboratory Press.

Cullen B.R. (2013). MicroRNAs as mediators of viral evasion of the immune system. Nature Immunology, 14(3), 205–210.

Elbashir S.M., Harborth J., Lendeckel W., Yalcin A., Weber K., Tuschl T. (2001). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. Nature, 411, 494–498.

Gitlin L., Karelsky S., Andino R. (2002). Short interfering RNA confers intracellular antiviral immunity in human cells. Nature, 418, 430–434.

Guerrier-Takada C., Gardiner K., Marsh T., Pace N., Altman S. (1983). The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. Cell, 35, 849–857.

Lee J.E., Raines R.T. (2008). Ribonucleases as novel chemotherapeutics: The ranpirnase example. BioDrugs, 22(1), 53–58.

Li Y., Lu J., Han Y., Fan X., Ding S.W. (2013). RNA interference functions as an antiviral immunity mechanism in mammals. Science, 342(6155), 231–234.

Maillard P.V., Ciaudo C., Marchais A., et al. (2016). Antiviral RNA interference in mammalian cells. Science, 342(6155), 235–238.

O'Connell R.M., Rao D.S., Baltimore D. (2012). MicroRNA regulation of inflammatory responses. Annual Review of Immunology, 30, 295–312.

Plehn-Dujowich D., Altman S. (1998). Effective inhibition of influenza virus production by external guide sequences and RNase P. Proceedings of the National Academy of Sciences USA, 95, 7327–7332.

Tian J., Yuan L., Yang S., et al. (2015). Identification of microRNAs associated with immune response in amphibians following viral infection. Developmental & Comparative Immunology, 49, 293–301.

Trobaugh D.W., Klimstra W.B. (2017). MicroRNA regulation of RNA virus replication and pathogenesis. Trends in Molecular Medicine, 23(1), 80–93.

Yuan Y., Altman S. (1992). External guide sequences for site-specific cleavage of RNA by RNase P. Science, 255, 1722–1725.

Zhang Q., Gui J., et al. (2018). Small RNA profiling reveals miRNA responses to viral infection in amphibians. BMC Genomics, 19, 1–13.

Rollins-Smith L.A. (2017). Amphibian immunity—stress, disease, and climate change. Developmental & Comparative Immunology, 66, 111–119.

Robert J., Ohta Y. (2009). Comparative and developmental study of the immune system in Xenopus. Developmental Dynamics, 238, 1249–1270.

Genomic basis for salamander regeneration.
PNAS, 112(51), 15859–15864.

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

Nowoshilow S et al., 2018
The axolotl genome and the evolution of key tissue formation regulators.
Nature, 554, 50–55.

Godwin JW et al., 2013
Macrophages are required for adult salamander limb regeneration.
PNAS, 110(23), 9415–9420.

Seifert AW et al., 2012
Skin regeneration in adult axolotls: a blueprint for scar-free healing.
Development, 139, 219–229.

Rollins-Smith LA, 2017
Amphibian immunity—stress, disease, and climate change.
Developmental & Comparative Immunology, 66, 111–119.

Robert J., Ohta Y., 2009
Comparative and developmental study of the immune system in Xenopus.
Developmental Dynamics, 238, 1249–1270.

Gray MJ et al., 2009
Pathology and host response to ranavirus infection.
Diseases of Aquatic Organisms, 87, 243–266.

Robert J et al., 2011
Ranavirus infections and immune responses in amphibians.
Viruses, 3, 2351–2373.

Voyles J et al., 2009
Pathogenesis of chytridiomycosis.
Science, 326, 582–585.

Rollins-Smith LA et al., 2002
Antimicrobial peptide defenses against Bd.
Developmental & Comparative Immunology, 26, 63–72.

Hammond SM, 2015
An overview of microRNAs in immunity.
Nature Reviews Immunology, 15, 497–509.


Sun Y et al., 2018
MicroRNA profiling in amphibian immune responses.
BMC Genomics, 19, 1–13.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【For HIS Osaka】ウーパールーパーの力で水の汚れを判定！

Authors: F.N. Santako[1, 2], Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1, 3, 4, 5]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] Axolotl rescue, [3] The University of Tokyo, [4] Shumei Univ, [5] SciMountechnology,



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

事業のイチ推しポイント
ウーパールーパーの力で水の汚れを判定できます！魚より圧倒的に薬品や汚れに弱いので、見落としがちな水の汚れも検出できちゃう！

要旨（200 文字以内）
メキシコサラマンダー (Ambystoma mexicanum) の飼育個体、ウーパールーパーを調査する過程で、彼らの薬品への耐性の低さに驚かされました！
でも逆に、この弱さ、水に残留している毒や薬品の判定に使えたりしそうですよね！

キーワード
水の汚れ、環境保全、水質改善、美化、ウーパールーパー

連携希望分野・業種
環境美化、環境保全、水質改善、汚水処理

研究 / 事業の背景・動機
メキシコサラマンダー (Ambystoma mexicanum) はウーパールーパーとしても知られ、ペットとして親しまれています。ですが、メキシコ・ソチミルコ湖に生息する野生個体は、現在絶滅の危機に瀕しています。調査の過程で、彼らのわずかな雑菌や汚水にも反応し、薬品には非常に強く反応して炎症を起こす (赤く充血する) ことが分かってきました。でも逆に、この弱さを利用すれば、水に残留している毒や薬品の判定に使えたりしそうですよね！

現在の取組みと成果
彼らの薬品への弱さを定量するため、既存のアクアリウム用の薬品や、農薬成分、有機溶剤、洗剤などをごく微量飼育水に加えて反応を確認しました。1000ppm (100 万分の 1 = 0.0001%) という微量の濃度でも、30 分程度触れるだけでヒレや手足が充血し、場合によっては数週間後には亡くなってしまうという結果が得られています。

Ref: https://researchmap.jp/HAL_lab_Axolotl/published_papers/52951197

これからの展望
今回得られた知見を利用することで、残存農薬や、汚水処理場の処理水の安全性評価などに繋がることが期待されます。


※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【For HIS Osaka】Protecting Your Skin!

The Secret of Axolotl Mucus (Mucin)

Authors: F.N. Santako[1, 2]; Sugiyama Axolotl Haruka[1], Yoshichika Yokoe[1, 3, 4, 5]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] Axolotl rescue, [3] The University of Tokyo, [4] Shumei Univ, [5] SciMountechnology,



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Highlight of the Research / Project

Water contamination can be detected using the sensitivity of axolotls!
Compared to fish, axolotls are far more sensitive to chemicals and pollutants, allowing them to detect subtle water contamination that might otherwise go unnoticed.

Abstract

During our investigation of captive Mexican salamanders (Ambystoma mexicanum), commonly known as axolotls, we were struck by their extremely low tolerance to chemicals.
Interestingly, this sensitivity may be advantageous, as it suggests their potential use as biological indicators for detecting residual toxins and chemicals in water.

Keywords

Water pollution, Environmental conservation, Water quality improvement, Environmental cleanup, Axolotl

Fields / Industries for Collaboration

Environmental cleanup, Environmental conservation, Water quality improvement, Wastewater treatment

Background and Motivation

The Mexican salamander (*Ambystoma mexicanum)*, commonly known as the axolotl, is widely kept as a pet. However, wild populations inhabiting Lake Xochimilco in Mexico are currently facing extinction.
Through our research, we found that axolotls respond even to minimal microbial contamination or polluted water, and show strong inflammatory reactions (e.g., redness and congestion) when exposed to chemicals.
This high sensitivity suggests that axolotls could potentially be used as biological indicators to detect residual toxins or chemical contaminants in water.

Current Work and Findings

To quantify their sensitivity to chemicals, we exposed axolotls to very low concentrations of aquarium chemicals, agricultural pesticides, organic solvents, and detergents in rearing water. Even at a concentration of 1000 ppm (1 part per million = 0.0001%), exposure for approximately 30 minutes resulted in visible inflammation, such as reddening of fins and limbs. In some cases, mortality was observed after several weeks.
Ref: https://researchmap.jp/HAL_lab_Axolotl/published_papers/52951197

Future Perspectives

These findings may contribute to applications such as detecting residual pesticides and evaluating the safety of treated water from wastewater treatment facilities.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

【Research】Mathematical Modeling of Surface Breathing Behavior in the Axolotl, *Ambystoma mexicanum*

Authors: Sugiyama Axolotl Haruka[1], Yoshichika Yokoe[1, 2, 3, 4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology,



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Abstract

The Mexican axolotl exhibits facultative air-breathing behavior, surfacing intermittently despite possessing well-developed external gills. This study proposes a mathematical model to describe the behavioral rule governing surface respiration. By integrating environmental oxygen concentration and metabolic demand, we demonstrate that surfacing frequency can be explained as an adaptive response to hypoxia. The model reproduces observed behavioral transitions and provides a framework for predicting axolotl respiration strategies under varying environmental conditions.

1. Introduction

Aquatic animals such as amphibians and fish often exhibit air-breathing behavior by surfacing. This behavior is generally associated with environmental oxygen availability and physiological constraints. In amphibians, multiple respiratory pathways—gills, lungs, and skin—coexist, allowing flexible adaptation to environmental changes [1, 5]. The Mexican Axolotl (*Ambystoma mexicanum*) is a neotenic amphibian that retains external gills but also possesses functional lungs and cutaneous respiration ability [1]. Under normoxic conditions, axolotls rely primarily on gill respiration; however, in hypoxic environments, they increase air-breathing behavior by surfacing to gulp air [2, 7]. Previous studies have shown that axolotls behave as “oxygen conformers,” where oxygen consumption correlates with ambient oxygen tension, and air-breathing increases under hypoxia [2, 7]. Similar behavioral modulation in response to hypoxia has been reported in other amphibians, suggesting a general principle linking respiration and behavior [3, 6].

In this study, we aim to construct a mathematical model that explains:

- Why axolotls surface to breathe
- How often they perform this behavior
- The governing law linking environment and behavior

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

2. Results

2.1 Model formulation

We define the following variables:

- $O_w(t)$: dissolved oxygen concentration in water
- O_a: oxygen concentration in air (constant)
- M: metabolic oxygen demand
- R_g: oxygen uptake rate via gills
- R_l: oxygen uptake rate via lungs (surface breathing)

Total oxygen intake:

$$R_{\text{total}} = R_g(O_w) + R_l \cdot S(t)$$

where:

- $S(t) \in \{0, 1\}$: surface behavior ($1 =$ surfacing event)

Gill respiration is modeled as:

$$R_g(O_w) = k_g \cdot O_w$$

Lung respiration is assumed constant per surfacing:

$$R_l = k_l$$

2.2 Behavioral switching rule

We define a threshold model:

$$S(t) = \begin{cases} 1 & \text{if } R_g(O_w) < M \\ 0 & \text{otherwise} \end{cases}$$

This implies that surfacing occurs when oxygen obtained from gills alone is insufficient.

2.3 Continuous probability model

To better reproduce real behavior, we extend to probabilistic dynamics:

$$P(\text{surfacing}) = \frac{1}{1 + e^{-\alpha(M - R_g(O_w))}}$$

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 1. The probabilistic behavioral switching rule as a function of internal oxygen levels. The surfacing probability $S(t)$ is modeled using a logistic function: $S(t) = 1/(1 + \exp(-\alpha(O_{crit} - O_b)))$. The graph illustrates how the sensitivity parameter α influences the transition sharpness between submerged and surfacing states. A higher α value results in a more discrete, threshold-like switching behavior near the critical oxygen level O_{crit} (indicated by the dashed vertical line).

2.4 Model behavior

- **High oxygen ($O_w \gg M/k_g$):**
  $\rightarrow R_g$ sufficient $\rightarrow S(t) \approx 0$
- **Low oxygen ($O_w \ll M/k_g$):**
  $\rightarrow R_g$ insufficient $\rightarrow S(t) \approx 1$
- **Intermediate oxygen:**
  $\rightarrow$ stochastic surfacing behavior

2.5 Comparison with observed behavior

Empirical observations show:

- Axolotls rarely surface in oxygen-rich water
- Surface breathing increases in hypoxic conditions
- Behavior changes continuously, not abruptly

These features are consistent with:

- Oxygen-dependent surfacing frequency [2, 7]
- Use of lungs as a “backup system” [1, 7]

Thus, the model successfully reproduces observed behavioral patterns.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl

3. Discussion

3.1 Biological interpretation

This model suggests that axolotl surfacing is governed by a supply-demand mismatch principle:

- Oxygen supply (gills) vs. metabolic demand
- Behavioral compensation via lung respiration

This aligns with experimental findings that:

- Air breathing increases when aquatic oxygen decreases [2, 7]
- Lung use supplements gill respiration under stress [1, 7]




Figure 2. Temporal evolution of physiological and environmental oxygen variables. This simulation demonstrates the emergent oscillatory dynamics described in Section S3.1. As metabolic demand M depletes the internal oxygen O_b (blue line), the surfacing probability S increases near the critical threshold O_{crit}. Each surfacing event (red shaded areas) provides a rapid influx of oxygen via the lungs (k_l), resetting the internal state. The environmental water oxygen O_w (green dashed line) exhibits a slow recovery toward saturation O_{sat} mediated by the diffusion rate D.

3.2 Extension of the model

Future improvements may include:

- Energy cost of surfacing
- Predation risk
- Buoyancy control (air gulping affects floating behavior)
- Time-dependent oxygen diffusion dynamics

3.3 Broader implications

This framework may generalize to:

- Other facultative air-breathing amphibians
- Fish species exhibiting hypoxia-driven behavior
- Behavioral physiology modeling in aquatic systems [3, 6, 7]

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4. References

[1] Hillman, S. S., Withers, P. C., Drewes, R. C., & Hillyard, S. D.
Ecological and Environmental Physiology of Amphibians.
Oxford University Press, 2009.
https://doi.org/10.1093/acprof:oso/9780198570325.001.0001

[2] Hochachka, P. W., & Somero, G. N.
Biochemical Adaptation: Mechanism and Process in Physiological Evolution.
Oxford University Press, 2002.
https://doi.org/10.1093/oso/9780195117035.001.0001

[3] Homma, I., & Phillips, A. G.
Respiratory Neurobiology.
In: Handbook of Clinical Neurology, Vol. 188. Elsevier, 2022.
https://doi.org/10.1016/B978-0-444-64076-5.00010-4

[4] Dejours, P.
Principles of Comparative Respiratory Physiology.
Elsevier/North-Holland Biomedical Press, 1981.
ISBN: 9780444802798

[5] Maina, J. N.
Structure, function and evolution of the gas exchangers: comparative perspectives.
Journal of Anatomy, 2002, 201(4), 281–304.

https://doi.org/10.1046/j.1469-7580.2002.00099.x
[6] Heisler, N. (Ed.)
Acid-Base Regulation in Animals.
Elsevier, 1986.
ISBN: 9780444806918

[7] Nilsson, G. E.
Respiratory Physiology of Vertebrates: Life With and Without Oxygen.
Cambridge University Press, 2010.
https://doi.org/10.1017/CBO9780511780813

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Supplementary Information

S1. Extended Mathematical Model

To describe the surface breathing behavior of the Mexican axolotl in a more mechanistic way, we extend the initial threshold model into a continuous dynamical system.

S1.1 Internal Oxygen Dynamics

We introduce the internal oxygen $O_b(t)$ level as a state variable:

$$\frac{dO_b}{dt} = k_g O_w + k_l S(t) - M$$

where:

- $O_b(t)$: internal oxygen level
- O_w: dissolved oxygen concentration in water
- k_g: gill respiration coefficient
- k_l: lung respiration coefficient
- $S(t)$: surfacing behavior
- M: metabolic oxygen consumption

S1.2 Behavioral Response Function

Instead of a binary switching rule, we model surfacing behavior as a continuous probabilistic response:

$$S(t) = \frac{1}{1 + e^{-\alpha(O_{crit} - O_b(t))}}$$

where:

- O_{crit}: critical oxygen threshold
- α: sensitivity parameter controlling behavioral sharpness

This formulation reflects a biologically plausible smooth transition in behavior.

S1.3 Environmental Oxygen Dynamics

To incorporate environmental feedback, we model water oxygen concentration dynamically:

$$\frac{dO_w}{dt} = D(O_{sat} - O_w) - c \cdot k_g O_w$$

where:

- D: oxygen diffusion/recovery rate
- O_{sat}: saturation oxygen level
- c: consumption scaling factor

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

S2. Numerical Simulation

We performed numerical simulations of the coupled system using a simple Euler integration scheme.

S2.1 Python Implementation

```
Python
import numpy as np
import matplotlib.pyplot as plt

# Parameters
dt = 0.01
T = 100
time = np.arange(0, T, dt)

Ow = 10.0     # Initial water oxygen
Ob = 10.0     # Initial body oxygen
M = 1.0       # Metabolic demand
kg = 0.1      # Gill coefficient
kl = 5.0      # Lung coefficient
alpha = 2.0 # Sensitivity
Ocrit = 5.0 # Critical threshold
D = 0.05      # Diffusion rate
Osat = 10.0 # Saturation level

Ob_list, Ow_list, S_list = [], [], []

for t in time:
    # Behavioral response (Logistic)
    S = 1 / (1 + np.exp(-alpha * (Ocrit - Ob)))

    # Dynamics
    dOb = kg * Ow + kl * S - M
    dOw = D * (Osat - Ow) - 0.1 * kg * Ow

    Ob += dOb * dt
    Ow += dOw * dt

    Ob_list.append(Ob)
    Ow_list.append(Ow)
    S_list.append(S)

# Visualization
plt.figure(figsize=(10, 6))
```

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

```
plt.plot(time, Ob_list, label="Body Oxygen (Ob)")
plt.plot(time, Ow_list, label="Water Oxygen (Ow)")
plt.plot(time, np.array(S_list) * 5, label="Surfacing (S) x5", alpha=0.3)
plt.legend()
plt.show()
```


S3. Emergent Dynamics

The model produces several key behaviors:

S3.1 Oscillatory Surfacing Behavior

Under intermediate oxygen conditions, the system exhibits oscillations:

- Decrease in internal oxygen
- Triggering of surfacing event
- Rapid recovery of internal oxygen
- Return to submerged state



Figre S3.1. Emergent Oscillatory Dynamics

This figure demonstrates the temporal evolution of internal oxygen levels (O_b) as a result of the coupled respiratory-behavioral model. The blue line represents the internal oxygen concentration, which gradually declines due to metabolic consumption (M) exceeding aquatic oxygen uptake. When O_b approaches the critical threshold, surfacing behavior is triggered (indicated by the sharp recovery phases). These results confirm the model's ability to reproduce the characteristic periodic air-breathing cycles observed in *Ambystoma mexicanum* under stable environmental conditions.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【Research】Propolis Supplementation Enhances Immune Resilience in the Axolotl, *Ambystoma mexicanum*

Authors: Sugiyama Axolotl Haruka[1], Yoshichika Yokoe[1, 2, 3, 4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology,



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Abstract

Bee-derived substances have long been recognized for their beneficial effects on human health, including antimicrobial, antioxidant, and immunomodulatory activities [1–3,7–9]. However, their application in amphibian biology and conservation remains poorly understood. The Mexican axolotl (Ambystoma mexicanum (Axolotl)), an endangered neotenic amphibian, exhibits relatively weak immune defenses, making it highly susceptible to environmental stress and infection.

In this study, we evaluated the effects of three bee-derived compounds—honey, propolis, and royal jelly (royalactin)—on growth and immune function. Juvenile Axolotls were fed pellet diets coated with each compound from post-hatching day 30 up to day 200. Royal jelly significantly promoted growth and enhanced mucus secretion while suppressing inflammatory responses. Honey moderately enhanced growth but induced inflammatory responses under stress. In contrast, propolis did not affect growth but significantly reduced inflammatory cytokine levels and extended lifespan under high-risk conditions.

Introduction

Bee-derived substances such as honey, propolis, and royal jelly have been widely utilized in human health applications due to their bioactive properties [1–3,7–9]. Honey is rich in carbohydrates and provides rapid metabolic energy, while also exhibiting antimicrobial properties [1–3]. Propolis is a resinous mixture containing flavonoids and phenolic compounds with strong anti-inflammatory and antimicrobial activities [7–9]. Royal jelly contains bioactive proteins such as royalactin and growth factor-like molecules, which influence development and cellular proliferation [4–6].

Royal jelly is a complex secretion produced by worker bees and is characterized by a unique composition of proteins, lipids, vitamins, and bioactive molecules. Its major protein components, collectively known as major royal jelly proteins (MRJPs), account for approximately 80–90% of its total protein content and play essential roles in nutritional and physiological regulation [6]. Among these, royalactin has been identified as a key factor influencing developmental pathways, including growth and differentiation, partly through activation of epidermal growth factor receptor (EGFR)-related signaling [4,5]. In addition, royal jelly contains bioactive lipids such as 10-hydroxy-2-decenoic acid (10-HDA), which has been reported to exhibit antimicrobial and immunomodulatory activities. The combined action of these components suggests that royal jelly functions not only as a nutrient source but also as a potent regulator of growth and immune-related processes, making it a strong candidate for comparative functional analysis with other bee-derived substances.

The Mexican axolotl (Ambystoma mexicanum (Axolotl)) is a neotenic amphibian that retains larval features such as external gills throughout its life. Despite possessing multiple respiratory pathways (gills, lungs, and skin), Axolotls are known to have relatively limited immune defenses, making them vulnerable to infection and environmental deterioration. This represents a major challenge for conservation efforts.

Previous studies in vertebrates suggest that immune modulation can be achieved through dietary supplementation of bioactive compounds [1,7,9]. However, comparative evaluation of different bee-derived substances in amphibians has not been systematically investigated.

In this study, we aimed to determine:

- **Which bee-derived component most effectively enhances immune function**
- **Whether growth and immune effects can be dissociated**
- **How these effects influence survival under environmental stress**

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

To address this, honey, propolis, and royal jelly were administered under identical conditions, and their effects on growth, immune response, and survival were quantitatively compared.

Materials and Methods

Animals and Rearing Conditions

Juvenile Axolotls were used. Individuals were maintained under controlled aquatic conditions (18–20°C, 12 h light/dark cycle) in dechlorinated water.

Experimental Design

Animals were randomly divided into four groups:

- Control (no supplementation)
- Honey-treated
- Propolis-treated
- Royal jelly-treated

Each group received pelletized feed coated with the respective substance dissolved in rearing water. Feeding began at post-hatching day 200.

Sampling points: Day 60 (30 days treatment)

Measured Parameters

- Growth rate (body length, body mass)
- Survival rate
- Inflammatory cytokine levels (e.g., IL-like markers)
- Mucus secretion (histological staining)

In this assay, all individuals used were unmated males and females.




Figure 1. Day 60 Comprehensive Analysis of Gene Expression Dynamics in male Gonads

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Table.1 Royal Jelly Composition

Category	Component	Content (Typical Range)	Function / Notes
Water	Water	60–70%	Major component; affects stability
Proteins	Total protein	9–18%	Core bioactive fraction
	MRJP1 (Royalactin)	40–50% of protein	Caste differentiation, EGFR activation
	MRJP2–9	30–50% of protein	Immunity, metabolism
	Enzymes	Few %	Antimicrobial, metabolic roles
	Antimicrobial peptides	Trace	Defense function
Lipids	Total lipids	3–8%	Rich in unusual fatty acids
	10-HDA	1–3% (w/w)	Unique marker, antimicrobial, signaling
	Other fatty acids	1–4%	Medium-chain, hydroxy fatty acids
Carbohydrates	Fructose	3–8%	Energy source
	Glucose	3–8%	Energy source
	Oligosaccharides	Trace	Possible gut-related effects
Vitamins	Vitamin B5 (Pantothenic acid)	5–20 mg/100 g	Highly abundant, metabolism
	Vitamin B2 (Riboflavin)	1–3 mg/100 g	Energy metabolism
	Vitamin B1 (Thiamine)	0.5–1.5 mg/100 g	Neural function
	Vitamin B6	0.5–1 mg/100 g	Amino acid metabolism
	Niacin	3–10 mg/100 g	Redox reactions
	Folate	0.1–0.5 mg/100 g	Cell proliferation
Minerals	Potassium	200–400 mg/100 g	Electrolyte balance
	Calcium	20–60 mg/100 g	Structural, signaling
	Magnesium	10–40 mg/100 g	Enzyme cofactor
	Iron	1–5 mg/100 g	Oxygen transport
	Zinc	1–3 mg/100 g	Enzyme, immunity
Bioactive compounds	Acetylcholine	1–5 mg/100 g	Neurotransmission
	Polyphenols	Trace	Antioxidant
	Organic acids (pH 3.5–4.5)	Trace	Antimicrobial environment
Other indicators	Ash	0.8–3%	Total mineral content
	pH	3.5–4.5	Acidic nature
	Dry matter	~30–40%	Nutrient density

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Results

Growth Effects

Royal jelly treatment resulted in the highest growth rate across all time points [4–6]. Honey-treated animals showed moderate growth enhancement, likely due to increased carbohydrate availability [1–3]. Propolis treatment did not significantly affect growth compared to controls. In addition, RNA samples were collected from both epithelial tissues and gonadal tissues to evaluate tissue-specific responses to each treatment.




Figure 2. Day 60 Comprehensive Analysis of Gene Expression Dynamics in male Gonads

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Immune Responses under Stress Conditions

Honey group

- ・Elevated inflammatory cytokine levels
- ・Increased mucus secretion
- ・No significant reduction in lifespan

Royal jelly group

- ・Suppressed inflammatory responses
- ・Strong increase in mucus secretion
- ・Significant lifespan extension

Propolis group

- ・Suppressed inflammatory cytokines
- ・Minimal effect on mucus secretion
- ・Significant lifespan extension

Furthermore, sex-dependent differences were observed in treatment responses. Female individuals exhibited stronger effects of EGF-related signaling and royal jelly supplementation compared to males, suggesting a higher sensitivity to growth factor-mediated pathways.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 3. Day 60 Comprehensive Analysis of Gene Expression Dynamics in male Gonads




Figure 4. Day 60 Comprehensive Analysis of Gene Expression Dynamics in male epithelium.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 5. Day 60 Comprehensive Analysis of Gene Expression Dynamics in female Gonads




Figure 6. Day 60 Comprehensive Analysis of Gene Expression Dynamics in female epithelium

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




Figure 7. Effect on miRNA expression levels involved in inflammation and regeneration-1

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl




Figure 8. Effect on miRNA expression levels involved in inflammation and regeneration-2

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 9. Effect on miRNA expression levels involved in inflammation and regeneration-3

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 10. Effect on miRNA expression levels involved in inflammation and regeneration-4

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Survival Analysis

Under high-risk conditions, both royal jelly and propolis groups showed improved survival compared to control. The propolis group demonstrated improved survival without increased metabolic cost (no growth change).




Figure 11. urvival rate of the *Axolotl*

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Discussion

Functional Differences among Bee-derived Substances

The results demonstrate clear functional separation:

- ・Honey promotes growth primarily through metabolic energy supply [1–3]
- ・Royal jelly exerts pleiotropic effects via royalactin and growth factor signaling [4–6]
- ・Propolis acts mainly as an immunomodulatory agent [7–9]

Mechanistic Interpretation

The anti-inflammatory effects observed in the propolis group are consistent with known actions of flavonoids and phenolic compounds [7–9]. Unlike royal jelly, propolis did not enhance mucus secretion, suggesting that it acts on systemic immune regulation rather than barrier enhancement.

Royal jelly appears to enhance both physical (mucus) and molecular (cytokine suppression) defenses, indicating a broader regulatory role.

Implications for Conservation

These findings suggest:

- ・Propolis is suitable for immune support without altering growth dynamics
- ・Royal jelly is effective for combined growth and immune enhancement
- ・Honey is useful for energy supplementation but not immune enhancement

Thus, propolis may be particularly useful in conservation settings where infection risk is high.

Conclusion

Propolis supplementation enhances immune resilience and survival in Axolotls without affecting growth, indicating its role as an immunomodulatory agent. This property may be valuable for improving the success of captive breeding and conservation programs.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure S1. Endogenous viral sequence in the *Axolotl*

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl

References

[1] Bogdanov, S., Jurendic, T., Sieber, R., & Gallmann, P.
Honey for nutrition and health: a review.
Journal of the American College of Nutrition, 2008, 27(6), 677–689.
https://doi.org/10.1080/07315724.2008.10719745

[2] Al-Waili, N., Salom, K., Butler, G., & Al Ghamdi, A. A.
Honey and microbial infections: a review supporting the use of honey for microbial control.
The Scientific World Journal, 2011, 11, 766–787.
https://doi.org/10.1100/tsw.2011.80

[3] Mandal, M. D., & Mandal, S.
Honey: its medicinal property and antibacterial activity.
Asian Pacific Journal of Tropical Biomedicine, 2011, 1(2), 154–160.
https://doi.org/10.1016/S2221-1691(11)60016-6

[4] Kamakura, M.
Royalactin induces queen differentiation in honeybees.
Nature, 2011, 473, 478–483.
https://doi.org/10.1038/nature10093

[5] Buttstedt, A., Moritz, R. F. A., & Erler, S.
Royalactin is not a royal making of a queen.
Nature, 2016, 537, E10–E12.
https://doi.org/10.1038/nature18004

[6] Fujita, T., Kozuka-Hata, H., Ao-Kondo, H., et al.
Proteomic analysis of the royal jelly and characterization of the functions of its derivation glands in the honeybee
Journal of Proteome Research, 2013 Jan 4;12(1):404-11. doi: 10.1021/pr300700e. Epub 2012 Nov 29.
https://doi.org/10.1021/pr400700e

[7] Sforcin, J. M.
Propolis and the immune system: a review.
Journal of Ethnopharmacology, 2007, 113(1), 1–14.
https://doi.org/10.1016/j.jep.2006.05.012

[8] Banskota, A. H., Tezuka, Y., & Kadota, S.
Recent progress in pharmacological research of propolis.
Phytotherapy Research, 2001, 15(7), 561–571.
https://doi.org/10.1002/ptr.1029

[9] Silva-Carvalho, R., Baltazar, F., & Almeida-Aguiar, C.
Propolis: a complex natural product with a plethora of biological activities.
Evidence-Based Complementary and Alternative
Evid Based Complement Alternat Med. 2015 May 27;2015:206439
https://doi.org/10.1155/2015/206439

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【Review】Nucleotide-Based Immune Strategies

Authors: Sugiyama Axolotl Haruka[1], Yoshichika Yokoe[1, 2, 3, 4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology,



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Abstract

RNA-based immune mechanisms represent an evolutionarily conserved and highly versatile defense strategy against viral and bacterial pathogens. These systems range from sequence-specific RNA interference (RNAi) to interferon-mediated antiviral pathways and small RNA-guided silencing. While mammals rely predominantly on interferon signaling and RNA sensing pathways, amphibians exhibit a hybrid system incorporating both RNAi-like and vertebrate innate immune responses. Arthropods, particularly insects, utilize RNAi as a primary antiviral mechanism. This review comprehensively summarizes RNA-mediated immune strategies across mammals, amphibians, and arthropods, highlighting molecular components, evolutionary transitions, and functional diversity.

1. Introduction

RNA-based immunity predates adaptive immunity and represents a core defense strategy conserved across eukaryotes [2,23]. These systems rely on the recognition of foreign nucleic acids, particularly viral RNA, and the subsequent activation of degradation or silencing pathways.

RNA functions in immunity can be categorized into:

- **Sensors**: Detection of pathogen-associated RNA
- **Effectors**: Silencing or degradation of target RNA
- **Regulators**: Fine-tuning immune responses (e.g., miRNA)

Major RNA-mediated immune pathways include:

- RNA interference (RNAi)
- Interferon-mediated antiviral responses
- Pattern recognition receptor (PRR)-mediated RNA sensing
- Small RNA-mediated gene regulation

2. RNA-Based Immunity in Mammals

2.1 Overview of Mammalian RNA Immunity

Unlike invertebrates, mammals primarily rely on **protein-mediated antiviral systems**, especially the interferon (IFN) pathway [6,10]. However, RNA remains central as:

- a **trigger molecule** (viral RNA detection)
- a **regulatory molecule** (miRNA, siRNA)

2.2 RNA Recognition Systems

RIG-I-like Receptors (RLRs)

Key cytoplasmic sensors include:

- RIG-I (recognizes 5′-triphosphate RNA)
- MDA5 (recognizes long dsRNA)

These receptors initiate signaling via MAVS, leading to IFN production [4,5].

2.3 Toll-like Receptors (TLRs)

Endosomal RNA sensors:

- TLR3 → dsRNA
- TLR7/8 → ssRNA

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

These receptors activate NF-κB and IRF pathways.

2.4 Interferon Response

Upon RNA detection:

1. IRF3/7 activation
2. Type I IFN secretion
3. Induction of interferon-stimulated genes (ISGs)

Major ISGs include:

- PKR → inhibits translation
- OAS/RNase L → RNA degradation
- IFIT → blocks viral RNA translation

[6,10]

2.5 RNAi in Mammals

Although present, RNAi plays a **limited antiviral role**:

- Active in embryonic stem cells [8]
- Suppressed in differentiated cells due to IFN dominance

Experimental evidence shows RNAi emerges when IFN signaling is disrupted [7].

2.6 microRNA in Immunity

miRNAs regulate:

- cytokine production
- immune signaling pathways
- viral replication

Viruses can encode miRNAs to manipulate host immunity [9].

2.7 Summary Table (Mammals)

Category	Factor	Function
RNA sensor	RIG-I	Detects 5′ppp RNA
RNA sensor	MDA5	Detects dsRNA
adaptor	MAVS	Signal transduction
cytokine	IFN-α/β	Antiviral signaling
effector	PKR	Translation inhibition
effector	OAS/RNase L	RNA degradation
RNAi	Dicer	dsRNA processing
RNAi	AGO2	RNA cleavage
regulatory	miRNA	Immune modulation

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

3. RNA-Based Immunity in Amphibians

3.1 Evolutionary Significance

Amphibians represent a **transitional immune state**:

- possess IFN system (vertebrate trait)
- retain potential RNAi contributions (ancestral trait)

[11]

3.2 Interferon System in Amphibians

Amphibians exhibit:

- multiple IFN subtypes
- strong antiviral responses to ranaviruses

[12,13]

3.3 RNAi Mechanisms

Evidence suggests:

- dsRNA processing into siRNAs
- antiviral RNA silencing activity

Though less dominant than in insects, RNAi likely contributes to antiviral defense.

3.4 microRNA Regulation

miRNAs regulate:

- immune gene expression
- inflammation
- antiviral signaling

[14]

3.5 Amphibian Antiviral Responses

Studies in *Xenopus* show:

- IFN-dependent viral clearance
- RNA-based regulatory fine-tuning

[12]

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

3.6 Summary Table (Amphibians)

Category	Factor	Function
RNA sensor	RIG-I homologs	Viral detection
cytokine	Type I IFN	Antiviral signaling
RNAi	Dicer	siRNA production
RNAi	AGO	RNA silencing
regulatory RNA	miRNA	Immune regulation
effector	ISGs	Viral inhibition

4. RNA-Based Immunity in Arthropods

4.1 Overview

Arthropods rely almost exclusively on:

RNA interference as primary antiviral immunity

[18]

4.2 siRNA Pathway

Core antiviral mechanism:

1. Viral dsRNA recognition
2. Cleavage by Dicer-2
3. Loading into AGO2
4. Target RNA degradation

[17]

4.3 piRNA Pathway

Functions:

- transposon silencing
- antiviral defense in germline

Amplification via “ping-pong” cycle [21]

4.4 Systemic RNAi

RNA signals spread across tissues:

- enables organism-wide immunity
- mediated by dsRNA uptake systems

[20]

4.5 Viral Suppression of RNAi

Viruses encode:

- RNAi suppressor proteins
- dsRNA masking mechanisms

[2]

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

4.6 Summary Table (Arthropods)

Category	Factor	Function
enzyme	Dicer-2	dsRNA processing
effector	AGO2	RNA cleavage
cofactor	R2D2	siRNA loading
piRNA protein	Piwi	Genome defense
systemic	SID-1-like	RNA transport
amplification	ping-pong cycle	Signal amplification

5. Comparative Evolutionary Analysis

5.1 Functional Transition

Feature	Arthropods	Amphibians	Mammals
RNAi antiviral	Strong	Moderate	Weak
IFN system	Absent	Present	Strong
RNA sensing	Dicer-based	Mixed	PRR-based

5.2 Evolutionary Interpretation

- RNAi = ancestral antiviral system
- IFN = later innovation in vertebrates
- mammals prioritized **broad, rapid response over specificity**

[23,24]

6. Viral Countermeasures

Viruses evolved strategies such as:

- RNAi suppressors
- IFN pathway inhibition
- RNA structure masking

Examples include:

- dsRNA sequestration
- viral miRNAs
- decoy RNAs

7. Integration of RNA and Protein Immunity

Modern immunity is layered:

- RNA sensing triggers IFN
- miRNA fine-tunes immune signaling
- RNAi acts as backup system

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

8. Applications

8.1 Medicine

- siRNA therapeutics
- mRNA vaccines
- antiviral RNA drugs

8.2 Agriculture

- RNAi pest control
- virus-resistant insects

8.3 Evolutionary Biology

- RNA world hypothesis linkage
- transition to protein-based immunity

9. Future Perspectives

Key questions:

- Why was RNAi suppressed in mammals?
- Can RNAi be therapeutically reactivated?
- How do amphibians maintain hybrid systems?

10. Conclusion

RNA-based immunity is a foundational biological defense mechanism that has diversified across evolution. Arthropods retain RNAi as their primary antiviral strategy, amphibians exhibit a hybrid immune system, and mammals rely predominantly on interferon-mediated responses while maintaining residual RNA-based regulatory mechanisms.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

References

[1] Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E., & Mello, C. C.
Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*.
Nature, 1998, 391, 806–811.
https://doi.org/10.1038/35888

[2] Ding, S. W., & Voinnet, O.
Antiviral immunity directed by small RNAs.
Cell, 2007, 130(3), 413–426.
https://doi.org/10.1016/j.cell.2007.07.039

[3] Cullen, B. R.
RNA interference: antiviral defense and genetic tool.
Nature Immunology, 2006, 7(6), 563–567.
https://doi.org/10.1038/ni1332

[4] Kato, H., Takeuchi, O., Sato, S., et al.
Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses.
Nature, 2006, 441, 101–105.
https://doi.org/10.1038/nature04734

[5] Loo, Y. M., & Gale, M.
Immune signaling by RIG-I-like receptors.
Immunity, 2011, 34(5), 680–692.
https://doi.org/10.1016/j.immuni.2011.05.003

[6] Schneider, W. M., Chevillotte, M. D., & Rice, C. M.
Interferon-stimulated genes: a complex web of host defenses.
Annual Review of Immunology, 2014, 32, 513–545.
https://doi.org/10.1146/annurev-immunol-032713-120231

[7] Maillard, P. V., Van der Veen, A. G., Deddouche-Grass, S., et al.
Inactivation of the type I interferon pathway reveals long double-stranded RNA-mediated RNA interference in mammalian cells.
EMBO Journal, 2016, 35(23), 2505–2518.
https://doi.org/10.15252/embj.201694875

[8] Li, Y., Lu, J., Han, Y., Fan, X., & Ding, S. W.
RNA interference functions as an antiviral immunity mechanism in mammals.
Science, 2013, 342(6155), 231–234.
https://doi.org/10.1126/science.1241911

[9] Umbach, J. L., & Cullen, B. R.
The role of RNAi and microRNAs in animal virus replication and antiviral immunity.
Genes & Development, 2009, 23(10), 1151–1164.
https://doi.org/10.1101/gad.1793309

[10] Sadler, A. J., & Williams, B. R.
Interferon-inducible antiviral effectors.
Nature Reviews Immunology, 2008, 8, 559–568.
https://doi.org/10.1038/nri2314

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

[11] Robert, J., & Ohta, Y.
Comparative and developmental study of the immune system in amphibians.
Developmental & Comparative Immunology, 2009, 33(3), 273–283.
https://doi.org/10.1016/j.dci.2008.08.006
[12] Grayfer, L., De Jesús Andino, F., & Robert, J.
The amphibian (Xenopus laevis) type I interferon response to frog virus 3.
Developmental & Comparative Immunology, 2014, 42(1), 104–113.
https://doi.org/10.1016/j.dci.2013.06.015
[13] De Jesús Andino, F., Chen, G., Li, Z., et al.
Innate immune responses to ranavirus infection in amphibians.
Viruses, 2012, 4(11), 3105–3126.
https://doi.org/10.3390/v4113105
[14] Huang, B., Zhang, L., Du, Y., et al.
MicroRNA regulation in amphibian immune responses.
Fish & Shellfish Immunology, 2018, 77, 146–154.
https://doi.org/10.1016/j.fsi.2018.03.036
[15] Small, H. J., & Peddie, S.
Molecular biology of amphibian antiviral immunity.
Viruses, 2014, 6(7), 2624–2645.
https://doi.org/10.3390/v6072624
[16] Hannon, G. J.
RNA interference.
Nature, 2002, 418, 244–251.
https://doi.org/10.1038/418244a
[17] Wang, X. H., Aliyari, R., Li, W. X., et al.
RNA interference directs innate immunity against viruses in adult Drosophila.
Science, 2006, 312(5772), 452–454.
https://doi.org/10.1126/science.1125694
[18] Bronkhorst, A. W., & van Rij, R. P.
The long and short of antiviral defense: small RNA-based immunity in insects.
Current Opinion in Virology, 2014, 7, 19–28.
https://doi.org/10.1016/j.coviro.2014.03.010
[19] Blair, C. D., & Olson, K. E.
The role of RNA interference (RNAi) in arbovirus-vector interactions.
Viruses, 2015, 7(2), 820–843.
https://doi.org/10.3390/v7020820
[20] Saleh, M. C., Tassetto, M., van Rij, R. P., et al.
Antiviral immunity in Drosophila requires systemic RNA interference spread.
Nature, 2009, 458, 346–350.
https://doi.org/10.1038/nature07712
[21] Miesen, P., Joosten, J., & van Rij, R. P.
PIWI-interacting RNAs in insects: biogenesis and function.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Annual Review of Entomology, 2016, 61, 1–17.
https://doi.org/10.1146/annurev-ento-010715-023949
[22] Schnettler, E., Tykalová, H., Watson, M., et al.
Induction and suppression of tick cell antiviral RNAi responses by tick-borne flaviviruses.
Nucleic Acids Research, 2014, 42(14), 9436–9446.
https://doi.org/10.1093/nar/gku657
[23] Obbard, D. J., Gordon, K. H., Buck, A. H., & Jiggins, F. M.
The evolution of RNAi as a defence against viruses and transposable elements.
Philosophical Transactions of the Royal Society B, 2009, 364, 99–115.
https://doi.org/10.1098/rstb.2008.0168
[24] tenOever, B. R.
The evolution of antiviral defense systems.
Cell Host & Microbe, 2016, 19(2), 142–149.
https://doi.org/10.1016/j.chom.2016.01.006
[25] Barrangou, R., Fremaux, C., Deveau, H., et al.
CRISPR provides acquired resistance against viruses in prokaryotes.
Science, 2007, 315(5819), 1709–1712.
https://doi.org/10.1126/science.1138140
[26] Gantier, M. P., & Williams, B. R.
The response of mammalian cells to double-stranded RNA.
Cytokine & Growth Factor Reviews, 2007, 18(5–6), 363–371.
https://doi.org/10.1016/j.cytogfr.2007.06.016
[27] Hornung, V., & Latz, E.
Intracellular DNA and RNA sensing pathways in innate immunity.
Immunity, 2010, 32(4), 453–465.
https://doi.org/10.1016/j.immuni.2010.03.011

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

【Research】Metabolic State Transition and Longevity Strategy in Axolotls: A Proteus-like Mode Hypothesis

Authors: F.N. Santako[1, 5]; Sugiyama Axolotl Haruka[1], Yoshichika Yokoe[1, 2, 3, 4]

[1] NPO Research Group_HAL-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ, [4] SciMountechnology, HAL lab Axolotl Rescue[5]



SNS and Mail Address

Researchmap: https://researchmap.jp/HAL_lab_Axolotl

X: https://x.com/lab_new2

ResearchGate: https://www.researchgate.net/profile/Sugiyama-Haruka

Contact: sugiyama.haruka.axoltl@gmail.com

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

Abstract

The Axolotl is a well-established model for regeneration and developmental biology. Under stress conditions, Axolotl often exhibit a marked reduction in body mass and a transition to a slender, elongated morphology. This phenotype has traditionally been interpreted as pathological, associated with infection, malnutrition, or incomplete metamorphosis. However, the persistence of this morphology even after recovery suggests an alternative interpretation.Here, we propose that the slender phenotype represents a strategic physiological transition into a low-metabolic, longevity-associated state, analogous to the phenotype observed in the olm. By integrating comparative physiology, transcriptomics, and metabolic analyses, this study aims to elucidate whether Axolotl can undergo a reversible or irreversible shift toward a survival-oriented mode characterized by reduced energy expenditure and enhanced lifespan potential.

Background

Phenotypic plasticity in amphibians allows for adaptive responses to environmental stressors, including nutrient limitation, hypoxia, and temperature fluctuations. In Axolotl, stress-induced morphological changes often manifest as reduced body mass and elongation. These changes are typically considered pathological. However, the irreversibility of this state in many individuals raises the possibility that it reflects a stable physiological reprogramming rather than damage.The olm, a cave-dwelling amphibian, exhibits extreme longevity (often exceeding 70–100 years), low metabolic rates, and remarkable resistance to environmental stress. These characteristics are associated with reduced oxidative damage, altered insulin/IGF signaling, and efficient mitochondrial function. Similar strategies are observed in other long-lived organisms, suggesting conserved mechanisms of metabolic suppression and lifespan extension. We hypothesize that Axolotl may possess an unrecognized capacity to transition into a comparable physiological state—a “Proteus-like mode”—characterized by metabolic downregulation and enhanced survival under adverse conditions.

Figure 1. Long-lived amphibians related to the “olm” (Proteus-like lineage and comparable taxa)

(A) Proteus anguinus (Olm)



The olm is a cave-dwelling salamander native to the Dinaric karst. It is one of the longest-lived amphibians known, with documented lifespans exceeding 70 years and estimates suggesting over 100 years under natural conditions. This species exhibits extremely low metabolic rates, reduced activity, and resistance to starvation, making it a model for energy-conserving longevity strategies.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

(B)

Andrias japonicus (Japanese giant salamander)



Legend: The Japanese giant salamander is one of the largest amphibians in the world. Individuals have been reported to live **50–80 years**, with some anecdotal records suggesting even longer lifespans in protected environments. Its slow metabolism, low activity, and cold-water habitat are considered contributing factors to its longevity.

(C)

Cryptobranchus alleganiensis (Hellbender)



Legend: The hellbender is a large aquatic salamander from North America. Wild individuals commonly reach **30 years**, while captive individuals have been reported to live **over 50 years**. Their longevity is associated with low metabolic demand and stable aquatic environments.

(D)

Necturus maculosus (Mudpuppy)



Legend: The mudpuppy is a fully aquatic salamander retaining external gills throughout life, similar to Axolotl. It can live **20–30 years** in the wild and longer in captivity. Its neotenic lifestyle and relatively low metabolic rate make it a useful comparative model for studying longevity in aquatic amphibians.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Representative species exhibiting extended lifespan and low metabolic strategies are shown. The olm is particularly notable, with reported lifespans exceeding 70–100 years, making it one of the longest-lived amphibians. Other large or fully aquatic salamanders, including Japanese giant salamander, hellbender, and mudpuppy, also exhibit extended longevity (20–80+ years). These species share key physiological traits such as reduced metabolic rate, aquatic or hypoxic environments, and energy-conserving life-history strategies, providing a comparative framework for understanding longevity-associated phenotypes in amphibians.




Figure 2. Morphological transition of Axolotl toward a Proteus-like phenotype.

(A) A healthy Axolotl in the “normal mode,” characterized by a robust body and well-developed tissues. (B–D) Individuals exhibiting a slender, elongated “Proteus-like” morphology.

We hypothesized that, if the elongated body form of the olm is associated with its exceptional longevity, a similar phenotype observed in Axolotl may not be merely pathological. Instead, this morphological shift could represent a strategic transition into an energy-conserving, longevity-associated metabolic mode. To test this possibility, we conducted a detailed physiological and molecular investigation.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Results and Discussions

Until now, it had been assumed that these individuals died simply due to pseudo-metamorphosis; however, a general decline in overall metabolic activity was observed. In individuals that had experienced reproduction, a marked reduction in immune function was evident, suggesting a condition analogous to “poor postpartum recovery.” Among such individuals, as well as those that had experienced prolonged infection, some were able to survive while maintaining a markedly emaciated body condition. Although this phenotype is often associated with infection, there are also cases in which individuals recover to a healthy state and survive long-term while retaining this body form. Based on these observations, it is possible that individuals subjected to severe physiological stress intentionally shift into a metabolic “off-mode,” enabling long-term survival through reduced energy expenditure and gradual recovery. Under conditions of significantly low water temperature, individuals in this off-mode tend to exhibit extended lifespans, further supporting the idea that this state represents a survival strategy for individuals under near-lethal stress. However, even if environmental conditions improve dramatically, this off-mode cannot be rapidly reversed. Once individuals enter this energy-saving state, they appear constrained to continue living within its limitations, revealing a critical drawback of this strategy.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 3. Traits also observed in Axolotls with an “olm-like” body morphology.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




Figure 4. Dynamics of inflammation- and regeneration-related miRNA expression depending on mating status.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！
HAL lab.axolotl




Figure 5. In low-temperature environments, the “energy-saving mode” can even be advantageous for survival.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

References

1. Voss SR et al., Salamander genomics and regeneration, *Annual Review of Genomics and Human Genetics*, 2015, 16: 161–185, https://doi.org/10.1146/annurev-genom-090413-025429
2. Nowoshilow S et al., The Axolotl genome and regeneration, *Nature*, 2018, 554: 50–55, https://doi.org/10.1038/nature25458
3. McCusker C & Gardiner DM, The Axolotl model for regeneration, *Methods in Molecular Biology*, 2011, 770: 139–159, https://doi.org/10.1007/978-1-61779-210-6_7
4. Tanaka EM, The molecular and cellular basis of limb regeneration, *Cell*, 2016, 165: 1598–1608, https://doi.org/10.1016/j.cell.2016.05.026
5. Galliot B & Ghila L, Regeneration in animals, *Cold Spring Harbor Perspectives in Biology*, 2010, 2: a000505, https://doi.org/10.1101/cshperspect.a000505
6. Voituron Y et al., Extreme lifespan of Proteus anguinus, *Biology Letters*, 2011, 7: 105–107, https://doi.org/10.1098/rsbl.2010.0539
7. Hervant F et al., Metabolic adaptation in cave amphibians, *Journal of Experimental Biology*, 2001, 204: 1143–1150, https://doi.org/10.1242/jeb.204.6.1143
8. Bulog B et al., Biology of Proteus anguinus, *Acta Biologica Slovenica*, 2000, 43: 55–69, URL: https://www.dlib.si
9. Aljančič G et al., Olm ecology and longevity, *Natura Sloveniae*, 2007, 9: 21–36, URL: https://www.natura-sloveniae.si
10. Speakman JR, The physiological costs of reproduction, *Philosophical Transactions of the Royal Society B*, 2008, 363: 375–398, https://doi.org/10.1098/rstb.2007.2145
11. López-Otín C et al., The hallmarks of aging, *Cell*, 2013, 153: 1194–1217, https://doi.org/10.1016/j.cell.2013.05.039
12. Kenyon CJ, The genetics of ageing, *Nature*, 2010, 464: 504–512, https://doi.org/10.1038/nature08980
13. Hulbert AJ et al., Metabolism and longevity, *Physiological Reviews*, 2007, 87: 1175–1213, https://doi.org/10.1152/physrev.00047.2006
14. Barja G, Free radicals and aging, *Trends in Neurosciences*, 2004, 27: 595–600, https://doi.org/10.1016/j.tins.2004.07.005
15. Feder ME & Burggren WW, Environmental physiology of amphibians, *University of Chicago Press*, 1992, URL: https://press.uchicago.edu
16. Wells KD, The Ecology and Behavior of Amphibians, *University of Chicago Press*, 2007, URL: https://press.uchicago.edu
17. Tattersall GJ & Ultsch GR, Metabolism in aquatic amphibians, *Journal of Thermal Biology*, 2008, 33: 200–210, https://doi.org/10.1016/j.jtherbio.2008.02.001
18. Ramsey JJ et al., Energy metabolism and lifespan, *Experimental Gerontology*, 2000, 35: 283–290, https://doi.org/10.1016/S0531-5565(00)00098-3
19. Storey KB, Metabolic regulation in stress tolerance, *Comparative Biochemistry and Physiology*, 2015, 181: 1–10, https://doi.org/10.1016/j.cbpa.2014.10.001
20. Austad SN, Comparative biology of aging, *Journal of Gerontology*, 2009, 64A: 199–201, https://doi.org/10.1093/gerona/gln060
21. Buffenstein R, Negligible senescence in long-lived animals, *Age*, 2008, 30: 135–146, https://doi.org/10.1007/s11357-008-9051-5

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

【研究報告】魚類・両生類の試料用途としてのダンゴムシの可能性

Research: Potential Application of Pillbugs as Feed or Experimental Organisms in Ichthyological and Amphibian Research

著:横江 諠衡 [1, 2, 3, 4]; 杉山 遥 [1], 竜岡 竜騎 [5], 阿保郎 太朗 [6], F.N. 三太子 [1]


[1.] NPO 研究グループ_ハルラボ (ウーパールーパー研究室), [2.] 東京大学, [3.] 秀明大学, [4.] サイマン・テクノロジー,

[5.] トラフサンショウウオラボ, [6.]ハルラボ・ファーム

Authors: Sugiyama Axolotl Haruka[1]; Yoshichika Yokoe[1,2,3,4]; F.N. Santako[1]

[1] NPO Research Group_HARU-Lab (Axolotl Research Laboratory), [2] The University of Tokyo, [3] Shumei Univ,

[4] SciMountechnology, [5] Ambys Lab, [6] Hal lab farm



※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

要旨

近年、魚類および両生類の飼育・実験において、安定供給が可能で栄養組成の制御が容易な試料生物の探索が進められている。本研究では、陸生等脚類であるダンゴムシ（Armadillidium spp.）に着目し、魚類・両生類の試料用途および餌料生物としての可能性を検討した。

ダンゴムシは甲殻類に属し、外骨格に由来するカルシウムを豊富に含有する点が特徴である。カルシウムは脊椎動物において骨格形成、成長促進、神経・筋機能の維持に不可欠な元素であり、とくに成長期の魚類・両生類では摂取量が成長率に大きく影響することが知られている。実際に、メキシコサラマンダー（Ambystoma mexicanum、以下アホロートル）においては、餌中カルシウム量の増加により成長率が顕著に向上することが報告されている。また、この成長促進効果は、無機カルシウムの添加に限らず、エビやカニ類などの甲殻類系餌料の摂取によっても同様に得られることが示唆されている。

これらの知見を踏まえると、ダンゴムシはカルシウム供給源として機能する可能性が高く、魚類・両生類、とりわけアホロートルのような成長段階でカルシウム要求量の高い種に対して、有効な補助餌料あるいは試料生物となり得る。さらに、ダンゴムシは比較的容易に大量飼育が可能であり、環境条件や餌内容を制御することで栄養組成の標準化も期待できる。

以上より、ダンゴムシは魚類・両生類の飼育・実験において、成長促進効果の評価、カルシウム代謝研究、あるいは代替甲殻類餌料としての利用が見込まれる。本要旨は、今後の基礎的飼育試験および栄養学的解析に向けた位置づけを示すものである。

Abstract

In recent years, there has been growing interest in identifying alternative model and feeder organisms for fish and amphibian husbandry and experimental research, particularly those that are easy to culture and nutritionally controllable. In this study, we focus on terrestrial isopods (pill bugs; Armadillidium spp.) and evaluate their potential as experimental materials and dietary resources for fish and amphibians.

Pill bugs are crustaceans characterized by a calcium-rich exoskeleton. Calcium plays a crucial role in vertebrates, contributing to skeletal formation, growth promotion, and the maintenance of neuromuscular function. In fish and amphibians, especially during juvenile stages, dietary calcium intake is known to strongly influence growth performance. Notably, in the Mexican salamander (Ambystoma mexicanum, axolotl), increased calcium intake has been reported to result in a marked enhancement of growth rate. Importantly, similar growth-promoting effects are observed not only with inorganic calcium supplementation but also through the consumption of crustacean-based diets, such as shrimp or crab-derived feed.

Based on these findings, pill bugs may serve as an effective calcium source and represent a promising supplemental feeder organism for fish and amphibians, particularly for species with high calcium demands during growth, such as the axolotl. In addition, pill bugs can be mass-cultured with relative ease, and their nutritional composition may be standardized by controlling environmental and dietary conditions.

In conclusion, pill bugs have considerable potential as experimental and dietary organisms in fish and amphibian research, including applications in growth promotion studies, calcium metabolism research, and as an alternative crustacean-based feed source. This abstract provides a conceptual framework for future controlled feeding experiments and nutritional analyses.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

第 1 章　総論：ダンゴムシとワラジムシとは何か

一般に「ダンゴムシ」「ワラジムシ」と呼ばれる生物は、**節足動物門・甲殻亜門・等脚目（Isopoda）・陸生等脚類（Oniscidea）**に属する動物である。
昆虫やムカデ類（多足類）と混同されやすいが、分類学的にはエビやカニに近い甲殻類である。
陸生等脚類は、甲殻類の中で完全に陸上生活に適応した唯一のグループであり、水中生活を起源とする祖先から、呼吸・繁殖・水分保持の仕組みを大きく変化させてきた。

Organisms commonly known as pillbugs and woodlice belong to the phylum Arthropoda, subphylum Crustacea, order Isopoda, and suborder Oniscidea (terrestrial isopods).
Although often mistaken for insects or myriapods, they are taxonomically closer to crabs and shrimps.
Terrestrial isopods are unique among crustaceans in that they have fully adapted to life on land, evolving specialized mechanisms for respiration, reproduction, and water conservation.

第 2 章　分類と代表的な属・種

日本で一般的に見られるグループは以下の通りである。

- ダンゴムシ類（丸まる）
  - *Armadillidium vulgare*（オカダンゴムシ）
- ワラジムシ類（丸まらない）
  - *Porcellio scaber*（ワラジムシ）
  - *Porcellionides pruinosus* など

「ダンゴムシ」と「ワラジムシ」は学術分類ではなく、行動形態（丸まるか否か）に基づく和名的区分である。

Common terrestrial isopods in Japan include:

- Pillbugs (conglobating forms)
  - *Armadillidium vulgare*
- Woodlice (non-conglobating forms)
  - *Porcellio scaber*
  - *Porcellionides pruinosus*, etc.

The terms “pillbug” and “woodlouse” are not strict taxonomic categories but are based on behavioral and morphological traits, particularly the ability to roll into a ball.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




図 1. アンブロスおよびシウダ・ペガソーオドネルーカンテラ・デル・トラペロ産のワラジムシ類の外部形態（現地で撮影した写真）。

Fig. 1. Habitus of woodlice from Ambroz and Ciudad Pegaso - O'Donnell - Cantera del Trapero (photographs taken in situ).

出典: (PDF) Los isópodos terrestres (Crustacea, Isopoda, Oniscidea) del entorno de las Lagunas de Ambroz: una zona urbana en proceso de renaturalización en el municipio de Madrid (España). Woodlice (Crustacea, Isopoda, Oniscidea) from the environment of the Lagunas de Ambroz: an urban zone in the process of renaturalization in the municipality of Madrid (Spain)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl




図 2. ***Armadillidium cf. arcangelii***：オス（背面）およびメス（側面）。

Fig. 2. Armadillidium cf. arcangelii: male (dorsal) and female (lateral).

出典: (PDF) Los isópodos terrestres (Crustacea, Isopoda, Oniscidea) del entorno de las Lagunas de Ambroz: una zona urbana en proceso de renaturalización en el municipio de Madrid (España). Woodlice (Crustacea, Isopoda, Oniscidea) from the environment of the Lagunas de Ambroz: an urban zone in the process of renaturalization in the municipality of Madrid (Spain)

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

第 3 章　外部形態と防御戦略

体は背腹に扁平で、外骨格に覆われる。体節は以下に分かれる。

- 頭部
- 胸部（7 節、各節に 1 対の脚）
- 腹部（付属肢は呼吸・生殖に関与）

ダンゴムシは**団子化（conglobation）**により捕食者から身を守る。一方、ワラジムシは丸まらず、高速移動・隙間潜行・集団化で防御する。

The body is dorsoventrally flattened and covered with a rigid exoskeleton. It consists of:

- Head
- Thorax (7 segments with one pair of legs each)
- Abdomen (appendages modified for respiration and reproduction)

Pillbugs defend themselves by conglobation (rolling into a ball), whereas woodlice rely on rapid movement, crevice hiding, and aggregation behavior.

第 4 章　呼吸・水分保持の生理学

陸生等脚類は水中呼吸器官である腹肢を改変し、

- 擬気管（pleopodal lungs）
- 湿潤表面でのガス交換

を行う。

乾燥には弱く、湿度依存性が非常に高い。集団で固まる行動は、水分蒸散を抑える生理的適応である。

Terrestrial isopods modified their ancestral aquatic pleopods into pleopodal lungs or gas-exchange surfaces.

They are highly sensitive to desiccation, and aggregation behavior helps reduce water loss under dry conditions.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

第 5 章　食性と生態的役割

主に以下を摂食する。

- 落葉・枯死植物
- 腐植
- 菌類・微生物膜

これにより、**分解促進者（デトリタス食者）**として土壌形成や養分循環に重要な役割を果たす。

They feed mainly on:

- Leaf litter and dead plant material
- Humus
- Fungi and microbial biofilms

Thus, they function as key detritivores, accelerating decomposition and nutrient cycling in terrestrial ecosystems.

第 6 章　生殖・発生

雌は腹部に育房（marsupium）を形成し、卵と幼生を保護する。
発生は直接発生で、孵化後の幼生（manca）は成体に近い形態を持つ。

Females form a marsupium (brood pouch) on the ventral side, where eggs and embryos develop. Development is direct, and hatchlings (mancae) resemble miniature adults.

第 7 章　進化と陸上化

陸生等脚類は海生等脚類から派生し、

- 呼吸器の改変
- 水分保持機構
- 行動的適応

を経て陸上化した。甲殻類の陸上進化のモデル群とされる。

Terrestrial isopods evolved from marine ancestors and represent a classic model of terrestrial adaptation in crustaceans, involving respiratory, physiological, and behavioral innovations.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

第 8 章　人間との関係・研究利用

- 学校教育（走性・行動実験）
- 分解者研究
- 毒性試験・重金属蓄積研究
- 魚類・両生類の餌料・試料候補

飼育が容易で、実験動物としての潜在性が高い。

They are used in:

- Educational experiments
- Decomposition and soil ecology studies
- Toxicology and heavy-metal accumulation research
- Potential feed or experimental organisms for fish and amphibians

Their ease of maintenance makes them valuable model invertebrates.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

参考文献

総説・専門書

- Minelli, A. (Ed.). *The Myriapoda*. Brill
- Schmalfuss, H. (2003). *World Catalog of Terrestrial Isopods*
- Hornung, E. (2011). Evolutionary adaptation of oniscidean isopods

生理・生態

- Zimmer, M. (2002). Nutrition in terrestrial isopods
- Hassall, M. et al. (2018). Aggregation behavior in woodlice

行動・教育利用

- Cloud, J. (2010). *Isopods in the Classroom*
- Warburg, M. R. (1993). *Evolutionary Biology of Land Isopods*

Warburg, M. R. (1993)
Evolutionary Biology of Land Isopods
Springer
→ 陸生等脚類研究の古典的名著。進化・生理・行動を包括。

Schmidt, C. (2008)
Phylogeny of the terrestrial Isopoda (Oniscidea)
Invertebrate Systematics
→ 分子・形態両面からの系統解析。

Schmalfuss, H. (2003)
World Catalog of Terrestrial Isopods (Isopoda: Oniscidea)
Stuttgarter Beiträge zur Naturkunde
→ 種リスト・分類の決定版（分類研究必携）。

Hornung, E. (2011)
Evolutionary adaptation of oniscidean isopods to terrestrial life
ZooKeys
→ 陸上適応のレビューとして非常に重要。

Ⅱ. 生理・水分保持・呼吸

Edney, E. B. (1968)
Water Balance in Land Arthropods
Springer
→ 等脚類の乾燥耐性研究の基盤文献。

Wright, J. C. & Ting, K. (2006)
Respiratory physiology of terrestrial isopods
Comparative Biochemistry and Physiology A

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

Greenaway, P. (1985)
Calcium balance and moulting in terrestrial crustaceans
Biological Reviews

Carefoot, T. H. (1984)
Nutrition and growth of terrestrial isopods
Biological Reviews

Ⅲ. 行動学（走性・集合・学習）

Fraenkel, G. & Gunn, D. L. (1961)
The Orientation of Animals
Dover
→ ダンゴムシ走性実験の理論的基盤。

Hassall, M., Turner, J. G., & Rands, M. R. W. (1987)
Aggregation and dispersal in terrestrial isopods
Symposia of the Zoological Society of London

Cloudsley-Thompson, J. L. (1956)
Studies in diurnal rhythms of woodlice
Journal of Experimental Biology

Ⅳ. 生態・分解・物質循環

Zimmer, M. (2002)
Nutrition in terrestrial isopods (Isopoda: Oniscidea)
Pedobiologia
→ 腐植食・微生物共生の定番レビュー。

Zimmer, M. & Topp, W. (2000)
Species-specific utilization of microbial food
Oecologia

David, J. F. (2014)
The role of terrestrial isopods in litter decomposition
Biological Reviews

Ⅴ. 毒性・環境指標・重金属蓄積

Dallinger, R. (1994)
Invertebrate organisms as biological indicators
Environmental Toxicology and Chemistry

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

Hopkin, S. P. (1990)
Species-specific differences in metal accumulation
Environmental Pollution

Paoletti, M. G. & Hassall, M. (1999)
Woodlice as bioindicators of heavy metals
Agriculture, Ecosystems & Environment

Ⅵ. 発生・繁殖・育房（marsupium）

Suzuki, S. & Futami, K. (1980)
Studies on marsupial development in terrestrial isopods
Journal of Crustacean Biology

Heath, D. J. (1973)
Marsupium formation in Oniscidea
Journal of Zoology

Ⅶ. 遺伝・分子生物学

Michel-Salzat, A. & Bouchon, D. (2000)
Mitochondrial genome variation in Armadillidium
Genetics

Cordaux, R. et al. (2011)
Wolbachia and sex determination in isopods
Trends in Genetics

Ⅷ. 飼育・実験動物・教育利用

Cloud, J. (2010)
Isopods in the Classroom
NSTA Press
→ 教育・実験用途に特化。

Pennak, R. W. (1989)
Fresh-water Invertebrates of the United States
Wiley
→ 餌生物・実験試料文脈で引用されることが多い。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

Ⅸ. 日本語文献

内田亨 監修
『動物系統分類学 7 甲殻類』
中山書店

西村三郎 編
『原色検索 日本海岸動物図鑑』
保育社

日本動物学会 編
『動物学事典』
丸善出版

高校・大学向け教材
文部科学省検定 生物実験書（走性・行動実験）

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！
HAL lab.axolotl

魚類・両生類の餌としての甲殻類のメリット

1. はじめに

魚類および両生類の飼育・研究において、餌料の質は成長、健康状態、生理機能、さらには実験再現性に直接影響を及ぼす重要な要素である。とりわけ成長期個体では、タンパク質、脂質、ミネラル、微量元素のバランスが成長率や生存率を左右することが広く知られている（Lovell, 1989；Davis et al., 1993）。これらの観点から、甲殻類を主成分または補助成分とする餌料は、古くから魚類・両生類の飼育および実験に利用されてきた。

実際、アルテミアやミジンコは仔魚・幼生飼育の標準餌料として確立されており、エビ類やカニ類由来餌料は成長促進や生存率向上を目的として広く用いられている（Sargent et al., 1997）。近年では、陸生等脚類であるワラジムシやダンゴムシも、栄養供給源および実験試料として注目されつつある。

本総説では、アルテミア、ミジンコ、エビ類、カニ類、等脚類（ワラジムシ・ダンゴムシを含む）などの甲殻類餌料に共通する利点を整理し、魚類・両生類における栄養学的・生理学的意義を概説する。

---

2. 高タンパク質・高消化性

甲殻類は一般に高タンパク質であり、必須アミノ酸組成が脊椎動物の要求量とよく一致している。例えば、アルテミアおよびミジンコは、仔魚期の海産魚・淡水魚において筋肉形成と成長を効果的に促進することが報告されている（Conceição et al., 2010）。

両生類においても、エビ類や甲殻類ミールを含む餌料は、ミミズ単独給餌と比較して成長速度が速く、体重増加が顕著であることが示されている（Wright & Whitaker, 2001）。これらの結果は、甲殻類由来タンパク質が高い消化吸収性を有し、摂餌行動を強く誘発することに起因すると考えられる。

---

3. カルシウムおよびミネラル供給源としての役割

甲殻類の最大の特徴の一つは、外骨格に由来する豊富なカルシウム含量である。カルシウムは魚類・両生類において骨格形成、鰭条や四肢の発達、神経・筋機能の維持に不可欠な元素である（Ogino & Takeda, 1978）。とくに両生類では、餌由来カルシウムの摂取量が成長率や骨格健全性に強く影響することが知られている。メキシコサラマンダー（*Ambystoma mexicanum*）では、カルシウム強化餌の投与により体長および体重増加が有意に促進され、骨密度の上昇や骨格異常の減少が確認されている（Wright & Whitaker, 2001）。さらに、無機カルシウム添加に代えて、エビ類や甲殻類ミールを含む餌料を与えた場合にも、同様の成長促進効果が得られることが報告されている。同様の傾向は、イベリアトゲイモリ（*Pleurodeles waltl*）やアフリカツメガエル（*Xenopus laevis*）においても観察されており、甲殻類餌料の摂取は四肢骨形成の健全化に寄与することが示唆されている（Browne et al., 2003）。

---

4. 脂質・必須脂肪酸の供給

甲殻類は、DHA や EPA などの**高度不飽和脂肪酸（HUFA）**を含むことが多く、これらは魚類の脳および視覚系の発達に不可欠である（Sargent et al., 1997）。アルテミアを脂肪酸強化した餌料は、仔魚の生残率および成長率を大きく改善することが知られている。

両生類においても、脂質を含む甲殻類餌料は細胞膜形成や免疫機能の維持に寄与し、幼生期の健全な発達を支える重要な要因であると考えられている。

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

5. 行動・摂餌刺激効果

生きた甲殻類餌料は、動きや化学刺激によって摂餌行動を強く刺激する。魚類仔魚では、ミジンコやアルテミアの遊泳運動が捕食行動を誘発し、摂餌開始率を向上させることが報告されている（Kolkovski, 2001）。両生類においても、動く甲殻類餌料は拒食傾向の個体に対して有効であり、人工飼料への移行前段階として重要な役割を果たす。

---

6. 実験・研究用途における利点

甲殻類はサイズや栄養状態を比較的容易に制御できるため、

- **成長・栄養試験**
- **ミネラル代謝研究**
- **毒性・環境影響評価**

といった研究において再現性の高い試料餌料として利用可能である（Davis et al., 1993）。とくに陸生等脚類（ワラジムシ・ダンゴムシ）は、低コストで大量飼育が可能であり、カルシウム代謝や微量元素蓄積の研究材料としての応用が期待されている。

---

7. おわりに

以上より、甲殻類は高タンパク質、豊富なカルシウムおよび脂肪酸供給、摂餌刺激効果といった複数の利点を併せ持つ餌料生物であり、魚類・両生類の飼育および研究において極めて有用である。今後は、種ごとの栄養組成の詳細解析や、成長・生理機能への定量的影響評価が進むことで、より最適化された甲殻類餌料の利用が期待される。

参考文献 / References

- Browne, R. K., et al. (2003). Amphibian nutrition and captive husbandry. *Herpetological Journal*.
- Conceição, L. E. C., et al. (2010). Amino acid requirements of fish larvae. *Aquaculture*.
- Davis, D. A., et al. (1993). Calcium and phosphorus nutrition in fish. *Journal of Nutrition*.
- Kolkovski, S. (2001). Digestive enzymes in fish larvae and live feed. *Aquaculture*.
- Lovell, R. T. (1989). *Nutrition and Feeding of Fish*. Van Nostrand Reinhold.
- Ogino, C., & Takeda, H. (1978). Mineral requirements in fish. *Bulletin of the Japanese Society of Scientific Fisheries*.
- Sargent, J., et al. (1997). Essential fatty acids in fish nutrition. *Aquaculture*.
- Wright, K. M., & Whitaker, B. R. (2001). *Amphibian Medicine and Captive Husbandry*. Krieger Publishing.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

Advantages of Crustaceans as Feed for Fish and Amphibians

1. Introduction

In the husbandry and experimental research of fish and amphibians, diet quality is a critical factor that directly influences growth, health status, physiological function, and experimental reproducibility. In particular, during growth stages, the balance of proteins, lipids, minerals, and trace elements strongly affects growth rate and survival (Lovell, 1989; Davis et al., 1993). From this perspective, crustaceans used as primary or supplemental feed organisms have long played an important role in the culture and experimental use of fish and amphibians.Artemia and Daphnia are well established as standard live feeds for larval fish and amphibians, while shrimp- and crab-derived feeds are widely used to improve growth performance and survival (Sargent et al., 1997). In recent years, terrestrial isopods such as woodlice and pill bugs have also attracted attention as novel nutritional resources and experimental feed organisms.

This review summarizes the common advantages of crustacean-based feeds—including Artemia, Daphnia, shrimp, crabs, and terrestrial isopods (woodlice and pill bugs)—and discusses their nutritional and physiological significance in fish and amphibians.

2. High Protein Content and Digestibility

Crustaceans are generally characterized by **high protein content**, and their essential amino acid profiles closely match the nutritional requirements of vertebrates. For example, Artemia and Daphnia have been shown to effectively promote muscle development and growth in larval stages of both marine and freshwater fish (Conceição et al., 2010).In amphibians, diets containing shrimp or crustacean meal result in faster growth rates and greater body mass gain compared with earthworm-only diets (Wright & Whitaker, 2001). These findings suggest that crustacean-derived proteins exhibit high digestibility and bioavailability, while also effectively stimulating feeding behavior compared with formulated diets.

3. Calcium and Mineral Supply

One of the most distinctive characteristics of crustaceans is their high calcium content derived from the exoskeleton. Calcium is essential for skeletal formation, development of fin rays and limbs, and maintenance of neuromuscular function in both fish and amphibians (Ogino & Takeda, 1978).In amphibians, dietary calcium intake is known to have a particularly strong influence on growth rate and skeletal integrity. In the Mexican salamander (*Ambystoma mexicanum*), supplementation with calcium-enriched diets significantly enhances body length and body mass, increases bone mineral density, and reduces the incidence of skeletal deformities (Wright & Whitaker, 2001). Importantly, similar growth-promoting effects have been observed when calcium supplementation is replaced by crustacean-based feeds containing shrimp or crustacean meal.

Comparable results have been reported in other amphibians, including the Iberian ribbed newt (*Pleurodeles waltl*) and the African clawed frog (*Xenopus laevis*), where crustacean-rich diets contribute to improved limb bone development and overall skeletal health (Browne et al., 2003). These findings collectively suggest that crustacean-based feeds provide a biologically effective source of calcium and minerals for amphibians.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

4. Lipids and Essential Fatty Acids

Live crustacean feeds strongly **stimulate feeding behavior** through movement and chemical cues. In fish larvae, the swimming behavior of Artemia and Daphnia effectively triggers prey capture responses and increases feeding initiation rates (Kolkovski, 2001).Similarly, in amphibians, actively moving crustacean prey is effective in overcoming feeding reluctance, especially during early husbandry stages. Consequently, crustacean feeds are often used as an intermediate step before transitioning to formulated diets.

5. Stimulation of Feeding Behavior

Live crustacean feeds strongly **stimulate feeding behavior** through movement and chemical cues. In fish larvae, the swimming behavior of Artemia and Daphnia effectively triggers prey capture responses and increases feeding initiation rates (Kolkovski, 2001).

Similarly, in amphibians, actively moving crustacean prey is effective in overcoming feeding reluctance, especially during early husbandry stages. Consequently, crustacean feeds are often used as an intermediate step before transitioning to formulated diets.

6. Advantages for Experimental Applications

Because crustaceans can be cultured under controlled conditions, their size and nutritional status can be relatively easily standardized. As a result, they serve as **highly reproducible experimental feed organisms** in studies involving:

- **growth and nutritional evaluation,**
- **mineral metabolism, and**
- **toxicological and environmental impact assessment (Davis et al., 1993).**

In particular, terrestrial isopods such as woodlice and pill bugs are inexpensive, easy to mass-culture, and rich in calcium, making them promising experimental models for studies on calcium metabolism and trace element accumulation.

7. Conclusion

In summary, crustaceans offer multiple advantages as feed organisms for fish and amphibians, including high protein content, abundant calcium and essential fatty acids, and strong feeding stimulation. These characteristics make crustaceans valuable for both husbandry and experimental research. Future studies focusing on detailed nutritional profiling and quantitative assessments of physiological effects will further optimize the use of crustacean-based feeds.

参考文献 / References

- Browne, R. K., et al. (2003). Amphibian nutrition and captive husbandry. *Herpetological Journal*.
- Conceição, L. E. C., et al. (2010). Amino acid requirements of fish larvae. *Aquaculture*.
- Davis, D. A., et al. (1993). Calcium and phosphorus nutrition in fish. *Journal of Nutrition*.
- Kolkovski, S. (2001). Digestive enzymes in fish larvae and live feed. *Aquaculture*.
- Lovell, R. T. (1989). *Nutrition and Feeding of Fish*. Van Nostrand Reinhold.
- Ogino, C., & Takeda, H. (1978). Mineral requirements in fish. *Bulletin of the Japanese Society of Scientific Fisheries*.
- Sargent, J., et al. (1997). Essential fatty acids in fish nutrition. *Aquaculture*.
- Wright, K. M., & Whitaker, B. R. (2001). *Amphibian Medicine and Captive Husbandry*. Krieger Publishing.

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

餌料別・成分比較表（乾物 1 g あたりの代表値）

※　以下、乾物換算（dry matter, DM）で表示

単位

- Ca，ミネラル：**mg/g DM**
- タンパク質・脂質・糖質：**mg/g DM**
- ビタミン：**μg/g DM（代表的なもの）**
- 「—」＝報告例が乏しい／無視できる量

灰分（Ash）とは？

- 無機成分の総量（Ca, P, Mg, Na, K，微量元素を含む）
- **骨・外骨格形成に直結**
- 甲殻類で特に高い
- 灰分：**mg/g DM**

餌料別：灰分量と Ca:P 比

- Ca:P：**重量比**

※ 理想的な Ca:P 比

・両生類・爬虫類：1.5–2 : 1 以上

・成長期・再生期（アホロートル）：2 : 1 以上が望ましい

※ **HUFA（高度不飽和脂肪酸）**：炭素鎖中に複数の二重結合を含む脂肪酸の一種

カルシウム含有量順（高 → 低）

ダンゴムシ
＞ ワラジムシ
＞ ひかりウーパー＋Ca 粉末
＞ オキアミ
＞ ひかりウーパー（通常）
＞ アルテミア
＞ フタホシコオロギ
＞ イエコオロギ
＞ ショウジョウバエ（成虫）
＞ ショウジョウバエ（幼虫）
＞ アカムシ

表 1: 甲殻類飼料が含有する栄養成分比較

① 通常餌

餌料	Ca	タンパク質	脂質	糖質	灰分	備考
ひかりウーパー	12–18	450–500	80–100	120–150	90–110	人工飼料
ひかりウーパー＋Ca粉末	30–60	450–500	80–100	120–150	120–180	Ca添加量依存

② 甲殻類

餌料	Ca	タンパク質	脂質	糖質	灰分	特徴
ダンゴムシ	70–120	350–420	60–80	40–60	200–250	外骨格Ca非常に多い
ワラジムシ	50–90	330–400	50–70	50–70	180–220	ダンゴムシより低Ca
アルテミア	5–15	500–600	80–120	50–80	70–90	HUFA強化可
オキアミ	20–40	600–700	100–150	30–50	120–150	Ca＋EPA/DHA

③ 昆虫類

餌料	Ca	タンパク質	脂質	糖質	灰分	Ca:P
アカムシ	1–3	550–650	50–80	80–120	40–60	極端に低Ca
ショウジョウバエ（成虫）	2–5	450–520	120–150	100–130	50–70	高脂質
ショウジョウバエ（幼虫）	1–3	400–480	180–220	80–110	40–60	Ca欠乏
イエコオロギ	2–6	600–650	120–150	50–80	60–80	Ca:P不良
フタホシコオロギ	2–7	580–630	130–160	40–70	60–90	同上

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

ハルラボにおける知見

アホロートルに甲殻類や昆虫類を餌として与え、生育率や生存率の違いを比較した。検討の結果、ダンゴムシだけに突出した効果が見られた訳ではなかったが、成長率の顕著な増大が確認された。これは、甲殻類を餌として摂取することで、必要量の Ca を効率よく摂取することが可能となったことに起因することが示唆された。実際に野生個体も、積極的に甲殻類や昆虫類を摂取することで、効率的に生育率を向上させ、最大体調を 40.0cm 近くにまで大きくできた可能性が考えられた。一方で、彼らが保持している腸内細菌の影響が炎症を誘導することから、継続的に与え続けると寿命の短縮につながってしまう場合がある点には注意したい。

Findings from Our Lab.

The axolotls were fed crustaceans and insects, and differences in growth rate and survival were compared. Although pill bugs did not show a uniquely outstanding effect compared to other prey, a marked increase in growth rate was observed overall. This suggests that consuming crustaceans as feed enables efficient intake of the required amount of calcium (Ca). Indeed, wild individuals may actively consume crustaceans and insects to enhance growth efficiency, potentially reaching a maximum body length of nearly 40.0 cm. On the other hand, it should be noted that the gut microbiota they harbor can induce inflammation; therefore, continuous feeding of such diets may lead to a shortened lifespan.




図 1. 甲殻類を摂取したアホロートルの成長率・生存率

※各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




図 2. 甲殻類を摂取したアホロートルにおける miRNA 発現動態-1

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




図 3. 甲殻類を摂取したアホロートルにおける miRNA 発現動態-2

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




図 4. 甲殻類を摂取したアホロートルにおける miRNA 発現動態-3

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。




図 5. 甲殻類を摂取したアホロートルにおける内在性配列の発現動態

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。
はるらぼ！ HAL lab.axolotl

参考文献

[1] Lovell, R. T., 1989, Nutrition and Feeding of Fish, Van Nostrand Reinhold, —, URL: https://books.google.com/books?id=Z0cFAQAAIAAJ

[2] Davis, D. A., Gatlin, D. M., et al., 1993, Journal of Nutrition, 123: 124–131, DOI:10.1093/jn/123.suppl_2.124

[3] Ogino, C., Takeda, H., 1978, Bulletin of the Japanese Society of Scientific Fisheries, 44: 101–106, DOI:10.2331/suisan.44.101

[4] Sargent, J., McEvoy, L., Estevez, A., et al., 1997, Aquaculture, 155: 117–127, DOI:10.1016/S0044-8486(97)00123-4

[5] Conceição, L. E. C., Yúfera, M., Makridis, P., et al., 2010, Aquaculture, 304: 1–13, DOI:10.1016/j.aquaculture.2010.03.016

[6] Kolkovski, S., 2001, Aquaculture, 200: 181–201, DOI:10.1016/S0044-8486(01)00700-3

[7] Wright, K. M., Whitaker, B. R., 2001, Amphibian Medicine and Captive Husbandry, Krieger Publishing, —, URL: https://books.google.com/books?id=9zYzAQAAMAAJ

[8] Browne, R. K., et al., 2003, Herpetological Journal, 13: 1–10, URL: https://www.thebhs.org/publications/the-herpetological-journal/volume-13-number-1

[9] Kohl, K. D., Carey, H. V., 2016, Journal of Experimental Biology, 219: 2019–2031, DOI:10.1242/jeb.136143

[10] Warne, R. W., Kirschman, L., Zeglin, L., 2017, Frontiers in Microbiology, 8: 2478, DOI:10.3389/fmicb.2017.02478

[11] Zimmer, M., 2002, Pedobiologia, 46: 589–607, DOI:10.1078/0031-4056-00158

[12] Carefoot, T. H., 1984, Biological Reviews, 59: 1–30, DOI:10.1111/j.1469-185X.1984.tb00404.x

[13] Hornung, E., 2011, ZooKeys, 176: 17–43, DOI:10.3897/zookeys.176.1929

[14] Schmalfuss, H., 2003, Stuttgarter Beiträge zur Naturkunde, URL:https://www.naturkundemuseumbw.de/sites/default/files/forschung/publikationen/beitraege_naturkunde/serie_a/640.pdf

[15] 荻野珍吉，武田博, 1978，日本水産学会誌, 44: 101–106, DOI:10.2331/suisan.44.101

[16] 西村三郎 編, 1992，原色検索 日本海岸動物図鑑，保育社, URL: https://www.hoikusha.co.jp

[17] 内田亨 監修, 1985，動物系統分類学 第 7 巻 甲殻類，中山書店, —, URL: https://www.nakayamashoten.co.jp

[18] 日本動物学会 編, 2018，動物学事典，丸善出版, URL: https://www.maruzen-publishing.co.jp

[19] 文部科学省, 2019，高等学校生物実験書, —, URL: https://www.mext.go.jp

[20] 松井正文, 2005，両生類学，東京大学出版会, —, URL: https://www.utp.or.jp

※ 各検証については、内容と必要に応じて、専門家の方々のご指導・ご協力のもと実施しております。
※一般家庭で実施不可能な試験が複数あり。一般的な飼育技術用へ応用可能な場合は、その旨を明記いたします。

はるらぼ！ HAL lab.axolotl

奥付

ウーパールーパーだより vol 8.0

発行日・・・ 2026/4/19

編集・著者　杉山 遥 (PN; Ph.D) ほか

発行元　ウーパールーパー研究室_はるらぼ, HAL_Lab_Axolotl

問い合わせ先　sugiyama.haruka.axoltl@gmail.com または X (Twitter) の DM へ

X (旧 Twitter) URL　https://twitter.com/lab_new2 (@lab_new2)



CC BY NC ND

これからもウパ活は継続しますよ。

by　杉山 遥



ウーパールーパー研究室・はるらぼ

sugiyama.haruka.axoltl@gmail.com

since 2020～（@lab_new2）

X (旧 Twitter): https://twitter.com/lab_new2

※注意

本誌の許可なき改変・商業利用は、

我々は一切認めておりません。